Patient Education



Anal Fissure

Thank you for allowing us to care for you. We recommend the following treatment for you.

Regulate bowel movements to make them soft, well-formed, easily passable/evacuable without straining. The stools should not be hard or liquid.

Below are some suggestions on things you can do to help regulate your stool:

- 1. Eat a high fiber diet (fresh fruits, vegetables, whole grains, bran, etc.)
- 2. Take Fiber supplements (MetamucilTM, KonsylTM, BenefiberTM, FiberConTM, or Fiber GummiesTM) twice to three times daily, with a goal of 20-30 grams of soluble fiber daily.
- 3. Drink 8-10 (8oz) glasses of water daily

Acute management:

- 1. Diltiazem (or other topical ointment) applied with a gloved finger to the anus (do not go inside) 4-6 times daily
- 2. Continue this until instructed to stop

Avoid irritants such as:

- 1. Caffeinated, alcoholic, citrus, spicy, or acidic foods or drinks
- 2. Perfumes and/or dyes
- 3. Excess wiping (1-2 times only)
- 4. Rubbing or scratching
- 5. Scented soaps or scented baths
- 6. Scented or dyed/colored toilet paper/tissue

Perianal Care:

- 1. Wipe with witch hazel/aloe wipes (1-2 wipes is sufficient)
- 2. Dry the perianal area with a hair dryer on low heat after showering.
- 3. Avoid bathing.
- 4. Apply Zinc Oxide, Calmoseptine, Balneol, or another diaper rash ointment (i.e. Boudreaux's or Triple Paste) to the perianal area circumferentially (but not in anal canal) twice a day.
- 5. Place cotton or cotton ball near the perianal area to absorb moisture or discharge.

For discomfort:

- 1. Sitz bath (or bidet, or hand-held shower irrigation with warm water) sparingly, up to twice a day for a maximum of 10-15 minutes at a time.
- 2. Can add Epsom salt to your sitz bath if desired
- You may take a non-aspirin pain reliever such as TylenolTM, or GenapapTM as needed. Please follow dosing recommendations on bottle. *Some other common pain relievers can affect blood thinning*. Examples include:
 - Aspirin
 - Ibuprofen (AdvilTM, MotrinTM)
 - Naproxen (AleveTM, NaprosynTM)

Please talk with your health care provider before taking these pain relievers