

Okyspor TT



Inoculum of mycorrhizal fungi with Trichoderma



COMPOSITION

Content in Mycorrhizae

Enzymes

Fungi and bacteria of the rhizosphere

Trichoderma harzianum and **Trichoderma gamsii**



Appearance
Powder



Application
Foliar / Radical



Packaging
1 - 5 Kg



DESCRIPTION

Okyspor TT is an organic product based on Mycorrhizae of the genus *Glomus* associated with rhizosphere fungi and bacteria including *Trichoderma harzianum* and *Trichoderma gamsii* in prevalence.

Thanks to the richness of its components and the ready activation of its consortium, Okyspor TT allows to achieve a great effectiveness, immediately after its application, both on the ground and on vegetation. Okyspor TT can be used also for the preventive treatment of seeds and seedlings in nurseries and open field.

Okyspor TT enhances plant defence mechanisms, then stimulates the development and vigor of vegetation.

Trichoderma is naturally characterized by the ability to produce natural antibiotics, which active to control a high number of pathogenic fungi.

Thanks to its organic characteristics, Okyspor TT can be used in an integrated or organic defense strategy to reduce the chemical pressure on the environment and to optimize the level of residues on the crop.

Furthermore, the particular multi-site action mechanism facilitates the management of anti-resistance strategies.

Okyspor TT enhances flowering and fruiting, optimizing fertilizers use, especially Phosphorus.



DOSES AND METHOD OF USE

Crops	Doses	Application	
Short cycle crops (30 - 40 days)	1 - 2 Kg/ha	First	0,5 - 1 Kg of Okyspor TT at sowing or transplanting
		Second	0,5 - 1 Kg of Okyspor TT after 10/15 days
Long cycle crops (60 - 90 days)	1 - 3 Kg/ha	First	0,5 - 1 Kg of Okyspor TT at sowing or transplanting
		Second	1 kg di Okyspor TT after 10/15 days
		Third	0,5 - 1 kg di Okyspor TT after 40/50 days