



SUPPLEMENT

FOR ANIMAL USE ONLY

Dexiron 200

Reg. No. G2813 (Act 36/1947)

Injection

For the prevention and treatment of iron deficiency anaemia in piglets and calves

■ STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.
Protect from light.

■ COMPOSITION

Iron dextran complex 20 % m/v

■ WARNINGS

- For administration by intramuscular injection only. NOT for administration by intravenous or subcutaneous injection.
- Calves and piglets with a vitamin E deficiency and/or selenium deficiency should receive an injection with vitamin E and/or selenium prior to the administration of iron.
- DEXIRON 200 is contra-indicated in animals with severe liver damage or an acute kidney infection.
- Iron preparations should not be administered concomitantly with any of the tetracycline group of antibiotics.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ PRECAUTIONS

- Sows to be fed a selenium enriched diet prior to farrowing.
- Observe aseptic precautions. Ensure that needles and syringes are clean and sterile prior to use.
- Weeping from the injection site can be avoided by pushing the skin and underlying fat to one side with the thumb before inserting needle, and allowing a second or so to elapse before withdrawing the needle.

■ DIRECTIONS FOR USE

Use only as directed

Piglets

1 ml solution per animal (1,5 – 2,0 kg body mass) administered by intramuscular injection in the neck, just below the ear, on the third day of life.

Calves

2 – 4 ml solution per animal administered by intramuscular injection, in the first week of life.

■ PRESENTATION

100 ml glass vial

VIRBAC RSA (Pty) Ltd
(Reg. No. 1990/003743/07)
Private Bag X115
Halfway House 1685
South Africa
Tel: +27 12 657 6000

Namibia Reg. No. V02/19.3/10 