

ENDOPARASITICIDE FOR ANIMAL USE ONLY

Prodose<sup>®</sup> Blue

Reg. No. G2090 (Act 36/1947)

#### ROUNDWORM AND MILK TAPEWORM REMEDY FOR CATTLE

(2)<sup>†</sup>

CAUTION

#### STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

# COMPOSITION

## WARNINGS

- Do not slaughter animals for human consumption within 12 days of dosing.
- Do not dose lactating cows whose milk or milk products are intended for human consumption.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Management practice should ensure that drenching procedures are carefully carried out. It is important not to exceed the recommended dosage in cows during the first month of pregnancy.

## ■ DIRECTIONS FOR USE - Use only as directed Shake well before use.

Dosage: The dose is 1 ml per 10 kg body mass, for example:

Body mass	Body mass
150 kg	15 ml
200 kg	20 ml
250 kg	25 ml
300 kg	30 ml

## Efficacy: Roundworms: PRODOSE BLUE controls the following:

Worm species	Immatures	Adult
Wireworm	*	*
Brown stomachworm	*	*
Cattle bankruptworm	*	*
Hookworm	*	*
Nodular worm	*	*

Tapeworms: Milk tapeworm (*Moniezia* spp.): Class 1 (100 % effective in 80 % of the treated animals).

Ovicidal: Roundworm eggs present in the animal at the time of dosing are prevented from hatching.

## PRESENTATION

200 ml, 1L, 5L and 10L polypropylene bottles.

## REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd (Reg No. 1990/003743/07) Private Bag X115 Halfway House 1685 South Africa Tel: +27 12 657 6000