

**ENDOPARASITICIDE  
FOR ANIMAL USE ONLY**

②+④†

## Prodose® Orange

Reg. No. G2101 (Act 36/1947)

**NASAL BOT, LIVER FLUKE, ROUNDWORM AND MILK TAPEWORM  
REMEDY FOR SHEEP AND GOATS WITH RESIDUAL PROTECTION  
AGAINST RE-INFESTATION OF WIREWORM AND HOOKWORM.**

### CAUTION

#### ■ STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

#### ■ COMPOSITION

Albendazole ..... 1,90 % m/v  
Closantel (as sodium) ..... 3,94 % m/v

#### ■ WARNINGS

- Do not use in milk goats whose milk is intended for human consumption.
- Allow 30 days between last dosing and slaughter for human consumption.
- Handle with care.
- Management practices should always ensure that all drenching procedures are carefully carried out. It is particularly important not to exceed the recommended dosage in ewes during mating or during the first month of pregnancy.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

#### ■ PRECAUTIONS

- Do not repeat treatments more frequently than every 5 weeks.
- As with all worm remedies it is advisable to ensure animals have been immunised against pulpy kidney disease before drenching.
- Prevent contamination of food, food utensils, etc.
- Wash hands thoroughly after use.
- Do not mix with other products.**

#### ■ DIRECTIONS FOR USE - Use only as directed

Residues of the compound are blood-bound. Meat and organs are virtually free from residues and are therefore suitable for human consumption 30 days after treatment. Shake before use. For oral use only

Body mass	Dosage volume
10 kg	2 ml
20 kg	4 ml
30 kg	6 ml
40 kg	8 ml
50 kg	10 ml
60 kg	12 ml

#### Efficacy:

**1. Nasal Worm:** Controls all stages (1<sup>st</sup> 2<sup>nd</sup> and 3<sup>rd</sup> instar larvae).

**2. Roundworms:** PRODOSE ORANGE controls the following:

Worm species	Immatures	Adult	† Presistency
Wireworm	*	*	5 weeks
Brown stomachworm	*	*	-
Bankruptworm	*	*	-
Long-necked bankruptworm	*	*	-
Hookworm	*	*	2 weeks
Nodular worm	*	*	-
Large-mouthed bowelworm	*	*	-

Definition: \* (Control (>90% effective))

† Interval between treatment and re-infestation.

**3. Liver fluke:** PRODOSE ORANGE controls the following:

Live fluke	6 weeks	Adult
	**	*
Giant liver fluke	8 weeks	Adult
	**	*

Definition: \* Control (>90% effective)

\*\* Aids in Control (60 - 89 % effective)

**Tapeworm:** Milk tapeworm: Class 1 (100 % effective in 80 % of the treated animals).

**Ovicidal:** Roundworm eggs present in the animal at the time of dosing are prevented from hatching.

#### ■ PRESENTATION

250 ml, 500 ml, 1 L, 5 L, 10 L and 25 L polypropylene bottles.

#### ■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd  
(Reg No. 1990/003743/07)  
Private Bag X115  
Halfway House  
1685  
South Africa  
Tel: +27 12 657 6000