

Virbac

ECTOPARASITICIDE

FOR EXTERNAL ANIMAL USE ONLY

Amipor®

Reg. No. G2058 (Act 36/1947)

POUR-ON REMEDY FOR CATTLE, SHEEP, GOATS AND GAME

Controls ticks, house flies, stable flies and face flies on cattle.

Kills biting and sucking lice on cattle.

Controls Karoo paralysis ticks on sheep.

Kills red lice on sheep and goats.

Controls ticks on game.

OXPECKER COMPATIBLE

CAUTION

■ STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

■ COMPOSITION

Amitraz 1 % m/v

Cypermethrin 1 % m/v

Piperonyl Butoxide 5 % m/v

■ WARNINGS

- For external use only.
- Animals must not be slaughtered within 18 days of the last treatment. No withdrawal period for milk usage.
- **Do not use on horses: toxic to horses.**
- Handle with care: poisonous when swallowed.
- Calves under 6 months of age must be treated strictly according to body mass.
- Signs of salivation and lethargy may occur with overdosage. These signs are transient and will pass with no ill effect.
- Store in a safe place away from food and feed.
- Toxic to bees, other insects and fish.
- Keep out of reach of children, uninformed persons and animals.

- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ PRECAUTIONS

- Do not mix AMIPOR with any other product.
- Individual animals may show signs of salivation; this is transient and will pass with no ill effect.
- Avoid skin and eye contact. Wash hands immediately with soap and water after accidental contact. Contact with the product may cause a burning sensation of the skin and eyes which may last for 24 hours. This reaction is transient and without harmful effect.
- A mild skin sensitivity may occur in certain individual animals.
- Wash hands thoroughly before eating, drinking or smoking.
- Prevent contamination of food, eating utensils, drinking water, rivers, dams, grazing, etc.
- Destroy empty containers after use. NEVER re-use containers for any other purpose.

NOTE TO PRACTITIONER

In case of accidental poisoning or ingestion, consult a medical practitioner and make this label available to him/her. No specific antidote; treat symptomatically.

■ DIRECTIONS FOR USE

Use only as directed

For external use only

Cattle

For the control of ticks, house flies, stable flies and face flies. Kills biting and sucking lice.

Game

For the control of ticks.

Cattle and Game

Use 10 ml per 100 kg body mass

Application table

Body mass (kg)	Dose (ml)	Doses per 200 ml
50	5	40
100	10	20
200	20	10
300	30	6,5
400	40	5
500	50	4

Apply 10 ml per 100 kg body mass to back midline of cattle/game being treated. Application should be done by means of a suitable applicator from the head to the base of the tail. Treat weekly for three weeks; thereafter, treat according to challenge.

Sheep and goats

Controls Karoo paralysis ticks on sheep.

Use 2 ml per 10 kg body mass.

For Karoo paralysis tick apply approximately a quarter of the dose in each of the four 'ampits' (the axillae and the groin).

Kills red lice in sheep and goats. Use 4 ml per 10 kg body mass. Apply the full dose as a stripe down the midline of the back. Treat sheep/goats twice, 3 weeks apart.

■ PRESENTATION

Polypropylene bottles containing 200 ml, 1 L, 5 L and 20 L solution

■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd

(Reg No. 1990/003743/07)

Private Bag X115

Halfway House 1685

South Africa

Tel: +27 12 657 6000

Namibia Reg. No. V06/19.3.9/75

NSO

Botswana Reg. No. W130656