

## VOERMOL SUPERFOS

Class: Phosphate, Salt and Trace Mineral Supplement for Ruminants

**VOERMOL SUPERFOS** is a molasses-based mineral supplement for cattle and sheep on green veld and pastures. It can also be used as a transitional lick from green to dry pastures:

- is ready mixed and ready to use;
- contains a wide range of minerals and trace minerals which promote reproduction and growth;
- part of the trace minerals are supplied in organic form;
- contains relatively little salt and will therefore not result in excessive salt intake;
- is molasses based and will not blow away;
- is suitable for ruminant game.



### COMPOSITION OF VOERMOL SUPERFOS

REG NO V17422 (Act 36/1947)

		(g/kg)			(mg/kg)
Crude Protein*	(min)	150	Manganese		1 200
Urea	(max)	47.4	Copper		315
Moisture	(max)	120	Cobalt		11
Calcium	(min/max)	60/100	Iron		750
Phosphorus	(min)	50	Iodine		25
			Zinc		1 200
			Selenium		10
*90.7 % derived from NPN					

Packaging – 50 kg

See urea warning.

**RECOMMENDED INTAKE** (g/animal/day)

Cattle: 120 – 240

Sheep: 20 – 40

**OTHER:** Use **Voermol Superfos** as an ingredient in different feed mixtures to supplement phosphorus and trace-minerals.



**VOERMOL**

## UREA WARNING

### UREA

The majority of licks contain urea that can be dangerous under certain conditions. During the formulation and manufacturing process of **Voermol** licks, great care is taken to minimise the risks of feeding urea.

The possibilities of urea poisoning can be further reduced by taking note of the following:

- use licks as prescribed;
- limit lick intake to the prescribed maximum levels;
- feed the licks together with sufficient grazing and/or roughage;
- take special care during feed scarcity (drought conditions) to prevent high lick intake;
- prevent excessive lick intake in the initial stages (salt hunger) by allowing the animals access to **Voermol Rumevite 6P** for the first 14 days;
- do not let licks get wet.

**See urea warning on the back of the bag.**

### UREA WARNING

This farm feed contains NPN (Non-Protein Nitrogen) sources and must therefore be fed strictly according to instructions for use.

- Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf/sheep or 2 – 4 bottles per head of cattle (1 bottle = 750 ml).
- Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
- Do not feed this farm feed indiscriminately with other NPN containing farm feeds. Consult an animal scientist.
- Mix concentrate thoroughly with the prescribed ingredients.
- This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
- Keep lick troughs filled and prevent gluttonous eating by hungry animals. A constant daily intake can help prevent poisoning.
- Before feeding a NPN containing lick, feed an ordinary salt / phosphate lick for at least 7 days.

### GMO STATEMENT

These animal feeds may contain ingredients derived from GMO (genetically modified organisms) origin.



**VOERMOL**