

PRODUCTION SUPPLEMENTS

VOERMOL PRODUCTION LICK

Class: Protein, Mineral and Energy Supplement for Ruminants

VOERMOL PRODUCTION LICK is a complete, ready-mixed lick that provides protein, energy, minerals, vitamins and trace minerals for the optimum production of the grazing animal:

- contains grains and molasses as energy sources;
- helps to prevent excessive mass loss in cows after calving;
- helps to prevent pregnancy toxaemia in ewes during late gestation and can also be used for flush feeding of ewes;
- ensures weight gain of young growing animals on good quality veld;
- part of the trace minerals are supplied in organic form which is highly absorbable and promotes optimal reproduction and growth;
- contains high levels of vitamin A, D3 and E. Vitamin E in combination with selenium is a powerful anti-oxidant which strengthens the animal's immunity. Vitamin D3 helps with the absorption of phosphorus and calcium;
- contains medicaments which promote energy usage from pastures.



PRODUCTION LICK

V10108

COMPOSITION OF VOERMOL PRODUCTION LICK

REG NO V 10108 (Act 36/1947)

(g/kg)

(mg/kg)

Crude Protein* (min)	250	Manganese	300
Urea (max)	55	Copper	50
Crude Fibre (max)	80	Cobalt	0.98
Moisture (max)	160	Iron	150
Calcium (max)	30	Iodine	4
Phosphorus (min)	8	Zinc	290
Potassium (min)	18	Selenium	0.9
Magnesium (min)	5	Vitamin A (IU/kg)	10 000
Sulphur (min)	4	Vitamin D3 (IU/kg)	2 000
		Vitamin E (mg/kg)	50
		Energy est. (MJ ME/kg)	8
*67.4 % derived from NPN		Contains medicaments	

Packaging – 50 kg

See urea warning.

FEEDING INSTRUCTIONS

Use **Voermol Production Lick** as a protein and energy supplement to hay and/or veld.

RECOMMENDED INTAKE	(g/animal/day)
Cattle:	1 500 – 1 620*
Sheep:	250 – 270*

*Maximum allowable intake of lick.

NB. Contains medicaments and must therefore not be fed to horses.



VOERMOL

UREA WARNING

UREA

The majority of licks contain urea that can be dangerous under certain conditions. During the formulation and manufacturing process of **Voermol** licks, great care is taken to minimise the risks of feeding urea.

The possibilities of urea poisoning can be further reduced by taking note of the following:

- use licks as prescribed;
- limit lick intake to the prescribed maximum levels;
- feed the licks together with sufficient grazing and/or roughage;
- take special care during feed scarcity (drought conditions) to prevent high lick intake;
- prevent excessive lick intake in the initial stages (salt hunger) by allowing the animals access to **Voermol Rumevite 6P** for the first 14 days;
- do not let licks get wet.

See urea warning on the back of the bag.

UREA WARNING

This farm feed contains NPN (Non-Protein Nitrogen) sources and must therefore be fed strictly according to instructions for use.

- Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf/sheep or 2 – 4 bottles per head of cattle (1 bottle = 750 ml).
- Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
- Do not feed this farm feed indiscriminately with other NPN containing farm feeds. Consult an animal scientist.
- Mix concentrate thoroughly with the prescribed ingredients.
- This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
- Keep lick troughs filled and prevent gluttonous eating by hungry animals. A constant daily intake can help prevent poisoning.
- Before feeding a NPN containing lick, feed an ordinary salt / phosphate lick for at least 7 days.

GMO STATEMENT

These animal feeds may contain ingredients derived from GMO (genetically modified organisms) origin.



VOERMOL