

RUMEVITE CATTLE BLOCK

Class: Protein and Mineral Supplement for Cattle



**RUMEVITE
CATTLE BLOCK**

**V10932
N-FF 0627**

RUMEVITE CATTLE BLOCK is the easiest way of feeding maintenance supplements to all types of cattle;

- is ready to use and no further additives are needed;
- is a concentrated source of protein in block form which leads to a low cost per unit protein;
- it helps with digestion of dry roughage and pasture since it provides protein and trace minerals that stimulate the rumen bacteria;
- provides maximum trace mineral levels which are essential for optimum appetite and fertility, as well as optimum functioning of immune system.

COMPOSITION OF RUMEVITE CATTLE BLOCK

REG NO V10932 (Act 36/1947) N-FF 0627

		(g/kg)			(mg/kg)
Crude Protein*	(min)	400	Manganese		1 000
Urea	(max)	124	Copper		120
Crude Fibre	(max)	20	Cobalt		2.5
Moisture	(max)	140	Iron		300
Calcium	(max)	48	Iodine		12
Phosphorus	(min)	4	Zinc		800
Magnesium	(min)	19	Selenium		2.5
Sulphur	(min)	6.2	Vitamin A (IU/kg)		20 400
*89 % derived from NPN					

Packaging – 25 kg


See urea warning.

FEEDING INSTRUCTIONS

Cattle: Supply 1 **Rumevite Cattle Block** per 15 animals.

RECOMMENDED DAILY INTAKE

Cattle: 375 - 600 g/animal/day.



VOERMOL

UREA WARNING

UREA

The majority of licks contain urea that can be dangerous under certain conditions. During the formulation and manufacturing process of **Voermol** licks, great care is taken to minimise the risks of feeding urea.

The possibilities of urea poisoning can be further reduced by taking note of the following:

- use licks as prescribed;
- limit lick intake to the prescribed maximum levels;
- feed the licks together with sufficient grazing and/or roughage;
- take special care during feed scarcity (drought conditions) to prevent high lick intake;
- prevent excessive lick intake in the initial stages (salt hunger) by allowing the animals access to **Voermol Rumevite 6P** for the first 14 days;
- do not let licks get wet.

See urea warning on the back of the bag.

UREA WARNING

This farm feed contains NPN (Non-Protein Nitrogen) sources and must therefore be fed strictly according to instructions for use.

- Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf/sheep or 2 – 4 bottles per head of cattle (1 bottle = 750 ml).
- Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
- Do not feed this farm feed indiscriminately with other NPN containing farm feeds. Consult an animal scientist.
- Mix concentrate thoroughly with the prescribed ingredients.
- This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
- Keep lick troughs filled and prevent gluttonous eating by hungry animals. A constant daily intake can help prevent poisoning.
- Before feeding a NPN containing lick, feed an ordinary salt / phosphate lick for at least 7 days.

GMO STATEMENT

These animal feeds may contain ingredients derived from GMO (genetically modified organisms) origin.



VOERMOL