

YaraVita Zintrac 700

RECOMMENDED APPLICATION RATES AND TIMINGS

Apples, Pears: 1 l/ha at bud burst and after the end of flowering. Water rate: 500 l/ha minimum. Alternatively, 50 mls / 100 l at bud burst and after the end of flowering, based on a typical TRV of 2500 l/ha and a maximum TRV of 4000 l/ha.

Avocado: 1 to 2 l/ha during spring flush, then during summer flush and again during autumn flush (up to 60 days before harvest). Water rate: 500 l/ha minimum.

Bananas: Up to 1 l/ha. Spray as required. Repeat applications may be necessary where low or marginal levels of zinc exist. Water rate: 30-300 l/ha.

Brassicac: Up to 2 l/ha applied at 4-8 leaves. Water rate: 50-500 l/ha.

Canola: 0.5 to 1 l/ha at the 4 to 9 leaf stage. Water rate: 30 to 500 l/ha.

Cereals: 0.5 to 1 l/ha from 2 leaf stage to first node detectable (Craven 9 - 10/11), repeated once or twice at 10 to 14 day intervals in cases of moderate to severe deficiency. Water rate: 50 to 200 l/ha.

Citrus: 1 to 2 l/ha during spring flush and again during autumn flush. In both instances a repeat application 10-14 days later may be required where moderate to severe deficiency exists. Water rate: 500 l/ha minimum. Alternatively, 50 mls / 100 l during spring flush and again during autumn flush. In both instances a repeat application 10-14 days later may be required where moderate to severe deficiency exists, based on a typical TRV of 2500 l/ha and a maximum TRV of 4000 l/ha.

Grapevines: Up to 1 l/ha applied at flower buds visible and again at flower buds separated. Water rate: 200-1,000 l/ha.

Macadamia: 50 mls/100l to give a maximum of 2 l/ha per application. Apply at spring flush. A repeat application should be made at the same rate at autumn flush. Avoid flowering. Water rate: apply sufficient to provide good cover but avoid run-off. (Minimum: 500 l/ha).

Maize: Up to 1 l/ha at 4-8 leaves. Water rate: 200 l/ha.

Mango: 1 to 1.5 l/ha applied during dormancy and again at early flush. Also, 1 to 1.5 l/ha applied post-harvest. Water rate: 500 l/ha minimum.

Potatoes: Up to 1 l/ha, 1 week after 100% emergence. Water rate: 200 l/ha.

Stone Fruit (Apricots, Cherries, Nectarines, Peaches, Plums): 1 l/ha at winter bud and again at pink bud. Water rate: 500 l/ha minimum. Alternatively, 50 mls / 100 l at winter bud and again at pink bud, based on a typical TRV of 2500 l/ha and a maximum TRV of 4000 l/ha.

Strawberry (Field Grown): One application of 0.5 l/ha at green bud followed by two applications of 0.25 l/ha applied at white bud and post-harvest regrowth. Water rate: 200 to 500 l/ha.

Sugar Cane: Up to 1 l/ha when cane is between 60 and 120 cm tall. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: 30 to 200 l/ha. Alternatively: up to 2.5 l/ha 50 days after harvest or 70 days after sowing. Water rate: 30 l/ha.

Tomato (Field Grown): 0.3 to 1 l/ha when plants are at 4 to 6 leaf stage. Repeat at 10 to 14 day intervals if necessary. Water rate: 70 to 500 l/ha.

For further crop recommendations contact your Yara representative.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Store all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes (temperatures exceeding 40°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

