

ON COLD NEUTRONS TRAPPING IN NANOSTRUCTURES & NEUTRON LIFETIME CONSIDERATIONS

Maaza@tlabs.ac.za
Maazam@unisa.ac.za



“QUANTUM TRAPPING OF COLD NEUTRONS WITH PICOSECOND PRECISION”

Maaza@tlabs.ac.za
Maazam@unisa.ac.za



“QUANTUM TRAPPING OF COLD NEUTRONS WITH PICOSECOND PRECISION”

Maaza@tlabs.ac.za
Maazam@unisa.ac.za



**INTERNATIONAL YEAR OF
Quantum Science
and Technology**

ON COLD NEUTRONS TRAPPING IN NANOSTRUCTURES & NEUTRON LIFETIME CONSIDERATIONS

Maaza@tlabs.ac.za
Maazam@unisa.ac.za



OUTLINE



1- Neutron lifetime: Generalities

2-Background: Wave-Matter Quantum Neutron Optics,

3A-Trapping neutron in Single Fabry-Perot nanoresonator Heisenberg uncertainty governed precision: Neutron lifetime considerations,

3B-Free propagation of neutrons in coupled Fabry-Perot nanoresonators with ^{10}B

4- Conclusion/Foresight

NEUTRON LIFETIME & S.M.

- **Lifetime:** $\langle \tau_n \rangle$ Used to predict the ratio of protons to helium atoms in the primordial universe and to search for physics beyond the Standard Model of particle physics. Relative abundance H/He During Universe 1st few mins:

$$n/p = \exp(-\Delta m / kT_{\text{freeze}}) \approx 1/6.$$

PARTICLE PHYSICS

Physicists close in on neutron puzzle

Researchers are narrowing down their measurements of how long the subatomic particle survives on its own.

BY ALEXANDRA WITZE IN DENVER, COLORADO

Physicists are drawing nearer to answering a long-standing mystery of the Universe: how long a neutron lives.

Neutrons are electrically neutral particles that usually combine with protons to make up atomic nuclei. Some neutrons are not bound up in atoms; these free-floating neutrons decay radioactively into other particles in minutes.

But physicists can't agree on precisely how long it takes a neutron to die. Using one laboratory approach, they measure the average neutron lifetime as 14 minutes 39 seconds. Using a different approach, they get 8 seconds longer.

"We don't know why they're different," says Shannon Hoogerheide, a physicist at the National Institute of Standards and Technology (NIST) in Gaithersburg, Maryland. "We really need to understand and eliminate this discrepancy." She and other scientists debated new ways to solve the problem this month at a meeting of the American Physical Society in Denver, Colorado.

Pinpointing the lifetime of a neutron is important for understanding how much hydrogen, helium and other light elements formed in the first few minutes after the Universe was born 13.8 billion years ago. Scientists also think that pinning down the neutron's lifetime would help to constrain measurements of other subatomic particles.

One way of clocking the neutron's lifespan is to put some of the particles in a bottle and count how many are left after a period of time. This 'bottle' method has been tried at several laboratories, including the Los Alamos National Laboratory in New Mexico¹ and the Institut Laue-Langevin in Grenoble, France. On average, they come up with a neutron lifetime of 14 minutes 39 seconds.

The other way is to feed neutrons into a detector that counts the protons created as the neutrons decay. This 'beam' method has been used at NIST and at the Japan Proton Accelerator Research Complex in Tokai. The Japanese work has just begun, but the NIST team reported in 2013 that its neutrons live eight seconds longer, on average, than those in the bottle method².

That's a big problem, because the beam and bottle measurements don't overlap, even when their margins of error are taken into account. So physicists have been looking for ways to explain why neutrons might be disappearing from bottles faster than from beams.

One possibility is that one of the two methods is doing something wrong. In that case, researchers might want to combine beam

442 | NATURE | VOL 568 | 25 APRIL 2019 | © 2019 Springer Nature Limited. All rights reserved.

ON COLD NEUTRONS TRAPPING IN NANOSTRUCTURES & NEUTRON LIFETIME CONSIDERATIONS

The Nobel Prize in Physics 1979



Sheldon Lee Glashow
Prize share: 1/3



Abdus Salam
Prize share: 1/3



Steven Weinberg
Prize share: 1/3

The Nobel Prize in Physics 1979 was awarded jointly to Sheldon Lee Glashow, Abdus Salam and Steven Weinberg *"for their contributions to the theory of the unified weak and electromagnetic interaction between elementary particles, including, inter alia, the prediction of the weak neutral current"*.

Photos: Copyright © The Nobel Foundation.



The Abdus Salam
International Centre
for Theoretical Physics



The Spirit of Abdus Salam Award



Malik Maaza




Concetta Mosca



Adnan Shihab-Eldin

NEUTRON LIFETIME: β -DECAY



ON NEUTRON WAVE-PACKETS, NEUTRON TRAPPING IN NANOSTRUCTURE & NEUTRON LIFETIME

- $F_{\mu\nu}$: Salam-Weinberg-Glashow Model /Electroweak forces, W/Z + Strong forces (1976)
- H: Higgs Boson
- R: Gravitational component/ Space-time curvature

$$\mathcal{L} = \sqrt{g} \left\{ R - \frac{1}{4} F^{\mu\nu} F_{\mu\nu} + \theta F^{\mu\nu} \tilde{F}_{\mu\nu} + \bar{\Psi} \not{D} \Psi + Y_{ij} \Psi_i \Psi_j + h.c. - |D_\mu H|^2 - V(H) \right\}$$

$\cong$ Our Universe... so far


This equation describes in a simplified condensed manner the fundamental laws governing the Universe at our present understanding. It includes the known fundamental particles: ψ represents all the fundamental matter particles (fermions) such as electron, quarks, etc. $F_{\mu\nu}$ describes the electro-weak forces (Salam-Weinberg-Glashow model) including photons (quanta of light) as well as the W and Z bosons, and strong forces that provide the binding of quarks into protons and neutrons. H denotes the Higgs particle which gives masses to various particles and has been recently discovered in the LHC experiment at CERN. It was the last missing piece of evidence in the context of the Salam-Weinberg-Glashow model. Finally, the space-time curvature, R describes Einstein's theory of gravitational interactions with g representing the space-time metric.




The Abdus Salam
International Centre
for Theoretical Physics



NEUTRON: COSMOLOGY PERSPECTIVE



NEUTRON LIFETIME & S.M.




- Within the Standard Model, the **Cabibbo-Kobayashi-Maskawa matrix** describes the strength of transitions between quarks in weak interactions. The first row of the matrix is given by a series of

$$\begin{bmatrix} |V_{ud}| & |V_{us}| & |V_{ub}| \\ |V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}| & |V_{ts}| & |V_{tb}| \end{bmatrix} = \begin{bmatrix} 0.97370 \pm 0.00014 & 0.2245 \pm 0.0008 & 0.00382 \pm 0.00024 \\ 0.221 \pm 0.004 & 0.987 \pm 0.011 & 0.0410 \pm 0.0014 \\ 0.0080 \pm 0.0003 & 0.0388 \pm 0.0011 & 1.013 \pm 0.030 \end{bmatrix}.$$

Using those values, one can check the unitarity of the CKM matrix. In particular, we find that the first-row matrix elements give:

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9985 \pm 0.0005;$$

NEUTRON LIFETIME & S.M.



- Within the Standard Model, the **Cabibbo-Kobayashi-Maskawa matrix** describes the strength of transitions between quarks in weak interactions. The first row of the matrix is given by a series of


$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9994 \pm 0.0005.$$

Most precise value of $|V_{ud}|$ is derived from neutron decay

$$|V_{ud}|^2 = (4908.7 \pm 1.9) / \langle \tau_n \rangle (1 + 3\zeta^2)\zeta,$$

ζ , being the ratio of the axial-vector and vector couplings.

NEUTRON LIFETIME & S.M.




The average neutron lifetime derived from:

Beam approach was found to be $\langle \tau_n \rangle_{\text{Beam}} \sim 888.0 \pm 2.0 \text{ s}$


Bottle technique of about $\langle \tau_n \rangle_{\text{Bottle}} \sim 879.4 \pm 0.6 \text{ s}$

This difference of $\sim 10 \text{ s}$ persists for years up to now

COLD & ULTRA-COLD NEUTRONS



NEUTRON: β -DECAY & NUCLEAR-CHART PERSPECTIVE



β^+ DECAY

NEUTRON: ELEMENTS PERSPECTIVE

ELEMENTLERIN PERİYODİK TABLOSU



NEUTRON WAVE-MATTER: ODYSSEY-1987

Journal of Neutron Research 00 (20xx) 1–16
DOI 10.3233/JNR-220015
IOS Press

1

On the trapping of neutrons in Fabry–Pérot nano-structures and potential applications for cold neutron lifetime Investigations

M. Maaza^{a,b,*}, B. Pardo^{c,†}, D. Hamidi^{a,b}, M. Akbari^{a,b}, R. Morad^{a,b}, M. Henini^{a,b,d} and A. Gibaud^{a,b,e}

^a UNESCO-UNISA Africa Chair in Nanosciences-Nanotechnology, University of South Africa, PO Box 392, Pretoria, South Africa

^b Nanosciences African Network (NANOAFNET), iThemba Laboratories for Accelerators Based Sciences, iThemba LABS-National Research Foundation, PO Box 722, Somerset West, Western Cape Province, South Africa

^c Institut d'Optique Théorique & Appliquée, Université Paris-Saclay, France

^d Physics Department, University of Nottingham, Nottingham, UK

^e IMMM, UMR 6283 CNRS, Université of Le Maine, Bd O. Messiaen, 72085 Le Mans cedex 09, Le Mans, France

Abstract. Correlated to the neutron total reflection phenomenon is the so-called frustrated total reflection, also known as neutron channeling, observed with both thermal and cold neutrons. Within this contribution, such a phenomenon is validated in various additional distinctive Fabry–Pérot nano-resonating configurations; namely in: (i) dual reflection and transmission neutron Fabry–Pérot nano-resonator (Ni/V/Ni/Si substrate), (ii) isotope-based neutron Fabry–Pérot nano-resonator ($^{58}\text{Ni}/^{62}\text{Ni}/^{58}\text{Ni}/\text{Silicon}$ substrate), and (iii) multilayered neutron Fabry–Pérot nano-resonator of 8 superposed ($\text{B}_4\text{C}/\text{Ti}/\text{B}_4\text{C}$) single nano-resonators. While such Fabry–Pérot nano-resonators allow effective neutron trapping, the precision of the trapping time of free neutrons in such nano-resonators is governed by the Heisenberg uncertainty and hence offers, a priori, an additional attractive precise approach for potential lifetime investigations. Depending on the configuration of the Fabry–Pérot nano-resonators and the available cold neutron beam, the trapping time is found to be within the temporal regime of 3 to 19 ps. While the main intention of this contribution is to validate the possibility of trapping cold neutrons in nano-structured Fabry–Pérot resonators with a picosecond precision in various configurations, it is hoped that these preliminary results will attract the interest of the neutron lifetime community specifically and the neutron scattering community in general. The potential integration of such trapping method into the bottle or beam methods would elucidate the origin of the difference in neutron lifetime between the two approaches.

Keywords: Neutron trapping, Fabry–Pérot nano-resonators, frustrated total reflection, neutron waveguiding, neutron lifetime, total reflection, Zeldovich–Vinogradov configuration, B_4C , Fermi pseudo-potential



• JNR
~ 2024

NEUTRON WAVE-MATTER: ODYSSEY-1987

Nuclear Instruments and Methods in Physics Research A326 (1993) 531–537
North-Holland

**NUCLEAR
INSTRUMENTS
& METHODS
IN PHYSICS
RESEARCH**
Section A

New optical transmission device to produce high monochromatic and high polarized neutron beams based on the tunneling frustrated total reflection in neutron guides

M. Maaza

Laboratoire Léon Brillouin, Commissariat à l'Energie Atomique, Centre National de la Recherche Scientifique, Bat 563, Centre d'Etude Nucléaire de Saclay, 91191, Gif-sur-Yvette, France

B. Pardo and F. Bridou

Institut d'Optique Théorique et Appliquée, Bat 503, Université Paris Sud, Orsay, France

Received 23 July 1992 and in revised form 12 October 1992

It has been shown that the tunneling of neutrons can be used in classical neutron guides to extract high monochromatic and/or high polarized cold neutron beams. This can be achieved by implanting ^{58}Ni -Ti- ^{58}Ni or ^{58}Ni -Co- ^{58}Ni Fabry-Pérot interferometers under total reflection condition in the neutron guide. According to the computed results, this tunneling phenomenon of neutrons manifests itself by the existence of sharp resonances in the ^{58}Ni total reflection plateau. These resonances are localized at different spectral positions $\lambda_{\pm}^r$ for the monochromatization and λ_{+}^r , λ_{-}^r for the polarization. The narrow spectral bandwidth of the tunneling transmitted beam can reach values of the order of $\Delta\lambda \approx 0.01 \text{ \AA}$ and a strong transmissivity higher than 99%. The number and the spectral position of these resonances vary mainly with the characteristics of the spacer layer Ti or Co whereas their spectral bandwidths are imposed by the thickness of the reflecting ^{58}Ni layers of the Fabry-Pérot interferometer. To obtain low resonance orders, the use of a nonmagnetic substrate with a negative nuclear scattering length density and a perfect collimated polychromatic neutron beam is required.

• NIMB
{ 1993

NEUTRON WAVE-MATTER: ODYSSEY-1987

Physics Reports 514 (2012) 177–198



ELSEVIER

Contents lists available at [SciVerse ScienceDirect](#)

Physics Reports

journal homepage: www.elsevier.com/locate/physrep



Nano-structured Fabry–Pérot resonators in neutron optics & tunneling of neutron wave-particles

M. Maaza^{a,b,*}, D. Hamidi^a

^a *Nanosciences African Network, iThemba LABS, National Research Foundation of South Africa, 1 Old Faure road, Somerset West 7129, POBox 722, Somerset West, Western Cape Province, South Africa*

^b *Faculty of Sciences, Pretoria-Tshwane University of Technology, Staatsartillerie Road, Pretoria, South Africa*

ARTICLE INFO

Article history:

Accepted 9 January 2012

Available online 16 January 2012

editor: D.L. Mills

Keywords:

Neutrons

Neutron optics

Total reflection

Frustrated total reflection

Tunneling phenomenon

Resonance

ABSTRACT

Correlated to the quantum mechanics wave-particle duality, the optical analogy between electromagnetic waves and cold neutrons manifests itself through several interference phenomena particularly the so called Frustrated Total Reflection i.e., the tunneling process in Fabry–Pérot nano-structured cavities. Prominent resonant situations offered by this configuration allow the attainment of numerous fundamental investigations and surface-interface studies as well as to devise new kinds of neutron optics devices. This review contribution reports such possibilities in addition to the recently observed peculiar Goos–Hänchen longitudinal shift of neutron wave-particles which was predicted by Sir Isaac Newton as early as 1730.

© 2012 Elsevier B.V. All rights reserved.

NEUTRON WAVE-MATTER: ODYSSEY-1987

J. Appl. Cryst. (1993). **26**, 327–333

Tunneling Reflection with Polarized Slow Neutrons: Polarized-Neutron Total Frustrated Reflection

BY M. MÂAZA

*Laboratoire Léon Brillouin, CEA-CNRS, Bâtiment 563, Centre d'Etudes Nucléaires de Saclay,
91191 Gif-sur-Yvette, France*

AND B. PARDO AND F. BRIDOU

Institut d'Optique Théorique et Appliquée, Bâtiment 503, Université Paris Sud, Orsay, France

(Received 2 July 1992; accepted 2 October 1992)

Abstract

This work describes the conditions required to observe the tunneling phenomenon in the grazing-angle reflection configuration of both polarized and unpolarized monochromatic neutron waves. According to the computed results, this neutron-tunneling phenomenon manifests itself by the existence of sharp resonances in the total reflection plateau. Their angular positions depend on the polarization state of the incident neutron waves. We treat the case where all these tunneling resonances (high and low orders) in the total-reflection plateau are observed. This can be

complete when the evanescent wave meets the second medium and a proportion of the incident electromagnetic evanescent wave will appear in the second medium. The reflectivity at the first interface (air–first-medium interface) will be less than total. This, as Baumeister (1967) has pointed out, is very similar to the behavior of fundamental particles tunneling through a potential barrier; such is the case for electrons in quantum semiconductor wells. Practically, this evanescent-waves effect is applied to the study of surface plasmons in Kretschmann geometry (Otto, 1968; Abelès, 1971).

NEUTRON WAVE-MATTER: ODYSSEY-1987

Physics Letters A 181 (1993) 276–282
North-Holland

PHYSICS LETTERS A

New generation of neutron interferometers based on tunneling frustrated total reflection of polarized neutron waves

M. Mâaza

*Laboratoire Léon Brillouin, Commissariat à l'Energie Atomique, Centre National de la Recherche Scientifique,
Bat. 563, Centre d'Etudes Nucléaires de Saclay, 91191 Gif-sur-Yvette, France*

and

B. Pardo

Institut d'Optique Théorique et Appliquée, Bat. 503, Université Paris Sud, Orsay, France

Received 3 May 1993; accepted for publication 10 August 1993

Communicated by J.P. Vigiér

We present a new generation of neutron interferometers using a new splitting mode. The method is based on the tunneling frustrated total reflection of neutron waves. The splitting of an incident neutron beam into two coherent partial beams is achieved by using magnetized Fabry–Perot resonators. These resonators which work under total reflection conditions, are constituted of stacks of $^{58}\text{Ni}/\text{Co}/^{58}\text{Ni}/\text{bulk Co}$. More specifically, when the resonance condition of the Fabry–Perot resonators is fulfilled, resonant spin down waves are totally tunneled while the corresponding spin up waves are totally reflected. The number of these tunneling resonances changes mainly with the thickness of the spacer layer of Co while their band-pass is linked to the thickness of the reflecting ^{58}Ni layers.

NEUTRON WAVE-MATTER: ODYSSEY-1987



ELSEVIER

21 November 1994

PHYSICS LETTERS A

Physics Letters A 195 (1994) 1–8

Shearing neutron interferometry and possibilities of studying interfacial diffusion processes between two highly dilute solutions ☆

M. Mâaza ^{a,b}, B. Pardo ^a, S. Shamlal Mallick ^a, F. Bridou ^a, D.A. Korneev ^b,
L.P. Chernenko ^b, C. Sella ^c

^a *Institut d'Optique Théorique et Appliquée, Bât. 503, Université Paris Sud, Orsay, France*

^b *Frank Laboratory of Neutron Physics, Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russian Federation*

^c *Laboratoire de Physique des Matériaux, Centre National de la Recherche Scientifique,
1 place Aristide Briand, 92125 Meudon cedex, France*

Received 14 October 1993; revised manuscript received 6 September 1994; accepted for publication 13 September 1994

Communicated by J.P. Vigiér

Abstract

We deal with the adaptation of the classical shearing interferometry in neutron optics and suggest a shearing neutron interferometer device. A polarized monochromatic neutron plane wave-particle front is distorted in passing through a diffusion cell containing two liquids in accordance with the variation of the concentration in the solution. This distorted front is sheared into two coherent fronts. The related interference spectrum varies in time and the diffusion coefficient can be deduced. The proposed method would be convenient to measure diffusion coefficient for very low concentrations smaller than 10^{-3} g/cm^3 .

• Phys.Lett.A
{ 1994

• FLNP-JINR
{ 1993

NEUTRON WAVE-MATTER: ODYSSEY-1987



ELSEVIER

2 December 1996

PHYSICS LETTERS A

Physics Letters A 223 (1996) 145–148

Neutron tunneling and neutron lifetime in a Ni–V–Ni Fabry–Perot thin film resonator

M. Mâaza ^a, B. Pardo ^b, J.P. Chauvineau ^b, A. Raynal ^b, A. Menelle ^c, F. Bridou ^b

^a *Physics Department, University of the Witwatersrand, Wits 2050, Johannesburg, South Africa*

^b *Institut d'Optique Théorique & Appliquée, Bat. 503, Université d'Orsay 91403, Orsay cedex, France*

^c *Laboratoire Léon Brillouin, CEA-CNRS, Bat. 563, Centre d'Etudes Nucléaires de Saclay, Gif-sur-Yvette, France*

Received 13 June 1996; accepted for publication 16 September 1996

Communicated by P.R. Holland

Abstract

Observations are reported on the appearance of frustrated total reflection, i.e. the tunnelling phenomenon, with cold neutron wave-particles. This tunnelling resonance was observed in a thin film Ni–V–Ni Fabry–Perot cavity by time of flight grazing angle neutron reflectometry. This frustrated total reflection phenomenon was used to estimate the neutron lifetime, which varies from approximately 0.1 to 1 μ s.

NEUTRON WAVE-MATTER: ODYSSEY-1987



ELSEVIER

20 October 1997

PHYSICS LETTERS A

Physics Letters A 235 (1997) 19–23

Zeeman neutron tunneling in “Ni–Co–Ni” thin film resonators

M. Maaza^a, B. Pardo^b, J.P. Chauvineau^b, A. Menelle^c, A. Raynal^b, F. Bridou^b, J. Corno^b

^a *Physics Department, University of the Witwatersrand, Johannesburg, South Africa*

^b *Institut d'Optique Théorique & Appliquée, Bat. 503, Université d'Orsay, Orsay 91403, France*

^c *Laboratoire Leon Brillouin, Commissariat à l'Energie Atomique, Saclay, France*

Received 5 February 1997; revised manuscript received 2 June 1997; accepted for publication 4 July 1997

Communicated by J.P. Vigier

Abstract

Polarized neutron tunneling measurements are reported, achieved by using simple and double thin film Ni–Co–Ni Fabry–Perot resonators operating in the total reflection geometry in the polarized configuration. Due to the Zeeman magnetic interaction term in the cobalt layers mainly, a spatial separation of the spin up and down states results. The Zeeman tunneling splitting within such resonators can be exploited to produce monochromatic and polarized neutron beams in the range of $\lambda \geq 10$ Å. © 1997 Elsevier Science B.V.

Keywords: Zeeman interaction; Tunneling; Polarized neutrons



PERGAMON

Solid State Communications 111 (1999) 23–28

solid
state
communications

V–Ni multilayered monochromators and supermirrors for cold neutrons

M. Maaza^{a,*}, J.P. Chauvineau^b, B. Pardo^b, A. Raynal^b, A. Menelle^c, F. Bridou^b,
J. Corno^b

^a*Physics Department, University of the Witwatersrand, Wits 2050, Johannesburg, South Africa*

^b*Institut d'Optique Théorique and Appliquée, Université Paris-Sud, Bat. 503, 91403 Orsay Cedex, France*

^c*Laboratoire Leon Brillouin, Commissariat à l'Energie Atomique, 91191 Gif-sur-Yvette, France*

Received 12 December 1998; accepted 15 March 1999 by R.T. Phillips

Abstract

We present experimental neutron reflectivity results showing that it is possible to use V–Ni multilayered systems as cold neutron monochromators and supermirrors. Compared to the Ni–Ti optics devices, the V–Ni stacks present sharp interfaces characterised by a low interfacial roughness and very small interfacial diffusion. © 1999 Elsevier Science Ltd. All rights reserved.

Keywords: A. Nanostructures; C. X-ray scattering; D. Optical properties; E. Neutron scattering



PERGAMON

Solid State Communications 112 (1999) 177–181

solid
state
communications

www.elsevier.com/locate/ssc

Thermal stability of Co–Ti multilayered neutron polarizers

M. Mâaza^{a,*}, M. Spegel^b, C. Sella^c, B. Pardo^d, A. Menelle^e, J. Corno^d, R. Gaziel^a

^a*Physics Department, University of the Witwatersrand, Wits 2050, Johannesburg, South Africa*

^b*Atominstytut der Österreichischen Universitäten, Schüttelstrasse 115, A-1020 Vienna, Austria*

^c*Laboratoire D'Optique des Solides, Université de Paris VI, Jussieu-Paris, France*

^d*Institut d'Optique Théorique and Appliquée, Université Paris-Sud, Bat. 503, 91403 Orsay, France*

^e*Laboratoire Léon Brillouin, Laboratoire-commun CEA-CNRS, Bat.563 91191 Gif-sur-Yvette, France*

Received 12 December 1998; accepted 15 March 1999 by R.T. Phillips

Abstract

Thermal stability of multilayered Co–Ti neutron optic polarizers with a period of the order of 103 Å is investigated. The diffusion kinetics is determined by using the Du Mond and Youtz's method with grazing angle neutron reflectometry in the temperature range of 293–723 K. It was found that the diffusion is mainly directed from Co-layers towards the Ti-layers. The effective interdiffusion coefficient D_{eff} of cobalt into titanium is calculated from the rate of decrease of the first reflected Bragg peak related to the artificial periodicity of the multilayer with the annealing temperature T . The temperature dependence of D_{eff} is found to be described approximately by $D_{\text{eff}} \approx (D_0 \exp(-0.25 \text{ eV}/k_B T)) \text{ cm}^2 \text{ s}^{-1}$. © 1999 Elsevier Science Ltd. All rights reserved.

Keywords: A. Surfaces and interfaces; A. Thin films; E. Neutron scattering

NEUTRON WAVE-MATTER: ODYSSEY-1987

Journal of Neutron Research 00 (20xx) 1–16
DOI 10.3233/JNR-220015
IOS Press

1

On the trapping of neutrons in Fabry–Pérot nano-structures and potential applications for cold neutron lifetime Investigations

M. Maaza^{a,b,*}, B. Pardo^{c,†}, D. Hamidi^{a,b}, M. Akbari^{a,b}, R. Morad^{a,b}, M. Henini^{a,b,d} and A. Gibaud^{a,b,e}

^a UNESCO-UNISA Africa Chair in Nanosciences-Nanotechnology, University of South Africa, PO Box 392, Pretoria, South Africa

^b Nanosciences African Network (NANOAFNET), iThemba Laboratories for Accelerators Based Sciences, iThemba LABS-National Research Foundation, PO Box 722, Somerset West, Western Cape Province, South Africa

^c Institut d'Optique Théorique & Appliquée, Université Paris-Saclay, France

^d Physics Department, University of Nottingham, Nottingham, UK


^e IMMM, UMR 6283 CNRS, Université of Le Maine, Bd O. Messiaen, 72085 Le Mans cedex 09, Le Mans, France

Abstract. Correlated to the neutron total reflection phenomenon is the so-called frustrated total reflection, also known as neutron channeling, observed with both thermal and cold neutrons. Within this contribution, such a phenomenon is validated in various additional distinctive Fabry–Pérot nano-resonating configurations; namely in: (i) dual reflection and transmission neutron Fabry–Pérot nano-resonator (Ni/V/Ni/Si substrate), (ii) isotope-based neutron Fabry–Pérot nano-resonator ($^{58}\text{Ni}/^{62}\text{Ni}/^{58}\text{Ni}/\text{Silicon}$ substrate), and (iii) multilayered neutron Fabry–Pérot nano-resonator of 8 superposed ($\text{B}_4\text{C}/\text{Ti}/\text{B}_4\text{C}$) single nano-resonators. While such Fabry–Pérot nano-resonators allow effective neutron trapping, the precision of the trapping time of free neutrons in such nano-resonators is governed by the Heisenberg uncertainty and hence offers, a priori, an additional attractive precise approach for potential lifetime investigations. Depending on the configuration of the Fabry–Pérot nano-resonators and the available cold neutron beam, the trapping time is found to be within the temporal regime of 3 to 19 ps. While the main intention of this contribution is to validate the possibility of trapping cold neutrons in nano-structured Fabry–Pérot resonators with a picosecond precision in various configurations, it is hoped that these preliminary results will attract the interest of the neutron lifetime community specifically and the neutron scattering community in general. The potential integration of such trapping method into the bottle or beam methods would elucidate the origin of the difference in neutron lifetime between the two approaches.


Keywords: Neutron trapping, Fabry–Pérot nano-resonators, frustrated total reflection, neutron waveguiding, neutron lifetime, total reflection, Zeldovich–Vinogradov configuration, B_4C , Fermi pseudo-potential



COLD & ULTRA-COLD NEUTRONS




- **Cold Neutrons** (0 eV; 0.025 eV). Neutrons in thermal equilibrium with liquid deuterium. This spectrum is used for neutron scattering experiments.
- **Thermal Neutrons**. Neutrons in thermal equilibrium with a surrounding at 20°C (68°F) for Maxwellian distribution is **0.025 eV** (~2 km/s). This spectrum constitutes the most important spectrum in thermal reactors.
- **Epithermal Neutrons** (0.025 eV; 0.4 eV). Neutrons of kinetic energy above the thermal range. Reactor designs operate with an epithermal neutron spectrum. This design allows reaching a higher fuel breeding ratio than in thermal reactors.
- **Cadmium Neutrons** (0.4 eV; 0.5 eV). Neutrons of kinetic energy below the **cadmium cut-off** energy. One cadmium isotope, ^{113}Cd , absorbs neutrons strongly only if they are below ~0.5 eV (**cadmium cut-off energy**).
- **Epicadmium Neutrons** (0.5 eV; 1 eV). Neutrons of kinetic energy above the cadmium cut-off energy. These neutrons are not absorbed by cadmium.
- **Slow Neutrons** (1 eV; 10 eV).
- **Resonance Neutrons** (10 eV; 300 eV). **The resonance neutrons** are called resonance for their special behavior. At resonance energies, the **cross-sections** can reach peaks more than 100x higher than the base value of the cross-section. At these energies, the neutron capture significantly exceeds the probability of **fission**.



Neutrons of kinetic energy below the cadmium cut-off energy (~0.5 eV) are strongly absorbed by ^{113}Cd .
Source: JANIS (Java-based nuclear information software) www.oecd-nea.org/janis/

COLD & ULTRA-COLD NEUTRONS



NEUTRON OPTICS: BASICS

The propagation of a neutron through a medium is governed by the matter-wave Schrödinger equation. In a time independent potential, under steady state conditions, the Schrödinger equation in the position representation is,

$$\left[-\frac{\hbar^2}{2m}\nabla^2 + V(\mathbf{r}) \right] \Psi(\mathbf{r}) = E\Psi(\mathbf{r}),$$

$$V(\mathbf{r}) = \sum_j \frac{2\pi\hbar^2}{m} b_j \delta(\mathbf{r} - \mathbf{r}_j),$$

where $V(\mathbf{r})$ is the potential energy of the particle, and E is the total energy of the particle.

The equation can be written in a form that is similar to the Helmholtz equation in classical optics as,

$$\nabla^2 \Psi(\mathbf{r}) + K(\mathbf{r})^2 \Psi(\mathbf{r}) = 0,$$

with $K(\mathbf{r}) = \sqrt{2m[E - V(\mathbf{r})]}/\hbar$ is the medium dependent wavevector. $V(\mathbf{r})$ can take various forms including electromagnetic, gravitational, and nuclear. Of interest to us is the nuclear and magnetic interaction which will be explored in the next section in detail.

One common way to characterize them is via the refractive index, defined as [2, 3],

$$n \equiv \frac{K(\mathbf{r})}{k} = \sqrt{\frac{2m[E - V(\mathbf{r})]}{2mE}} \simeq 1 - \frac{\bar{V}}{2E},$$

$$V(\mathbf{r}) = \sum_j \frac{2\pi\hbar^2}{m} b_j \delta(\mathbf{r} - \mathbf{r}_j),$$

where $k = \sqrt{2mE}/\hbar$ is the wavevector in free space. The approximation in Eq. (1.7) is valid for thermal neutrons as the potential $V(\mathbf{r})$ can be expressed in a form equal to the optical potential, $\bar{V}$, which for most materials is of the order 10^{-5} eV .



NEUTRON OPTICS: BASICS

$$V_O = \frac{2\pi\hbar^2}{m}N\bar{b}, \text{ where } N\bar{b} = \sum_j b_j\delta(\mathbf{r} - \mathbf{r}_j). \quad (1.9)$$

$N\bar{b}$ is the local mean scattering length density, which represents the response of the overall system as multiple copies of a single atom of the same kind. Generally, $\bar{b}$ is complex with typical values in the femtometre scale. Moreover, this value leads to a scattering cross-section defined as $\bar{\sigma}_s = 4\pi|\bar{b}|^2$ and an absorption cross-section defined as $\bar{\sigma}_a = 4\pi\text{Im}[\bar{b}]^2k^{-1}$, where $\text{Im}[\bar{b}]$ is the imaginary component.

When a neutron beam is shined on an absorbing target of effective thickness D , the intensity of transmitted neutron is related to the incident intensity, I_0 , by [19],

$$I = I_0 e^{-\bar{\sigma}_a N D}. \quad (1.10)$$

Materials with high $\bar{\sigma}_a$ including lead, cadmium, gadolinium, are commonly used as neutron absorbers. For most materials, $\text{Re}[\bar{b}] > 0$, with the few exceptions of ^2H , ^{48}Ti , and ^{62}Ni

The general form of the complex refractive index from a spin-independent scattering material is [3, 2],



$$n \equiv 1 - \frac{\lambda^2 N}{2\pi} \sqrt{\bar{b}^2 - \left(\frac{\bar{\sigma}_r}{2\lambda}\right)^2} + i \frac{\bar{\sigma}_r N \lambda}{4\pi}, \quad (1.11)$$

X-RAYS OPTICS:

Untersuchungen zur Totalreflexion von Röntgenstrahlen¹⁾

Von Heinz Kiessig

(Mit 30 Figuren)

Einleitung

Nachdem der Nachweis erbracht worden war, daß auch die Röntgenstrahlen sich meßbar brechen und reflektieren lassen, sind in den letzten Jahren zahlreiche Arbeiten gemacht worden, die sich mit dem quantitativen Studium dieser Erscheinung befassen.²⁾ Das Ziel dieser Arbeiten war fast ausschließlich die Messung des Brechungsindex in Abhängigkeit von der Wellenlänge und den Materialien, um die Dispersionskurve für Röntgenstrahlen aufstellen zu können. Es zeigte sich, daß die Drude-Lorentzsche Dispersionstheorie nicht nur qualitativ anwendbar ist, sondern auch quantitativ die Verhältnisse gut wiedergibt. Da für Röntgenstrahlen der Brechungsindex n kleiner als 1 ist, aber nur wenig von 1 abweicht, setzt man üblicherweise

$$n = 1 - \delta$$

und gibt immer nur den Wert δ an. Es lautet dann die Lorentz-sche Dispersionsformel

$$\delta = \frac{e^2}{2\pi m} \sum \frac{N_K}{\nu^2 - \nu_K^2},$$

wobei

ν die Frequenz der einfallenden Strahlung,


ν_K die Eigenfrequenzen der Elektronen und

N_K die Anzahl der Elektronen mit der Eigenfrequenz ν_K in dem Einheitsvolumen bedeutet.

Die Summe ist über alle Eigenfrequenzen zu erstrecken.

1) Dissertation der Technischen Hochschule München.


2) Zusammenfassende Darstellungen über das vorliegende Material



Neutron Wave-Packet == Photon in S-polarization state

NEUTRON OPTICS: BASICS

Neutron Particle



Discontinuous scattering medium
Delta function type interaction

$$V_O = \frac{2\pi\hbar^2}{m} N\bar{b}, \text{ where } N\bar{b} = \sum_j b_j \delta(\mathbf{r} - \mathbf{r}_j).$$

Neutron Wave Packet



Continuous scattering medium

$$n \equiv 1 - \frac{\lambda^2 N}{2\pi} \sqrt{\bar{b}^2 - \left(\frac{\bar{\sigma}_r}{2\lambda}\right)^2} + i \frac{\bar{\sigma}_r N \lambda}{4\pi},$$

TOTAL REFLECTION “Fermi, 1946”,

PHYSICAL REVIEW

VOLUME 71, NUMBER 10

MAY 15, 1947

Interference Phenomena of Slow Neutrons

E. FERMI AND L. MARSHALL

Argonne National Laboratory and University of Chicago, Chicago, Illinois

(Received February 7, 1947)

Various experiments involving interference of slow neutrons have been performed in order to determine the phase of the scattered neutron wave with respect to the primary neutron wave. Theoretically this phase change is very close to either 0° or 180° . The experiments show that with few exceptions the latter is the case. The evidence is based on the following types of measurements: (a) measurement of the intensities of Bragg reflection of various orders of many crystals, and comparison with the theoretical values of the form factor; (b) total scattering cross section of gas molecules for wave-lengths long compared with the molecular dimensions; and (c) determination of the limiting angles for total reflection of neutrons on various mirrors. The elements Ba, Be, C, Ca, Cu, F, Fe, Mg, N, Ni, O, Pb, S, and Zn were found to scatter neutrons with 180° phase difference; Li and probably Mn scatter with zero phase difference. The five elements I, Br, Cl, K, and Na behave alike and the phase with which they scatter is tentatively identified as 180° . Coherent scattering cross sections have been determined for several elements.

INTERFERENCE OF SLOW NEUTRONS

675




FIG. 4. Monochromatic total reflection on mirrors.






FIG. 3. Spectrum filtered through BeO.


TOTAL REFLECTION PHENOMENON



TOTAL REFLECTION PHENOMENON


3A-NEUTRON OPTICS :TOTAL REFLECTION

- Quantum Mechanics governed phenomena
- Maximize the reactor capabilities via n-guides




NEUTRON OPTICS:N.WAVE-PACKET

WAVE BEHAVIOR OF 'n'






- Total Reflection “Fermi, 1946”,
- Polarization “Gukasov, 1954”,
- Thin Film Interference “Maier-Leibnitz, 1962”,
- Magnetic polarization/spin flipping “Drabkin, 1962”
- Prism Deflection “Landkammer/ Korpiun, 1966”,
- Neutron Interferometry “Rauch, 1973”,
- Supermirrors “Mezei, 1978”,
- Neutron Tunneling “Refl. mode, Steyerl, 1981”,
- Sagnac Effect “Werner, 1985”,
- Aharonov-Bohm Effect “Collela, 1993”,
- Neutron Tunneling “Ref./Trans, Polar” “Maaza & al, 1996”,
- Neutron Tunneling “Modelling” “Ignatovich & al, 1997”,
- Neutron Goos-Hanschen “de Haan & al, 2010,
- Neutron Holography “Sarenac, Pushin & al, 2016”.


FRUSTRATED TOTAL REFLECTION.




TOTAL REFLECTION of NEUTRON WAVE-PACKETS.




1-NEUTRON TRAPPING: NEUTRON CONFINEMENT IN NANO-F.P.



$$\tau \approx m / h k \Delta k$$



FRUSTRATED TOTAL REFLECTION OF NEUTRON WAVEPACKETS



Resonance order ν	Resonance wavelength λ_{res} (Å)	$\Delta\lambda_{res}$ (Å)	k (Å ⁻¹)	Δk (10 ⁻² Å ⁻¹)	Trapping time τ (ps)
1	17.38	1.09	3.561	2.267	19.35
2	12.99	0.85	4.837	3.165	10.36
3	10.56	0.73	5.950	4.113	6.48
4	9.02	0.61	6.966	4.711	4.83
5	7.87	0.61	7.984	6.188	3.21

The average neutron lifetime derived from:

$$\langle T_n \rangle_{\text{Beam}} \sim 888.0 \pm 2.0 \text{ s}$$

$$\langle T_n \rangle_{\text{Bottle}} \sim 879.4 \pm 0.6 \text{ s}$$

Trapping time in Fabry-Perot Nano-resonator'' $\pm 10^{-12} \text{ s}$

NEUTRON LIFETIME: BOTTLE TECH.

- QED: The Magneto-Gravitational UCN Trap
- Lifetime: $\tau_n = 877.7 \pm 0.6 \text{ s}$

Science

RESEARCH ARTICLES

Cite as: R. W. Pattie Jr. *et al.*, *Science*
10.1126/science.aan8895 (2018).

Measurement of the neutron lifetime using a magneto-gravitational trap and in situ detection

R. W. Pattie Jr.,¹ N. B. Callahan,² C. Cude-Woods,^{1,3} E. R. Adamek,² L. J. Broussard,⁴ S. M. Engel,⁶ D. E. Fellers,¹ W. Fox,² P. Geltenbort,⁷ K. P. Hickerson,⁸ M. A. Hoffbauer,¹ A. T. Hol MacDonald,¹ M. Makela,¹ C. L. Morris,¹ J. D. Ortiz,¹ J. Ramsey,¹ D. J. Salvat,¹¹ A. Saunders,⁹ Tang,¹ J. Vanderwerp,² B. Vogelaar,⁵ P. L. Walstrom,¹ Z. Wang,¹ W. Wei,¹ H. L. Weaver,¹ J. Zeck^{1,3}

¹Los Alamos National Laboratory, Los Alamos, NM 87545, USA. ²Center for Exploration of Energy and Matter and 47408, USA. ³Triangle Universities Nuclear Laboratory and North Carolina State University, Raleigh, NC 27695, USA. ⁴Department of Physics, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA. ⁵Institut Laue-Langevin, Grenoble, France. ⁶Kellogg Radiation Laboratory, California Institute of Technology, Pasadena, CA 91125, USA. ⁷Department of Physics and Astronomy, DePaul University, Chicago, IL 60614, USA. ⁸Department of Physics, University of Washington, Seattle, WA 98195-1560, USA. ⁹Joint Institute for Nuclear Research, Dubna, Moscow Region, 151400, Russia.

*Corresponding author. Email: asaunders@lanl.gov

†Present address: Sandia National Laboratories, Albuquerque, NM 87185, USA.

The precise value of the mean neutron lifetime, τ_n , plays an important role in nuclear physics and cosmology. It is used to predict the ratio of protons to helium atoms produced in the early universe. The search for physics beyond the Standard Model of particle physics. We report on a new method for measuring the neutron lifetime by using a magneto-gravitational trap and in situ detection. This method is different from previous trap experiments by levitating polarized ultracold neutrons in a magneto-gravitational trap and detecting them in situ. As a result of this approach and the improved detection efficiency, the lifetime reported here $[877.7 \pm 0.7 \text{ (stat)} + 0.4/-0.2 \text{ (sys)} \text{ seconds}]$ is more precise than the quoted uncertainties.




Fig. 1. Layout of the UCN beam line and trap used for these measurements.

NEUTRON LIFETIME: BEAM TECH.

- QED: Time of Flight Method

- Lifetime:

$$\tau_n = 885.7 \pm 2.0 \text{ s}$$



Vol 444 | 21/28 December 2006 | doi:10.1038/nature05390

nature

LETTERS

Observation of the radiative decay mode of the free neutron

Jeffrey S. Nico¹, Maynard S. Dewey¹, Thomas R. Gentile¹, H. Pieter Mumm¹, Alan K. Thompson¹, Brian M. Fisher², Isaac Kremsky², Fred E. Wietfeldt², Timothy E. Chupp³, Robert L. Cooper³, Elizabeth J. Beise⁴, Kristin G. Kiriluk⁴, James Byrne⁵ & Kevin J. Coakley⁶

The theory of quantum electrodynamics (QED) predicts that beta decay of the neutron into a proton, electron and antineutrino should be accompanied by a continuous spectrum of soft photons. While this inner bremsstrahlung branch has been previously measured in nuclear beta and electron capture decay, it has never been observed in free neutron decay. Recently, the photon energy spectrum and branching ratio for neutron radiative decay have been calculated using two approaches: a standard QED framework^{1–3} and heavy baryon chiral perturbation theory⁴ (an effective theory of hadrons based on the symmetries of quantum chromodynamics). The QED calculation treats the nucleons as point-like, whereas the latter approach includes the effect of nucleon structure in a systematic way. Here we observe the radiative decay mode of free neutrons, measuring photons in coincidence with both the emitted electron and proton. We determined a branching ratio of $(3.13 \pm 0.34) \times 10^{-3}$ (68 per cent level of confidence) in the energy region between 15 and 340 keV, where the uncertainty is dominated by systematic effects. The value is consistent with the predictions of both theoretical approaches; the characteristic energy spectrum of the radiated photons, which differs from the uncorrelated background spectrum, is also consistent with the calculated spectrum. This result may provide opportunities for more detailed investigations of the weak interaction processes involved in neutron beta decay.

The experimental setup for measuring the radiative decay of the neutron beam is shown in Figure 1. The neutron beam enters from the left and passes through a series of components: a B-field line, shielding, and a photon detector. The beam then passes through an electrostatic mirror and an electron/proton detector. The distance between the photon detector and the electron/proton detector is 50 cm.





Figure 1 | Detection scheme for measuring the radiative decay of the


FRUSTRATED TOTAL REFLECTION.




Resonance order ν	Resonance wavelength $\lambda_{res} (\text{\AA})$	$\Delta\lambda_{res}$ ($\text{\AA}$)	k ($\text{\AA}^{-1}$)	Δk ($10^{-2} \text{\AA}^{-1}$)	Trapping time τ (ps)
1	17.38	1.09	3.561	2.267	19.35
2	12.99	0.85	4.837	3.165	10.36
3	10.56	0.73	5.950	4.113	6.48
4	9.02	0.61	6.966	4.711	4.83

The evolution can be reasonably described by a quadratic curve $T_{\text{trapping}} \approx -0.084\lambda + 0.7\lambda^2$. Hence, by extrapolation, for ultra-cold neutrons with wavelength of 100 or 1000 nm, T_{trapping} would be of 7 μs and 700,084 μs

ULTRA-COLD NEUTRON SOURCES



OUTLINE

1-Wave-Matter Quantum
Neutron Optics,



2A-Trapping neutron in Single Fabry-Perot
nanoresonator Heisenberg uncertainty governed
precision: Neutron lifetime considerations,


2B-Free propagation of neutrons in coupled Fabry-
Perot nanoresonators with ^{10}B

2C- γ -Radiolysis: From Maxwell to Nanofluids


4- Conclusion/Foresight

F.P. NANO-RESONATORS COUPLING

$\Theta_i \approx 1 \text{ Deg}$



Neutron
capture:
 B_4C




NEUTRON WAVE-PACKET RESONANCE: BORON TRANSPARENCY





- Most important neutron absorbers: ^{10}B as $^{10}\text{B}_4\text{C}$ in control rods, or **Boric acid** as a coolant water additive in PWRs.
- Other important neutron absorbers that are used in nuclear reactors **Xe, Cd, Hf, Gd, Co, Sm,, Dy**, all of which usually consist of mixtures of various isotopes—some of which are excellent neutron-absorbers. These also occur in combinations such as **Mo_2B_5 , HfB_2 , TiB_2** ,

BORON ABSORPTION CROSS SECTION: REACTORS' BARS CONTROL




- Most important neutron absorbers: ^{10}B as $^{10}\text{B}_4\text{C}$ in control rods, or Boric acid as a coolant water additive in PWRs.

BORON ABSORPTION CROSS SECTION: BORON NEUTRON CAPTURE THERAPY




BORON ABSORPTION CROSS SECTION: BF₄ NEUTRON DETECTORS



TMSC-TAIWAN & Nano-Processors.



NEUTRON WAVE-PACKET RESONANCE: BORON TRANSPARENCY



Bragg diffraction: $2 \Lambda \sin\Theta_i \approx k\lambda$

Kiessig & resonance condition:

$$L_{\text{total}} Q_{\text{max}} \approx k' 2 \pi$$

NEUTRON WAVE-PACKET RESONANCE: BORON TRANSPARENCY

Resonance conditions:
Zeldovich-Vinogradov equation (X-Rays)

$$\Gamma = \frac{\Delta t_H}{\Delta t_H + \Delta t_L} = \frac{\Delta t_H}{d} \quad (4.7)$$

$$\tan(\pi \Gamma_{\text{opt}}) = \pi \left[\Gamma_{\text{opt}} + \frac{\beta_L}{\beta_H - \beta_L} \right] \quad (4.8)$$


(Vinogradov and Zeldovich, 1977)

(also see Borrmann, 1941)





- Sharp interfaces needed for scattering
- Thin high-Z layer to minimize absorption
- Low-Z layer best as a "spacer"


©2014, ZeldovichReflex



NEUTRON WAVE-PACKET RESONANCE: BORON TRANSPARENCY



NEUTRON WAVE-PACKET RESONANCE: BORON TRANSPARENCY



NEUTRON WAVE-PACKET RESONANCE: BORON TRANSPARENCY

Untersuchungen zur Totalreflexion von Röntgenstrahlen¹⁾

Von Heinz Kiessig

(Mit 30 Figuren)

Eiuleitung

Nachdem der Nachweis erbracht worden war, daß auch Röntgenstrahlen sich meßbar brechen und reflektieren können, sind in den letzten Jahren zahlreiche Arbeiten gemacht worden, die sich mit dem quantitativen Studium dieser Erscheinungen befassen.²⁾ Das Ziel dieser Arbeiten war fast ausschließlich die Messung des Brechungsindex in Abhängigkeit von der Wellenlänge und den Materialien, um die Dispersionskurve für Röntgenstrahlen aufstellen zu können. Es zeigte sich, daß die Drude-Lorentzsche Dispersionstheorie nicht nur qualitativ anwendbar ist, sondern auch quantitativ die Verhältnisse gut wiedergibt. Da für Röntgenstrahlen der Brechungsindex n kleiner als 1 ist, aber nur wenig von 1 abweicht, setzt man üblicherweise

$$n = 1 - \delta$$

und gibt immer nur den Wert δ an. Es lautet dann die Lorentz-sche Dispersionsformel

$$\delta = \frac{e^2}{2\pi m} \sum \frac{N_K}{\nu^2 - \nu_K^2},$$

wobei

ν die Frequenz der einfallenden Strahlung,



ν_K die Eigenfrequenzen der Elektronen und

N_K die Anzahl der Elektronen mit der Eigenfrequenz ν_K in dem Einheitsvolumen bedeutet.




Die Summe ist über alle Eigenfrequenzen zu erstrecken.

1) Dissertation der Technischen Hochschule München.


2) Zusammenfassende Darstellungen über das vorliegende Material



NEUTRON WAVE-PACKET RESONANCE: BORON TRANSPARENCY



NEUTRON WAVE-PACKET RESONANCE: BORON TRANSPARENCY




$$\Theta_i \approx 1 \text{ Deg}$$




$$D_{eff} \approx D_{B4C} / \sin \Theta$$


$$D_{eff} \approx 200 \text{ nm}$$




ULTRA-COLD NEUTRON SOURCES



EUROPEAN SPALLATION SOURCE: ULTRACOLD NEUTRONS SOURCE



CHUKRAN,NGIYABONGA, ENKOSI,
MERCİ, DANKİE, KE A LEBOGA,
ASANTE SANA,THANK YOU



OUTLINE

1-Wave-Matter Quantum
Neutron Optics,



2A-Trapping neutron in Single Fabry-Perot
nanoresonator Heisenberg uncertainty governed
precision: Neutron lifetime considerations,

2B-Free propagation of neutrons in coupled Fabry-
Perot nanoresonators with ^{10}B

2C- γ -Radiolysis: From Maxwell to Nanofluids

4- Conclusion/Foresight

R&D NETWORKS/CONT.INFRASTRUCTURE:

HIGH ENERGY STEREOSCOPIC SYSTEM: "HESS"



2000 2002 2003 2005 2010 2011 2012




R&D NETWORKS/CONT.INFRASTRUCTURE:



GAMMA RADIATION DETECTOR: "GAMKA"






γ -RADIOLYSIS: FROM MAXWELL'S CONCEPT TO NANOFUIDS





James Clerk Maxwell, 1867



γ -RADIOLYSIS: FROM MAXWELL'S CONCEPT TO NANOFUIDS



γ -RADIOLYSIS: FROM MAXWELL'S CONCEPT TO NANOFUIDS



- σ_{thermal} EG: ~ 0.327 W/mK
- GNS-EG: $\sim 6.25\%$
- AgNPs-GNs: $\sim 12.5\%$
- AgNPs-GNs-EG: 33%

OUTLINE

1-Wave-Matter Quantum
Neutron Optics,



2A-Trapping neutron in Single Fabry-Perot
nanoresonator Heisenberg uncertainty governed
precision: Neutron lifetime considerations,

2B-Free propagation of neutrons in coupled Fabry-
Perot nanoresonators with ^{10}B

2C- γ -Radiolysis: From Maxwell to Nanofluids

4- Conclusion/**Foresight**

NEUTRON LIFETIME & S.M.

- **Lifetime:** $\langle \tau_n \rangle$ Used to predict the ratio of protons to helium atoms in the primordial universe and to search for physics beyond the Standard Model of particle physics. Relative abundance H/He During Universe 1st few mins:

$$n/p = \exp(-\Delta m / kT_{\text{freeze}}) \approx 1/6.$$

PARTICLE PHYSICS

Physicists close in on neutron puzzle

Researchers are narrowing down their measurements of how long the subatomic particle survives on its own.

BY ALEXANDRA WITZE IN DENVER, COLORADO

Physicists are drawing nearer to answering a long-standing mystery of the Universe: how long a neutron lives.

Neutrons are electrically neutral particles that usually combine with protons to make up atomic nuclei. Some neutrons are not bound up in atoms; these free-floating neutrons decay radioactively into other particles in minutes.

But physicists can't agree on precisely how long it takes a neutron to die. Using one laboratory approach, they measure the average neutron lifetime as 14 minutes 39 seconds. Using a different approach, they get 8 seconds longer.

"We don't know why they're different," says Shannon Hoogerheide, a physicist at the National Institute of Standards and Technology (NIST) in Gaithersburg, Maryland. "We really need to understand and eliminate this discrepancy." She and other scientists debated new ways to solve the problem this month at a meeting of the American Physical Society in Denver, Colorado.

Pinpointing the lifetime of a neutron is important for understanding how much hydrogen, helium and other light elements formed in the first few minutes after the Universe was born 13.8 billion years ago. Scientists also think that pinning down the neutron's lifetime would help to constrain measurements of other subatomic particles.

One way of clocking the neutron's lifespan is to put some of the particles in a bottle and count how many are left after a period of time. This 'bottle' method has been tried at several laboratories, including the Los Alamos National Laboratory in New Mexico¹ and the Institut Laue-Langevin in Grenoble, France. On average, they come up with a neutron lifetime of 14 minutes 39 seconds.


The other way is to feed neutrons into a detector that counts the protons created as the neutrons decay. This 'beam' method has been used at NIST and at the Japan Proton Accelerator Research Complex in Tokai. The Japanese work has just begun, but the NIST team reported in 2013 that its neutrons live eight seconds longer, on average, than those in the bottle method².

That's a big problem, because the beam and bottle measurements don't overlap, even when their margins of error are taken into account. So physicists have been looking for ways to explain why neutrons might be disappearing from bottles faster than from beams.

One possibility is that one of the two methods is doing something wrong. In that case, researchers might want to combine beam

442 | NATURE | VOL 568 | 25 APRIL 2019 | © 2019 Springer Nature Limited. All rights reserved.

ULTRA-COLD NEUTRON SOURCES



EUROPEAN SPALLATION SOURCE: BRITHNESS



EUROPEAN
SPALLATION
SOURCE

Partners:


- JÜLICH Forschungszentrum
- cea
- necsa
- NEUTRONS FOR SOCIETY
- ESS Bilbao
- Science & Technology Facilities Council
- UK Research and Innovation
- FRM II Forschungs-Neutronenquelle Heinz Maier-Leibnitz
- ess EUROPEAN SPALLATION SOURCE
- wigner
- TU WIEN TECHNISCHE UNIVERSITÄT WIEN
- NATIONAL CENTRE FOR NUCLEAR RESEARCH SWIER
- INFN Istituto Nazionale di Fisica Nucleare
- PAUL SCHERRER INSTITUT PSI
- NRf iThemba LABS

Quick facts

- Timeline**
Jan 2019 – Dec 2021
- Budget**
4 999 592.50 €
- 15 Partners**
- 11 Countries**

BrightnESS² is funded by the European Union Framework Programme for Research and Innovation Horizon 2020, under grant agreement 823867

EUROPEAN SPALLATION SOURCE: ULTRACOLD NEUTRONS SOURCE




Acknowledgments

- UNESCO , UNISA & Ithemba-LABS/NRF
- Abdus Salam International Centre for Theoretical Physics, Trieste-Italy.
- Department of Science & Technology of South Africa, Pretoria-South Africa.
- Ministry of Foreign Affairs, Roma-Italy.
- Science & Technology Directorate, French Embassy, Pretoria-South Africa.
- ELETTRA Synchrotron Facility, Trieste-Italy
- African Laser Centre, Pretoria-South Africa.
- African Union-Science & Technology Commission, Addis Ababa-Ethiopia.
- National Research Foundation of South Africa, Pretoria-South Africa.
- Academy of Sciences for the Developing World, Trieste-Italy.
- Organization of Women in Science for the Developing World, Trieste-Italy.
- International Centre for Science & Technology-UNIDO, Trieste-Italy.
- Centre National pour la Recherche Scientifique, Paris-France.
- The EU-FP7 ICPCNANONET, Brussels-Belgium.
- The National Institute for Materials Sciences NIMS, Tsukuba-Japan.
- Nelson Mandela African University of Science & Technology, Abuja-Nigeria.
- l' Oreal-UNESCO Foundation, Paris-France.
- University of South Africa.
- Islamic Academy of Sciences, Amman-Jordan.
- iThemba LABS, Western Cape-South Africa.



TYPICAL NEUTRON GUIDE HALL/ REFLECTOMER



FRUSTRATED TOTAL REFLECTION.

VOLUME 44, NUMBER 20

PHYSICAL REVIEW LETTERS

19 MAY 1980

Observation of Quasibound States of the Neutron in Matter

K.-A. Steinhauser, A. Steyerl, and H. Scheckenhofer

Fakultät für Physik, Technische Universität München, D-8046 Garching, Germany

and

S. S. Malik

*Fakultät für Physik, Technische Universität München, D-8046 Garching, Germany, and Physics Department,
University of Rhode Island, Kingston, Rhode Island 02881*

(Received 29 February 1980)

Quasistationary states of ultracold neutrons have been observed in a double-hump potential barrier. This potential barrier was created by sandwiching a thin film of aluminum between two thin copper films. Measurements of reflection from and transmission through such composite films displayed clear resonances. The resonance positions and widths are in agreement with calculations.

The average neutron lifetime derived from:


Beam approach was found to be $\langle \tau_n \rangle_{\text{Beam}} \sim 888.0 \pm 2.0 \text{ s}$

Bottle technique of about $\langle \tau_n \rangle_{\text{Bottle}} \sim 879.4 \pm 0.6 \text{ s}$

Steyerl & al, "Trapping in Fabry-Perot Nano-resonator" $\pm 10^{-6} \text{ s}$

Trapping in Fabry-Perot Nano-resonator" $\pm 10^{-12} \text{ s}$

Neutron Diamond Jubilee, U2ACN2, Sept.2022, Lund-Sweden, UCN-ESS2022





TYPICAL NEUTRON GUIDE HALL



TOTAL REFLECTION PHENOMENON

3A-NEUTRON OPTICS :TOTAL REFLECTION

- Quantum Mechanics governed phenomena
- Maximize the reactor capabilities via n-guides





NUCLEAR SCATTERING AMPLITUDE.

3A-NEUTRON OPTICS : TOTAL REFLECTION

$V_{\text{Fermi}} \approx 10^{-5} \text{ eV},$
 $n \approx 1$

Grazing incidence

$\theta_i \approx 1 \text{ Deg}$



NEUTRON LIFETIME & S.M.

- **Lifetime:** $\langle \tau_n \rangle$ Used to predict the ratio of protons to helium atoms in the primordial universe and to search for physics beyond the Standard Model of particle physics. Relative abundance H/He During Universe 1st few mins:

$$n/p = \exp(-\Delta m / kT_{\text{freeze}}) \approx 1/6.$$

PARTICLE PHYSICS

Physicists close in on neutron puzzle

Researchers are narrowing down their measurements of how long the subatomic particle survives on its own.

BY ALEXANDRA WITZE IN DENVER, COLORADO

Physicists are drawing nearer to answering a long-standing mystery of the Universe: how long a neutron lives.

Neutrons are electrically neutral particles that usually combine with protons to make up atomic nuclei. Some neutrons are not bound up in atoms; these free-floating neutrons decay radioactively into other particles in minutes.

But physicists can't agree on precisely how long it takes a neutron to die. Using one laboratory approach, they measure the average neutron lifetime as 14 minutes 39 seconds. Using a different approach, they get 8 seconds longer.

"We don't know why they're different," says Shannon Hoogerheide, a physicist at the National Institute of Standards and Technology (NIST) in Gaithersburg, Maryland. "We really need to understand and eliminate this discrepancy." She and other scientists debated new ways to solve the problem this month at a meeting of the American Physical Society in Denver, Colorado.

Pinpointing the lifetime of a neutron is important for understanding how much hydrogen, helium and other light elements formed in the first few minutes after the Universe was born 13.8 billion years ago. Scientists also think that pinning down the neutron's lifetime would help to constrain measurements of other subatomic particles.

One way of clocking the neutron's lifespan is to put some of the particles in a bottle and count how many are left after a period of time. This 'bottle' method has been tried at several laboratories, including the Los Alamos National Laboratory in New Mexico¹ and the Institut Laue-Langevin in Grenoble, France. On average, they come up with a neutron lifetime of 14 minutes 39 seconds.

The other way is to feed neutrons into a detector that counts the protons created as the neutrons decay. This 'beam' method has been used at NIST and at the Japan Proton Accelerator Research Complex in Tokai. The Japanese work has just begun, but the NIST team reported in 2013 that its neutrons live eight seconds longer, on average, than those in the bottle method².

That's a big problem, because the beam and bottle measurements don't overlap, even when their margins of error are taken into account. So physicists have been looking for ways to explain why neutrons might be disappearing from bottles faster than from beams.

One possibility is that one of the two methods is doing something wrong. In that case, researchers might want to combine beam

442 | NATURE | VOL 568 | 25 APRIL 2019 | © 2019 Springer Nature Limited. All rights reserved.