Editors' Suggestion

Warm Inflation with the Standard Model

Kim V. Berghaus,[1] Marco Drewes,[2,3] and Sebastian Zell[4,5,2]
[1]Walter Burke Institute for Theoretical Physics, California Institute of Technology, 1200 E California Blvd, Pasadena, California 91125, USA
[2]Centre for Cosmology, Particle Physics and Phenomenology (CP3), Université catholique de Louvain, Chemin du Cyclotron 2, B-1348 Louvan-la Neuve, Belgium
[3]Physik–Department, Technische Universität München, James Franck Straße 1, D-85748 Garching, Germany
[4]Arnold Sommerfeld Center, Ludwig-Maximilians-Universität, Theresienstraße 37, 80333 München, Germany
[5]Max-Planck-Institut für Physik, Boltzmannstraße 8, 85748 Garching bei München, Germany

(Received 6 May 2025; revised 19 August 2025; accepted 9 September 2025)

Q1 We show for the first time that warm inflation is feasible with standard model (SM) gauge interactions alone. Our model consists of a minimal extension of the SM by a single scalar inflaton field with an axionlike coupling to gluons and a monomial potential. The effects of light fermions, which were previously argued to render warm inflation with the SM impossible, are alleviated by Hubble dilution of their chiral chemical potentials. Our model features only one adjustable combination of parameters and accommodates all inflationary observables. We briefly discuss implications for axion experiments, dark matter, and the strong CP problem.

DOI: 10.1103/9nn9-bsm9

Introduction—Cosmic inflation [1–4], an early period of accelerated cosmic expansion, provides the leading explanation for the observed near-homogeneity and isotropy of our Universe. However, it remains unknown what mechanism drove inflation, and how it is connected to microphysical theories of particle physics. Many models can effectively be parametrized by a single scalar field ϕ, the potential of which, $V(\phi)$, dominated the cosmic energy budget. However, simple renormalizable potentials with a single parameter have long been ruled out by observations of the cosmic microwave background (CMB) [5,6]. A plethora of models with more complicated potentials exist [7], but they usually contain more parameters, making an observational discrimination impossible. Moreover, for many models, it is not clear how they can be realized microphysically and related to theories of particle physics, in particular, the standard model (SM).

Some of these issues can be addressed by *warm inflation* [8], a variant of the inflationary paradigm in which dissipative effects maintain the presence of a thermal bath during inflation; see, e.g., Ref. [9] for a review. This allows for accelerated expansion on steeper potentials, alleviating the long-standing difficulty of protecting the flatness of the inflaton potential from quantum corrections [10–14]. Moreover, thermal fluctuations provide an additional source of scalar perturbations in the metric, effectively suppressing the tensor-to-scalar ratio r and, hence, rescuing models that would be observationally excluded in their absence. Finally, non-Gaussianities with a unique bispectral shape [15,16] provide a pathway toward discovering evidence of inflation.

Arguably, the main challenge of warm inflation has been to embed it in a microphysical theory [17]. The fundamental problem is that, in perturbative computations, the inflaton dissipation coefficient and thermal correction to the effective potential originate from the imaginary and real parts of the same Feynman diagrams, making it difficult to enhance the former without introducing sizable corrections to the latter that spoil the flatness of the potential. This issue can be avoided if the dissipation arises from an axionlike coupling to the topological winding number in non-Abelian gauge theory, which gives rise to a new form of thermal friction [18] known as sphaleron heating. It has been shown that sphaleron heating can indeed support warm inflationary dynamics [19] within a consistent quantum field theory framework [20], but previous realizations of this idea generally require new (dark) non-Abelian gauge interactions that serve no other evident purpose.

Here, we demonstrate that warm inflation can be induced directly by an axionlike coupling between the inflaton and quantum chromodynamics (QCD), the strong force of the SM. QCD has not previously been considered as a viable candidate for the required non-Abelian force due to the suppression of sphaleron heating by axial chemical potentials μ_A that it generates for the quarks [18,21,22]. In the present proposal, this problem is alleviated by the Hubble dilution of μ_A [23], which has been neglected in earlier

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

works but is essential during inflation. We construct the first model of warm inflation that is driven by known SM gauge interactions alone. Our proposal is compatible with all observations and contains only two unknown parameters, one in the quartic inflaton potential and the other one in the coupling to gluons that is potentially experimentally testable.

Warm inflation—We consider inflation to be warm if there is a thermal bath with a temperature T that exceeds the Hubble rate H during a period of accelerated expansion ($T > H$). To achieve this, the inflaton field must source particles at a rate that can compensate for their Hubble dilution, and their self-interactions must be fast enough to maintain thermal equilibrium. This microphysical process can be captured by an effective description in which a thermal friction coefficient [24] Υ extracts energy from the slow-rolling inflaton field ϕ. The homogeneous equations of motion of the inflaton expectation value and radiation density $\rho_R \equiv (\pi^2/30) g_* T^4$ read

$$\ddot{\phi} + (3H + \Upsilon)\dot{\phi} + V' = 0, \tag{1}$$

$$\dot{\rho}_R + 4H\rho_R = \Upsilon \dot{\phi}^2. \tag{2}$$

Here, dots denote derivatives with respect to time, prime denotes derivative with respect to ϕ, $H \equiv \dot{a}/a$ is the Hubble expansion, with a the scale factor, and g_* corresponds to the number of relativistic degrees of freedom in the thermal bath. The cases $Q \equiv \Upsilon/(3H) > 1$ and $Q < 1$ are commonly referred to as the strong and weak regime of warm inflation, respectively. In both regimes, the thermal bath can contribute to sourcing cosmological perturbations, but only in the strong regime does thermal friction dominate the slow-roll dynamics.

The Friedman equation governs the inflationary dynamics, $H^2 = (V(\phi) + \rho_R + \frac{1}{2}\dot{\phi}^2)/(3M_{\rm pl}^2) \approx V(\phi)/(3M_{\rm pl}^2)$, where $M_{\rm pl} = 2.43 \times 10^{18}$ GeV is the reduced Planck mass. A sustained period of accelerated expansion requires $V(\phi) \gg \rho_R, \dot{\phi}^2$, which can be assured by the usual slow-roll conditions, $\epsilon_V, \eta_V \ll 1$, if the slow-roll parameters are modified to be [26]

$$\epsilon_V \equiv \frac{M_{\rm pl}^2}{2(1+Q)}\left(\frac{V'}{V}\right)^2, \qquad \eta_V \equiv \frac{M_{\rm pl}^2}{(1+Q)}\frac{V''}{V}. \tag{3}$$

From Eq. (3), it is immediately apparent that in the strong regime of warm inflation the slow-roll conditions are more easily satisfied than in its cold counterpart, implying that a given number of e-folds,

$$N_* = \int H dt = \int_{\phi_{\rm end}}^{\phi_*} \frac{1}{M_{\rm pl}^2}\frac{V}{V'}(1+Q) d\phi, \tag{4}$$

can be achieved with a smaller field value ϕ_* or steeper potential (in the sense of larger V'/V). The slow-roll conditions suppress the first two terms in Eqs. (1) and (2), such that

$$\frac{\pi^2}{30} g_* T^4 \approx \frac{\Upsilon}{4H}\left(\frac{V'}{(3H+\Upsilon)}\right)^2 \tag{5}$$

defines the approximately constant steady-state temperature of the thermal bath T.

Sphaleron heating with the SM—Our model is defined by the extension of the SM [27]

$$\mathcal{L} = \frac{1}{2}\partial^\mu\phi\partial_\mu\phi - V(\phi) - \frac{\alpha_s}{8\pi}\frac{\phi}{f}G^a_{\mu\nu}\tilde{G}^{a\mu\nu}, \tag{6}$$

where $G^a_{\mu\nu}$ is the QCD field strength tensor, $\tilde{G}^{a\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\rho\sigma}G^a_{\rho\sigma}$ its dual, and $\alpha_s \equiv g^2/4\pi$ with g the QCD gauge coupling. The following considerations, however, hold for a general Yang-Mills SU(N_c) gauge group. We first neglect the presence of quarks and include their impact further below. For T larger than the confinement temperature, the equation of motion for the expectation value of ϕ reads [18]

$$\ddot{\phi} + 3H\dot{\phi} + V' = -\frac{\alpha_s}{8\pi f}\langle G^a_{\mu\nu}\tilde{G}^{a\mu\nu}\rangle. \tag{7}$$

The thermal average of $G^a_{\mu\nu}\tilde{G}^{a\mu\nu}$ induces a thermal friction coefficient $\Upsilon_{\rm sph}$ due to real-time sphaleron processes that lead to biased transitions between vacua with different Chern-Simons number $N_{\rm CS}$, driven by the inflaton velocity $\dot{\phi}$. The friction coefficient $\Upsilon_{\rm sph}$ is related to the SU(N_c) sphaleron rate $\Gamma_{\rm sph} \equiv \lim_{t\to\infty}(N_{\rm CS}(t) - N_{\rm CS}(0))^2/(Vt)$ (also called the Chern-Simons diffusion rate) via the fluctuation-dissipation theorem [16,18–20,28]:

$$\frac{\alpha_s}{8\pi f}\langle G^a_{\mu\nu}\tilde{G}^{a\mu\nu}\rangle \equiv \Upsilon_{\rm sph}\dot{\phi} = \frac{\Gamma_{\rm sph}}{2Tf}\frac{\dot{\phi}}{f}. \tag{8}$$

By comparing Eqs. (7) and (8) to Eq. (1), we can identify $\Upsilon = \Upsilon_{\rm sph}$. Using the estimate $\Gamma_{\rm sph} \approx \alpha_s^5 N_c^5 T^4$ [29], one finds [28]

$$\Upsilon_{\rm sph} \simeq (\alpha_s N_c)^5 \frac{T^3}{2f^2}, \tag{9}$$

where we neglected an additional prefactor of the order of 1 with a weak dependence on model parameters and dynamical quantities (see Refs. [20,29–31]).

The perturbative shift symmetry of the axionlike coupling in Eq. (6) protects the inflaton potential from thermal corrections [16,18–20,31,32] that would otherwise endanger the slow-roll dynamics [17]. However, since nonperturbative contributions to $V(\phi)$ from instantons vanish at temperatures much larger than the QCD confinement scale $\Lambda_{\rm QCD}$, where $\alpha_s < 1$ (see, e.g., Ref. [33]),

the coupling to QCD cannot generate a potential that drives inflation [34].

The relevant interaction with SM quarks ψ is given by

$$\mathcal{L}_{\rm QCD} \supset \sum_f \left(\frac{i}{2} \bar{\psi}_f \not{D} \psi_f + \text{H.c.} \right), \tag{10}$$

where f denotes the sum over SM flavors. The chiral anomaly of SU(N_c) leads to a nonconservation of the axial fermion current $j^\mu_A = \sum_f \bar{\psi}_f \gamma^\mu \gamma^5 \psi_f$ [18,36]:

$$\partial_t j^0_A = -N_f \left\langle \frac{\alpha_s G^a_{\mu\nu} \tilde{G}^{a\mu\nu}}{4\pi} \right\rangle, \tag{11}$$

where the axial charge density j^0_A corresponds to the difference in number density between right-handed and left-handed Dirac fermions. The quarks backreact onto the biased sphaleron transitions such that Eq. (8) changes to [18,29,37–41]

$$\frac{\alpha_s}{8\pi f} \langle G^a_{\mu\nu} \tilde{G}^{a\mu\nu} \rangle = \Upsilon_{\rm sph} \left(\dot{\phi} + \frac{6 f j^0_A}{N_c T^2} \right). \tag{12}$$

In particular, Eqs. (11) and (12) imply that, in the absence of a rolling axion, sphalerons relax the axial charge as $\partial_t j^0_A = -6(N_f/N_c)(\Gamma_{\rm sph}/T^3) j^0_A$ [29,37–41].

We define a chemical potential μ_A via

$$\begin{aligned} j^0_A &= 2 N_c N_f \int \frac{d^3 p}{(2\pi)^3} \left(\frac{1}{e^{(p-\mu_A)/T} + 1} - \frac{1}{e^{(p+\mu_A)/T} + 1} \right) \\ &= \frac{N_c N_f \mu_A T^2}{3}, \end{aligned} \tag{13}$$

where N_f denote the number of flavors, and we have taken the $\mu_A \ll T$ limit in the second line [42]. Using Eqs. (12) and (7) and the anomaly relation (11), we obtain the coupled evolution of the inflaton and the axial chemical potential for a quasi-steady-state temperature in an expanding universe [18]:

$$\ddot{\phi} + 3H\dot{\phi} + V' = -\Upsilon_{\rm sph} (\dot{\phi} + 2 N_f f \mu_A), \tag{14}$$

$$\dot{\mu}_A + 3H\mu_A = -\frac{6 f \Upsilon_{\rm sph}}{N_c T^2} (\dot{\phi} + 2 N_f f \mu_A). \tag{15}$$

Importantly, here we also take into account Hubble dilution of the chemical potential in the second line. When Hubble friction is neglected, there is no solution where $\ddot{\phi}$ vanishes asymptotically [18]; hence, slow roll cannot be sustained [19,21]. In a quasistatic situation, the effect of H can be incorporated in Eq. (14) by absorbing the term $\Upsilon_{\rm sph} 2 N_f f \mu_A$ into an effective dissipation coefficient [23]:

$$\Upsilon_{\rm eff} = \Upsilon_{\rm sph} \Big/ \left(1 + \frac{\Upsilon_{\rm sph}}{3H} \frac{N_f}{N_c} \frac{12 f^2}{T^2} \right). \tag{16}$$

Sphaleron heating is fully restored if $H/\Upsilon_{\rm eff} \gg 4 f^2 N_f / (T^2 N_c)$; i.e., the chemical potential is diluted faster than it builds up due to sphaleron transitions. Achieving this is generically easier in the weak regime of warm inflation where $\Upsilon_{\rm sph} < 3H$. Using the Friedmann equation and Eq. (9), we get

$$\Upsilon_{\rm eff} = N_c^5 \alpha_s^5 \frac{T^3}{2 f^2} \Big/ \left(1 + \frac{2 N_f N_c^4 \alpha_s^5 T}{\sqrt{V(\phi)/(3 M_{\rm pl}^2)}} \right). \tag{17}$$

Note that Eq. (17) can yield different functional dependencies of $\Upsilon_{\rm eff}$ on ϕ and T if the second term dominates the denominator, depending on the shape of $V(\phi)$:


$$\Upsilon_{\rm eff} \propto \frac{T^2}{f^2} \frac{\sqrt{V(\phi)}}{M_{\rm pl}}. \tag{18}$$

In addition to the gauge interactions in (10), quarks also have Yukawa couplings, $y_f H \overline{\psi_f} \psi_f$, which mediate chirality violation processes either by generating fermion masses [43] or through thermal scatterings [44,45]. Both make the SM flavors distinguishable. An equation of motion for each chemical potential μ_{A_f} can be obtained from (15) by omitting the factor N_f and adding a term $-\Gamma_{{\rm ch}f} \mu_A$ on the rhs, which can effectively be incorporated into $\Upsilon_{\rm eff}$ by expressing the factor N_f terms of a sum over flavors with the replacement $3H \to 3H + \Gamma_{{\rm ch}f}$. If all $\Gamma_{{\rm ch}f}$ are either much smaller or much larger than H and $\Upsilon_{\rm sph}$, one can simply continue using the unflavored equation (16) but set N_f to the number of SM flavors for which $\Gamma_{{\rm ch}f} \ll \Upsilon_{\rm sph}$, which for $\Gamma_{{\rm ch}f} \sim 10^{-2} y_f^2 T$ [44] amounts to $y_f^2 \lesssim 10^3 \alpha_s^5 N_c^4$. This condition is violated only for the top quark; hence, we use (16) with $N_f = 5$ [46]. We further set $N_c = 3$ and obtain α_s from running to the energy scale T with the known SM β functions [47,48].

Working model—We choose the quartic potential

$$V(\phi) = \lambda \phi^4 \tag{19}$$

to fully determine the Lagrangian (6) with two unknown model parameters f and λ. Matching the observed amplitude $A_s(k_*) \approx 2 \times 10^{-9}$ of scalar perturbations $\Delta_s^2(k) = A_s(k_*)(k/k_*)^{n_s(k_*)-1}$ at the pivot scale $k_* = 0.05\ \text{Mpc}^{-1}$ [5] establishes a relation between them, effectively leaving us with one single free parameter. Practically, we use Eqs. (4) and (5), with $\Upsilon = \Upsilon_{\rm eff}$ and (19) to numerically solve for the temperature T and, subsequently, $\Upsilon_{\rm eff}$ as a function of ϕ, λ, and f. Varying λ scans along values of $Q = \Upsilon_{\rm eff}/(3H)$. With this and the approximation of instantaneous reheating,



F1:1 FIG. 1. The predictions of warm inflation with the SM for a
F1:2 quartic inflaton potential. The allowed 68% and 95% contour
F1:3 regions correspond to the constraints from the combined analysis
F1:4 of Planck [5] and BICEP/Keck [6] data. Proposed and upcoming
F1:5 CMB experiments will improve this bound by about a factor of 4
F1:6 (AliCPT [51,52], Simons Observatory [53], and Simons Array
F1:7 [54]) to 20 (LiteBIRD [55] and CMB-S4 [56]).

we can compute N_*, A_s, n_s, and the tensor-to-scalar ratio $r \equiv \Delta_h^2/\Delta_s^2$ from the relations given in the Appendix. Since our model effectively has only one free parameter, we predict a line in the $n_s - r$ plane, as displayed in Fig. 1.

We find that our model is compatible with observations in both the strong regime and the weak regime, consistent with what was previously found for dark gauge groups [16]. For $Q \gg 1$, the denominator in Eq. (17) suppresses the efficiency of sphaleron heating ($\Upsilon_{\rm eff} \ll \Upsilon_{\rm sph}$), and $\Upsilon_{\rm eff}$ follows the scaling in Eq. (18). We do not compare to CMB observables in this regime, as we can no longer compute perturbations reliably with the formulas given in the Appendix. This suppression is, however, already alleviated for $Q \lesssim \mathcal{O}(10)$, i.e., within the strong regime. Column 7 in Table I shows that the deviation of (17) from $\Upsilon_{\rm sph}$ amounts to less than 10%. For smaller Q, the restoration becomes increasingly efficient, so that $\Upsilon_{\rm eff}$ further approaches $\Upsilon_{\rm sph}$ in the weak regime. Thermally produced cosmological perturbations become negligible for $Q < 10^{-4}$, practically recovering standard cold inflation. The predicted warm inflation temperatures lie below $T \lesssim 6 \times 10^{14}$ GeV, with temperatures decreasing for larger values of Q. The corresponding quartic couplings for which inflation is warm lie below $\lambda \lesssim 10^{-13}$. We provide a comprehensive list of all the relevant parameters discussed here in Table I.

In our model, the interaction of the inflaton ϕ with the SM is already fully specified. Therefore, the transition from inflation to radiation domination is calculable, and, in the strong regime, accelerated expansion directly evolves to a period dominated by a quark-gluon plasma. This is reflected in the fact that we can uniquely predict r as a function of n_s (see Fig. 1), with no additional uncertainty arising from unknown interactions of the inflaton with the SM. Moreover, the direct coupling of the inflaton to QCD can lead to observable signatures at low energies [57]. The values for the coupling $f \sim 10^{10}$–10^{13} GeV lie within reach of proposed searches for axionlike particles [62,63]. However, Hubble dilution alone suffices to ensure that $\phi \ll f$ at the temperature of the QCD phase transition, implying that ϕ oscillations around the minimum of the potential $V(\phi) \simeq \lambda\phi^4 + \Lambda_{\rm QCD}^4(1 - \cos(\phi/f - \theta))$ contribute only a small fraction to the dark matter density, where θ is the CP-violating angle in QCD. Since $\lambda\phi^4$ still gives an important contribution to the potential at low energies, satisfying the experimental bound on CP violation in the strong interaction requires $|\phi/f - \theta| \ll 10^{-10}$ at the minimum, i.e., the same amount of tuning as in QCD, and can potentially be achieved with a standard QCD axion [64–66].

Conclusion—We showed for the first time that warm inflation can be driven by sphaleron heating from SM gauge interactions alone. Our model comprises a minimal extension of the SM by a single scalar inflaton field ϕ with an axionlike coupling to gluons (6). A crucial point is that the suppressing effect which light fermions have on the efficiency of sphaleron heating is partially alleviated by the expansion of the Universe, leading to the effective friction coefficient in Eq. (16). Notably, this coefficient can have different functional dependencies on the inflaton field value for different effective potentials in Eq. (18). Additionally, it predicts a smooth transition from a warm inflationary period to a Universe dominated by a quark-gluon plasma. For a monomial quartic potential (19), the model contains only two parameters and is consistent with CMB observations as well as laboratory experiments. Its minimality in terms of both the potential during inflation and the clear connection to the SM make this scenario highly predictive and a target for tests with future CMB and large-scale structure surveys as well as laboratory experiments, offering a unique opportunity to probe the connection between cosmic inflation and particle physics.

Note added—Recently, CMB predictions of our model were further investigated in [67,68]. Both works confirm that our proposal is compatible with current observational bounds.

Acknowledgments—We are grateful to Mikko Laine, Alica Rogelj, and Mikhail Shaposhnikov for their valuable comments on the manuscript. K. V. B. thanks the U.S. Department of Energy, Office of Science, Office of High Energy Physics, under Award No. DE-SC0011632 and the Walter Burke Institute for Theoretical Physics. The work of M. D. was partially funded by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation)—SFB 1258–283604770. The work of S. Z. was supported by the European Research Council Gravites Horizon Grant AO No. 850 173-6. Funded by the European Union.

Views and opinions expressed are, however, those of the authors only and do not necessarily reflect those of the European Union or European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

Data availability—The data that support the findings of this article are not publicly available. The data are available from the authors upon reasonable request.

[1] Alexei A. Starobinsky, A new type of isotropic cosmological models without singularity, Phys. Lett. B **91**, 99 (1980).
[2] Alan H. Guth, The inflationary universe: A possible solution to the horizon and flatness problems, Phys. Rev. D **23**, 347 (1981).
[3] Andrei D. Linde, A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems, Phys. Lett. B **108**, 389 (1982).
[4] Viatcheslav F. Mukhanov and G. V. Chibisov, Quantum fluctuations and a nonsingular universe, JETP Lett. **33**, 532 (1981), http://jetpletters.ru/ps/1510/article_23079.pdf.
[5] Y. Akrami *et al.* (Planck Collaboration), Planck 2018 results. X. Constraints on inflation, Astron. Astrophys. **641**, A10 (2020).
[6] P. A. R. Ade *et al.* (BICEP and Keck Collaborations), Improved constraints on primordial gravitational waves using Planck, WMAP, and BICEP/Keck observations through the 2018 observing season, Phys. Rev. Lett. **127**, 151301 (2021).
[7] Jerome Martin, Christophe Ringeval, and Vincent Vennin, Encyclopædia inflationaris: Opiparous edition, Phys. Dark Universe **5–6**, 75 (2014).
[8] Arjun Berera, Warm inflation, Phys. Rev. Lett. **75**, 3218 (1995).
[9] Vahid Kamali, Meysam Motaharfar, and Rudnei O. Ramos, Recent developments in warm inflation, Universe **9**, 124 (2023).
[10] Edmund J. Copeland, Andrew R. Liddle, David H. Lyth, Ewan D. Stewart, and David Wands, False vacuum inflation with Einstein gravity, Phys. Rev. D **49**, 6410 (1994).
[11] P. Binetruy and G. R. Dvali, D term inflation, Phys. Lett. B **388**, 241 (1996).
[12] G. R. Dvali and S. H. Henry Tye, Brane inflation, Phys. Lett. B **450**, 72 (1999).
[13] Nima Arkani-Hamed, Hsin-Chia Cheng, Paolo Creminelli, and Lisa Randall, Pseudonatural inflation, J. Cosmol. Astropart. Phys. 07 (2003) 003.
[14] Daniel Baumann and Liam McAllister, *Inflation and String Theory*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England, 2015).
[15] Mar Bastero-Gil, Arjun Berera, Ian G. Moss, and Rudnei O. Ramos, Theory of non-Gaussianity in warm inflation, J. Cosmol. Astropart. Phys. 12 (2014) 008.
[16] Mehrdad Mirbabayi and Andrei Gruzinov, Shapes of non-Gaussianity in warm inflation, J. Cosmol. Astropart. Phys. 02 (2023) 012.
[17] Junichi Yokoyama and Andrei D. Linde, Is warm inflation possible?, Phys. Rev. D **60**, 083509 (1999).
[18] Larry D. McLerran, Emil Mottola, and Mikhail E. Shaposhnikov, Sphalerons and axion dynamics in high temperature QCD, Phys. Rev. D **43**, 2027 (1991).
[19] Kim V. Berghaus, Peter W. Graham, and David E. Kaplan, Minimal warm inflation, J. Cosmol. Astropart. Phys. 03 (2020) 034; 10 (2023) E02.
[20] M. Laine and S. Procacci, Minimal warm inflation with complete medium response, J. Cosmol. Astropart. Phys. 06 (2021) 031.
[21] Kim V. Berghaus, Peter W. Graham, David E. Kaplan, Guy D. Moore, and Surjeet Rajendran, Dark energy radiation, Phys. Rev. D **104**, 083520 (2021).
[22] Kim V. Berghaus, Matthew Forslund, and Mark Vincent Guevarra, Warm inflation with a heavy QCD axion, J. Cosmol. Astropart. Phys. 10 (2024) 103.
[23] Marco Drewes and Sebastian Zell, On sphaleron heating in the presence of fermions, J. Cosmol. Astropart. Phys. 06 (2024) 038.
[24] See, e.g., Refs. [18,20,25], and references therein for derivations within quantum field theory.
[25] Gilles Buldgen, Marco Drewes, Jin U. Kang, and Ui Ri Mun, General Markovian equation for scalar fields in a slowly evolving background, J. Cosmol. Astropart. Phys. 05 (2022) 039.
[26] Arjun Berera, Ian G. Moss, and Rudnei O. Ramos, Warm inflation and its microphysical basis, Rep. Prog. Phys. **72**, 026901 (2009).
[27] The inflaton-gluon interaction is mediated by a dimension-five operator and, therefore, must be viewed as an effective field theory (EFT) with a cutoff $\Lambda_{\rm cutoff} \equiv (\alpha_s/8\pi f)^{-1}$. This restricts the validity of our computations to the regime $T < \Lambda_{\rm cutoff}$. We verify *a posteriori* that all temperatures in Table I lie at least one order of magnitude below this EFT cutoff.
[28] Mikko Laine and Aleksi Vuorinen, *Basics of Thermal Field Theory* (Springer, New York, 2016), Vol. 925.
[29] Guy D. Moore and Marcus Tassler, The sphaleron rate in SU(N) gauge theory, J. High Energy Phys. 02 (2011) 105.
[30] M. Laine, L. Niemi, S. Procacci, and K. Rummukainen, Shape of the hot topological charge density spectral function, J. High Energy Phys. 11 (2022) 126.
[31] P. Klose, M. Laine, and S. Procacci, Gravitational wave background from vacuum and thermal fluctuations during axion-like inflation, J. Cosmol. Astropart. Phys. 12 (2022) 020.
[32] H. Kolesova, M. Laine, and S. Procacci, Maximal temperature of strongly-coupled dark sectors, J. High Energy Phys. 05 (2023) 239.
[33] Sidney Coleman, *Aspects of Symmetry: Selected Erice Lectures* (Cambridge University Press, Cambridge, England, 1985).

[34] More generally, successful warm inflation in the strong regime requires a potential that breaks the symmetry $\phi \to \phi + 2\pi f$ of the axionlike coupling in Eq. (6) [35].

[35] Sebastian Zell, No warm inflation from sphaleron heating with a vanilla axion, arXiv:2408.07746.

[36] Michael E. Peskin and Daniel V. Schroeder, *An Introduction to Quantum Field Theory* (Addison-Wesley, Reading, MA, 1995).

[37] S. Yu. Khlebnikov and M. E. Shaposhnikov, The statistical theory of anomalous fermion number nonconservation, Nucl. Phys. **B308**, 885 (1988).

[38] G. F. Giudice and Mikhail E. Shaposhnikov, Strong sphalerons and electroweak baryogenesis, Phys. Lett. B **326**, 118 (1994).

[39] V. A. Rubakov and M. E. Shaposhnikov, Electroweak baryon number nonconservation in the early universe and in high-energy collisions, Usp. Fiz. Nauk **166**, 493 (1996).

[40] Guy D. Moore, Motion of Chern-Simons number at high temperatures under a chemical potential, Nucl. Phys. **B480**, 657 (1996).

[41] Guy D. Moore, Computing the strong sphaleron rate, Phys. Lett. B **412**, 359 (1997).

[42] In terms of this chemical potential, we see that $\partial_t j^0_A$ is independent of N_c but scales with $N_f^2 \mu_A$. A heuristic explanation of this scaling is that the production rate of fermions, which is caused by sphaleron transitions and violates chiral symmetry, is proportional to the free energy $N_f \mu_A$ liberated by a sphaleron event and, in addition, each event changes j^0_A by N_f [41].

[43] A. Boyarsky, V. Cheianov, O. Ruchayskiy, and O. Sobol, Evolution of the primordial axial charge across cosmic times, Phys. Rev. Lett. **126**, 021801 (2021).

[44] Dietrich Bödeker and Dennis Schröder, Equilibration of right-handed electrons, J. Cosmol. Astropart. Phys. 05 (2019) 010.

[45] Moreover, SU(2) sphalerons provide an additional source of chirality violation [23], but this effect turns out to be negligible since $(N_c \alpha_s)^5 \gg (2\alpha_w)^5$, where α_w is the weak fine structure constant.

[46] For the bottom quark $\Gamma_{\mathrm{ch}f} \simeq \Upsilon_{\mathrm{sph}}$, provided α_s runs to sufficiently low values.

[47] Lohan Sartore and Ingo Schienbein, PyR@TE 3, Comput. Phys. Commun. **261**, 107819 (2021).

[48] Thermal corrections to the vacuum β functions are expected to be negligible at temperatures greatly exceeding the QCD confinement scale [49,50].

[49] Olaf Kaczmarek and Felix Zantow, Static quark anti-quark interactions in zero and finite temperature QCD. I. Heavy quark free energies, running coupling and quarkonium binding, Phys. Rev. D **71**, 114510 (2005).

[50] Jens Braun and Holger Gies, Running coupling at finite temperature and chiral symmetry restoration in QCD, Phys. Lett. B **645**, 53 (2007).

[51] Hong Li *et al.*, Probing primordial gravitational waves: Ali CMB polarization telescope, Natl. Sci. Rev. **6**, 145 (2019).

[52] Yang Liu, Lei Ming, Marco Drewes, and Hong Li, AliCPT sensitivity to cosmic reheating, arXiv:2503.21207.

[53] Peter Ade *et al.* (Simons Observatory Collaboration), The Simons Observatory: Science goals and forecasts, J. Cosmol. Astropart. Phys. 02 (2019) 056.

[54] A. Suzuki *et al.* (POLARBEAR Collaboration), The POLARBEAR-2 and the simons array experiment, J. Low Temp. Phys. **184**, 805 (2016).

[55] E. Allys *et al.* (LiteBIRD Collaboration), Probing cosmic inflation with the LiteBIRD cosmic microwave background polarization survey, Prog. Theor. Exp. Phys. **2023**, 042F01 (2023).

[56] Kevork Abazajian *et al.* (CMB-S4 Collaboration), CMB-S4: Forecasting constraints on primordial gravitational waves, Astrophys. J. **926**, 54 (2022).

[57] In contrast, the connection between cold inflation and particle physics can be probed only indirectly [58,59], with the notable exception of Higgs inflation [60] (see, e.g., Ref. [61]).

[58] Marco Drewes, Measuring the inflaton coupling in the CMB, J. Cosmol. Astropart. Phys. 09 (2022) 069.

[59] Marco Drewes and Lei Ming, Connecting cosmic inflation to particle physics with LiteBIRD, CMB-S4, EUCLID, and SKA, Phys. Rev. Lett. **133**, 031001 (2024).

[60] Fedor L. Bezrukov and Mikhail Shaposhnikov, The Standard Model Higgs boson as the inflaton, Phys. Lett. B **659**, 703 (2008).

[61] Mikhail Shaposhnikov, Andrey Shkerin, and Sebastian Zell, Quantum effects in Palatini Higgs inflation, J. Cosmol. Astropart. Phys. 07 (2020) 064.

[62] C. Antel *et al.*, Feebly-interacting particles: FIPs 2022 workshop report, Eur. Phys. J. C **83**, 1122 (2023).

[63] Ciaran O'Hare, cajohare/axionlimits: Axionlimits, https://cajohare.github.io/AxionLimits/ (2020).

[64] R. D. Peccei and Helen R. Quinn, *CP* conservation in the presence of instantons, Phys. Rev. Lett. **38**, 1440 (1977).

[65] Steven Weinberg, A new light boson?, Phys. Rev. Lett. **40**, 223 (1978).

[66] Frank Wilczek, Problem of strong *P* and *T* invariance in the presence of instantons, Phys. Rev. Lett. **40**, 279 (1978).

[67] Rudnei O. Ramos and Gabriel S. Rodrigues, Viability of warm inflation with Standard Model interactions, Phys. Rev. D **111**, 123527 (2025).

[68] M. Laine, S. Procacci, and A. Rogelj, Evolution of coupled scalar perturbations through smooth reheating. II. Thermal fluctuation regime, arXiv:2507.12849.

[69] Chris Graham and Ian G. Moss, Density fluctuations from warm inflation, J. Cosmol. Astropart. Phys. 07 (2009) 013.

[70] Mar Bastero-Gil, Arjun Berera, and Rudnei O. Ramos, Shear viscous effects on the primordial power spectrum from warm inflation, J. Cosmol. Astropart. Phys. 07 (2011) 030.

[71] Gabriele Montefalcone, Vikas Aragam, Luca Visinelli, and Katherine Freese, WarmSPy: A numerical study of cosmological perturbations in warm inflation, J. Cosmol. Astropart. Phys. 01 (2024) 032.

[72] M. Laine, S. Procacci, and A. Rogelj, Evolution of coupled scalar perturbations through smooth reheating. Part I. Dissipative regime, J. Cosmol. Astropart. Phys. 10 (2024) 040.

[73] Gabriel S. Rodrigues and Rudnei O. Ramos, WI2easy: Warm inflation dynamics made easy, J. Cosmol. Astropart. Phys. 09 (2025) 014.

[74] Javier Rubio and Eemeli S. Tomberg, Preheating in Palatini Higgs inflation, J. Cosmol. Astropart. Phys. 04 (2019) 021.

[75] Lars Husdal, On effective degrees of freedom in the early universe, Galaxies **4**, 78 (2016).

End Matter

Appendix—The spectrum of cosmic perturbations for warm inflation has not been derived within quantum field theory at the same level of rigor as for standard inflation, and the predictions presented in the literature exhibit minor discrepancies [16,22,67–73]. However, our main conclusions are not sensitive to these differences. We use the predictions for A_s from [16] (recently confirmed by [68]):

$$\Delta_s^2(\phi_*) = \frac{H^2(\phi_*)}{4\pi^2\dot{\phi}^2(\phi_*)}(H(\phi_*)^2 + H(\phi_*)T(\phi_*)F[Q(\phi_*)]), \tag{A1}$$

where [16,22]

$$F(Q_*) = 3.24\times10^{-4}Q_*^7 + 168Q_*\left(\frac{1}{3}\left(1+\frac{9Q_*^2}{25}\right)+\frac{2}{3}\tanh\frac{1}{30Q_*}\right). \tag{A2}$$

The resulting scalar index $n_s - 1 \equiv d\ln\Delta_s^2/dN$ cannot be written in closed form in all generality for the regime in which Q transitions from the weak to the strong regime, and we, thus, calculate it numerically.

The scalar-to-tensor ratio is

$$r = \frac{8\dot{\phi}_*^2}{M_{\rm pl}^2 H_* T_* F(Q_*)}. \tag{A3}$$

Equation (A2) was derived for $\Upsilon_{\rm sph}$ rather than $\Upsilon_{\rm eff}$, for which the temperature no longer scales as a simple power law. This is a subtle complication, as it has been shown that the scalar perturbations are quite sensitive to the scaling with temperature in the friction coefficient for $Q > 1$ [69,70]. However, in the absence of calculations of the scalar power spectrum given $\Upsilon_{\rm eff}$, we proceed with Eq. (A1) as the best available approximation, and column 7 in Table I confirms *a posteriori* that the error is small.

Finally, the number of inflationary e-foldings after the generation of CMB perturbations is (cf. [74])

$$N_* = \ln\left(\left(\frac{g_{*0}}{g_*}\right)^{1/3}\frac{T_0}{T_{RH}}\frac{H_*}{k_*}\right), \tag{A4}$$

to be matched with Eq. (4). Here, $g_{*0} = 3.94$ [75] is the effective number of degrees of freedom today, $T_0 \approx 2.7$ K corresponds to the present CMB temperature, H_* is the Hubble scale at CMB generation, and we take $k_* = 0.05\ \mathrm{Mpc}^{-1}$ as the pivot scale. Given that the inflaton interactions are fully specified in our model, reheating is, in principle, calculable. In Eq. (A4), we have taken the transition from inflation to radiation domination as instantaneous and believe this to be a good approximation due to the strong coupling $\sim\alpha T/f$ with SM degrees of freedom. Consequently, we set T_{RH} to the temperature of the thermal plasma when $\epsilon_V(\phi_{\rm end}) = 1$.

TABLE I. Parameters of interest for warm SM inflation. λ, the potential slope, is a free parameter and scans the parameter space along Q. ϕ_* and f are fixed by N_*, and $A_s \approx 2\times10^{-9}$. All other parameters are derived. The EFT cutoff is $\Lambda_{\rm cutoff} \equiv (\alpha_s/8\pi f)^{-1}$. We present values to three significant figures to facilitate reproducibility when using Eqs. (9) and (A2). However, when accounting for theoretical uncertainties, we estimate that only one significant figure can be considered as reliable for the derived quantities (two figures for n_s, theoretical uncertainty not shown in Fig. 1).

λ	$(\phi_*/M_{\rm pl})$	$f[10^{12}$ GeV]	$T[10^{12}$ GeV]	α_s	T/H	$(2N_f N_c^4\alpha_s^5 T/\sqrt{\lambda\phi_*^4/3M_{\rm pl}^2})$	$T/\Lambda_{\rm cutoff}$	N_*	Q	n_s	r
10^{-21}	5.02	0.146	8.75	0.0269	7820	8.99×10^{-2}	0.064	52.8	14.9	0.9696	2.19×10^{-11}
10^{-20}	6.19	0.264	19.1	0.0263	3551	3.63×10^{-2}	0.076	53.7	9.22	0.9681	5.02×10^{-10}
10^{-19}	8.22	0.518	44.1	0.0257	1474	1.34×10^{-2}	0.087	54.6	4.78	0.9681	1.54×10^{-8}
10^{-18}	11.1	1.05	101	0.0251	582	4.69×10^{-3}	0.097	55.6	2.19	0.9678	5.29×10^{-7}
10^{-17}	15.6	2.30	219	0.0246	202	1.46×10^{-3}	0.093	56.6	0.661	0.9672	2.01×10^{-5}
10^{-16}	20.0	4.95	329	0.0243	58.9	4.03×10^{-4}	0.064	57.5	0.0889	0.9709	5.34×10^{-4}
10^{-15}	21.4	8.71	381	0.0242	18.8	1.26×10^{-4}	0.042	58.2	0.0122	0.9803	6.96×10^{-3}