



by InspectorLab

3301 N.W. 55th St., Ft. Lauderdale, FL 33309 | 888-854-0477

PREPARED FOR: GHS-BALTIMORE

TEST ADDRESS: 1419 CLARKSON STREET BALTIMORE, MD 21230



CERTIFICATE OF MOLD ANALYSIS

PREPARED FOR

GHS-BALTIMORE

PHONE NUMBER: (410) 982-6340

EMAIL: GHSBALTO@GREENHOMESOLUTIONS.COM

TEST LOCATION:

CHRIS MARTIN

1419 CLARKSON STREET

BALTIMORE, MD 21230

CHAIN OF CUSTODY # 52958674

COLLECTED: THU APRIL 10, 2025

RECEIVED: MON APRIL 14, 2025

REPORTED: MON APRIL 14, 2025

APPROVED BY:

John D. Shane PhD
Laboratory Manager

VERSION: 1.0 (A VERSION NUMBER GREATER THAN ONE (1) INDICATES THAT THE DATA IN THIS REPORT HAS BEEN AMENDED)

EPA regulations or standards for airborne or surface mold concentrations have not been established. There are also no EPA regulations or standards for evaluating health effects due to mold exposure. Information about mold can be found at www.epa.gov/mold.

All samples were received in an acceptable condition for analysis unless noted specifically in the Comments section under a particular sample. All results relate only to the samples submitted for analysis and apply to the samples as received by the laboratory. Volumes, flowrates, areas or other information are supplied by the customer. This information can affect the validity of the results. Results have not been adjusted for field or laboratory unless otherwise noted. PriorityLab bears no responsibility for sample collection activities or analytical method limitations. No warranty is either express or implied and PriorityLab assumes no responsibility or liability for errors in public information utilized, statements from sources other than PriorityLab, or developments resulting from situations outside the scope of this analysis, nor for the purpose for which the client uses the analysis. The determinations in this report are outside the scope of the AIHA LAP, LLC scope of accreditation. Contractors or consultants reviewing this report must draw their own conclusions regarding further investigation or remediation deemed necessary. PriorityLab liability is limited to the cost of the sample analysis and may not exceed the amount of the fee paid by the client.

Reports are issued free of alterations or additions and PriorityLab does not accept liability of the tampering or unlawful alteration of documents sent. All reports are expressly and exclusively for PriorityLab clients and may not be reproduced by third parties. If this report is reproduced, it must be reproduced in full unless written permission is obtained from PriorityLab. PriorityLab keeps all client data secure and confidential and any information contained in reports or files will not be divulged unless permission is expressly given by the client submitting the sample(s) except where authorized by law and all PriorityLab

FOR MORE INFORMATION, PLEASE CONTACT PRIORITYLAB AT (888) 854-0477 OR EMAIL ASK@PRIORITYLAB.COM

PREPARED FOR: GHS-BALTIMORE

TEST ADDRESS: 1419 CLARKSON STREET BALTIMORE, MD 21230

Detailed Mold Analysis (WATER-INDICATING FUNGI, IF PRESENT, ARE SHOWN BELOW IN RED)

Analysis Method	Air Analysis	Air Analysis	Air Analysis	Surface Analysis
Lab Sample #	52958674-1	52958674-2	52958674-3	52958674-4
Sample Identification	22186860	22183647	22183783	TAPE 1
Sample Location	OUTSIDE	BASEMENT	FIRST FLOOR	BASEMENT
Sample Type / Metric	Breeze ST/75L	Breeze ST/75L	Breeze ST/75L	Tape Lift
Analysis Date	Mon April 14, 2025	Mon April 14, 2025	Mon April 14, 2025	Mon April 14, 2025
Determination	CONTROL	PROBLEM	PROBLEM	GROWTH

Fungal Types Identified	Raw Count	Spores / m ³	% of Total	Raw Count	Spores / m ³	% of Total	Raw Count	Spores / m ³	% of Total		Mold Present	
*INDOOR PROBLEM FUNGI												
Chaetomium	---	---	---	28	372	54	12	160	52		---	
Chlamydo spores	---	---	---	---	---	---	---	---	---		Present	
Hyphae	---	---	---	---	---	---	---	---	---		Present	
Stachybotrys	---	---	---	5	67	9	---	---	---		---	
**Non-Problem Fungi												
Ascospores	24	319	55	3	40	5	1	13	4		---	
Basidiospores	8	106	18	6	80	11	2	27	8		---	
Cladosporium	4	53	9	2	27	3	1	13	4		---	
Penicillium/Aspergillus	5	67	11	7	93	13	4	53	17		---	
Smut/Myxomycetes	---	---	---	---	---	---	1	13	4		---	
Unclassified Pigmented Spores	2	27	4	---	---	---	2	27	8		---	
Total Spore Count*	43	570	100	51	680	100	23	310	100		NA	
Minimum Detection Limit	14			14			14			1		
Comments/Definitions Raw Count: Actual number of spores observed and counted. Spores/m³: Spores per cubic meter. % of Total: Percentage of a particular spore in relation to total number of spores. Present = growth observed. --- : Spore type was not observed. * : Indicates to look above at the names in red under "indoor problem fungi".	CONTROL samples are normally taken outside a building to provide a baseline from which samples on the interior of the building are compared. Outside air is considered normal whatever the mold counts may be. LIGHT DEBRIS: The debris present in the sample likely had no effect on the accuracy of the mold count.			Mold concentrations in the air are ABNORMAL and based on the mold counts, you likely have a mold source from which spores are able to become airborne and are an exposure concern to the occupants. LIGHT DEBRIS: The debris present in the sample likely had no effect on the accuracy of the mold count.			Mold concentrations in the air are ABNORMAL and based on the mold counts, you likely have a mold source from which spores are able to become airborne and are an exposure concern to the occupants. LIGHT DEBRIS: The debris present in the sample likely had no effect on the accuracy of the mold count.			Presence of current or former MOLD GROWTH observed. EXPOSURE TO SPORES LIKELY and will continue if the growth is not removed. An active or intermittent water source will cause the mold to continue to grow if the water source is not eliminated.		

* **Indoor Problem Fungi** are generally capable of growing on wetted building materials.

** **Non-Problem Fungi** are less capable or do not grow on wetted building materials. They are commonly found in the air outside and infiltrate into indoor air naturally.

High numbers of any one of these spore types as compared to the Control sample may indicate that they are growing on wetted building materials indoors.

Spore types not listed in this report were not observed.

Background debris estimates the amount of non-spore particles. Increasing amount of debris will affect the accuracy of the spore counts. Total percent may not equal 100% due to rounding.

#Total Spore Counts are reported to 2 significant figures.

PREPARED FOR: GHS-BALTIMORE**TEST ADDRESS:** 1419 CLARKSON STREET BALTIMORE, MD 21230

Introduction

All spores found in indoor air are also normally found in outdoor air because most originate or live in the soil and on dead or decaying plants. Therefore, it is not unusual to find mold spores in indoor air. This Mold Glossary is only intended to provide general information about the mold found in the samples that were provided to the laboratory.

Ascospores

Outdoor Habitat: Soil and decaying vegetation, dead and dying insects. These spores constitute a large part of the spores in the air and can be found in the air in very large numbers in the spring and summer, especially during and up to three (3) days after a rain.

Indoor Habitat: Very few of fungi that produce ascospores grow indoors. Some fungi that produce ascospores are recognizable by their spores and when observed are listed under their own categories. Wetted wood and gypsum wallboard paper

Allergy Potential: Depends on the type of fungus producing the ascospores.

Disease Potential: Not normally pathogenic as a group

Toxin Potential: None known

Comments: Ascospores are produced from a very large group of fungi. Notable ascospores that are considered problematic for indoor environments are Chaetomium, Peziza, and Ascotracha. If these types of ascspores are observed they will be listed in the report under their own names.

Basidiospores

Outdoor Habitat: These are mushroom spores and are common everywhere outside, especially in the late summer and fall.

Indoor Habitat: Sometimes mushrooms can be observed growing in potted plants indoors.

Allergy Potential: Rarely reported, but some Type I (hay fever, asthma) and Type III (hypersensitivity pneumonitis) has been reported.

Disease Potential: None known

Toxin Potential: None known

Comments: Mushroom spores are commonly found indoors, especially when the outdoor spore count is high. When spores of this group are derived from wood rotting fungi, including dry rot (Serpula and Poria), they can be especially destructive to buildings. When spores from destructive types of mushrooms (dry and wet rot group) are observed in the sample they are listed under their own names on the report.

PREPARED FOR: GHS-BALTIMORE

TEST ADDRESS: 1419 CLARKSON STREET BALTIMORE, MD 21230

Chaetomium

Outdoor Habitat: Commonly found on paper products, cotton products, soil, decaying vegetation, wood and natural fiber textiles (such as jute-backed carpets, canvas, etc.) and similar materials. They are rarely identified in outdoor air. These spores can be disseminated by insects, wind and water splash, etc. It is also known as a soft-rot fungus for softwood and hardwood timber.

Indoor Habitat: Chaetomium is often found on a variety of substrates containing cellulose that are chronically wetted, including paper documents, wallpaper, textiles and construction materials like gypsum board (paper-coated sheet rock) and wood.

Chaetomium can develop quickly, covering a surface with substantial growth after two weeks.

Chaetomium globosum is the most commonly found species of Chaetomium indoors. It is not that unusual to find the occasional Chaetomium spore in the air indoors.

Allergy Potential: Type I (hay fever, asthma) potential. However, no allergens have yet been characterised. However, at least two potential allergens have been isolated.

Disease Potential: Rarely reported as human pathogen.

Toxin Potential: Several known

Comments: Chaetomium spores are easily disseminated when it becomes dry. However, Chaetomium spores do not remain airborne for long unless disturbed.

This genus is often associated with termite damaged and rotting wood. These spores will continue to be found in the air until this damaged wood is removed.

High numbers of spores of this genus is not normal for indoor environments and indicate a current or former water problem. Furthermore, since the spores are held together by mucilage and trapped by hairs, few become airborne until the mold has completely dried out or is mechanically disturbed during renovations remediation. It is, therefore, not uncommon to find low Chaetomium spore counts in pre-remediation air samples and relatively higher counts in post-remediation samples.

Chaetomium species colonize surfaces under similar conditions as Stachybotrys, Alternaria, Fusarium and Ulocladium.

HIGH CONCENTRATIONS AND LONG EXPOSURES TO CHAETOMIUM SHOULD BE AVOIDED.

PREPARED FOR: GHS-BALTIMORE**TEST ADDRESS:** 1419 CLARKSON STREET BALTIMORE, MD 21230***Chlamydospores*****Outdoor Habitat:** Soil and decaying vegetation**Indoor Habitat:** Wetted wood and gypsum wallboard paper, paper products.**Allergy Potential:** None known**Disease Potential:** None known**Toxin Potential:** None known**Comments:** Asexual resting spores of all fungi***Cladosporium*****Outdoor Habitat:** Cladosporium is one of the most common environmental fungi observed worldwide and is widely reported from soil and decaying vegetation.

Cladosporium herbarum and C. cladosporioides are among the most frequently encountered species, both in outdoor and indoor environments.

Indoor Habitat: Wetted wood and gypsum wallboard paper, paper products, textiles, rubber, window sills. Cladosporium has the ability to grow at low temperatures and can thus, grow on rubber gaskets and food in refrigerators.**Allergy Potential:** Type I (hay fever, asthma) - an important and common outdoor allergen**Disease Potential:** Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals. Cladosporium are some of the most common species reported as indoor contaminants, occasionally linked to health problems.**Toxin Potential:** Cladosporium has two known toxins (cladosporin and emodin). These toxins are not known to be highly toxic. There is no evidence in the literature of toxic effects associated to inhalation of Cladosporium conidia (spores) indoors.**Comments:** The most commonly reported spore in the outdoor air worldwide. This makes Cladosporium one of the most commonly reported and abundant spore types both indoors and outdoors. The prevalence of this spore can vary throughout the year, but is especially high in late summer and autumn, especially where cereal crops are commonly planted.

An important and common allergen source.

PREPARED FOR: GHS-BALTIMORE**TEST ADDRESS:** 1419 CLARKSON STREET BALTIMORE, MD 21230***Hyphae*****Outdoor Habitat:** Any cellulose-based substance that fungi can inhabit.**Indoor Habitat:** Wetted wood and gypsum wallboard paper, etc.**Allergy Potential:** Known to be allergenic.**Disease Potential:** None known**Toxin Potential:** None known

Comments: "Root-like" structures of fungal growth that can become airborne and can possibly be allergenic.
When hyphae are found growing on a surface and associated with fruiting bodies and/or fungal spores, they indicate that growth has taken. Sometimes hyphae grow and do not produce spores. Hyphae are generally not specific to any particular type of fungus or mold type.
A mass of hyphae on a surface is indicative of mold growth.

Penicillium/Aspergillus**Outdoor Habitat:** Soil and decaying vegetation, textiles, fruits. These spores are commonly observed and are a normal part of outside air.**Indoor Habitat:** Wetted wood and gypsum wallboard paper, textiles, leather, able to grow on many types of substrates.**Allergy Potential:** Type I (hay fever, asthma), Type III (hypersensitivity pneumonitis)**Disease Potential:** Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.**Toxin Potential:** Several known

Comments: Extremely common in indoor air in low to moderate amounts as compared to the outside air. This type of spore should not be present in very high numbers as compared to the outside (control) nor constitute an overwhelming percentage (e.g., 90% or greater) of the total spores in that room(s). However, this type of mold spore is not always detected in outside air and when diversity of mold types are low in the indoor sample(s), their percentage can be 90% or more. Therefore, when the raw numbers are low the determination would be NORMAL even if the percentage is high.

There is a wide range of what is a NORMAL amount of this type of mold spores in indoor air and 200 - 700 spores per cubic meter are commonly seen in homes.

These two genera are grouped together because they cannot be reliably differentiated into their respective genera based solely on spore morphology.

PREPARED FOR: GHS-BALTIMORE**TEST ADDRESS:** 1419 CLARKSON STREET BALTIMORE, MD 21230***Smut/Myxomycetes*****Outdoor Habitat:** Soil and decaying vegetation and wood, especially dead stumps and bark**Indoor Habitat:** Not normally known to grow indoors. However the Myxomycetes can sometimes be found on firewood inside the home and especially on wood paneling. Sometimes known to grow on wood framing inside walls, ceilings and woodwork in closets.**Allergy Potential:** Type I (hay fever, asthma), rare**Disease Potential:** None known**Toxin Potential:** None known**Comments:** These two groups are difficult to distinguish due to their "round and brown" morphology. Smuts are especially common in the outside environment and can be seen in indoor air samples even during the winter in homes because the spores enter homes. These spores can be recycled through the indoor environment all year in small amounts.

A large number of these types of spores indoors can mean that there are fruiting bodies inside the home due to excessive water, usually on a wood surface(s).

Stachybotrys**Outdoor Habitat:** Soil and decaying vegetation, especially straw**Indoor Habitat:** Wetted wood, gypsum wallboard paper, cardboard boxes and ceiling tiles. This type of mold needs significant water to grow and thrive**Allergy Potential:** Type I (hay fever, asthma)**Disease Potential:** None known**Toxin Potential:** Several known (including macrocyclic trichothecenes, satratoxin F, G, H)**Comments:** Spores can be dispersed into the air when old and dry, but are wet, slimy and heavy when actively growing and thus are not easily dispersed into the air. Significantly higher numbers of spores, as compared to outside background levels, of this genus are not normal for indoor environments and indicate a current or former water problem. It is not that unusual to find the occasional *Stachybotrys* spore in the air indoors. *Stachybotrys* has several mycotoxins and has been implicated as a causative agent in disease. HIGH CONCENTRATIONS AND LONG EXPOSURES TO STACHYBOTRYS SHOULD BE AVOIDED.

PREPARED FOR: GHS-BALTIMORE**TEST ADDRESS:** 1419 CLARKSON STREET BALTIMORE, MD 21230***Unclassified Pigmented Spores*****Outdoor Habitat:** None specified**Indoor Habitat:** None specified**Allergy Potential:** Although no specific allergic potential can be given, ALL spores have the potential to be allergenic.**Disease Potential:** None known**Toxin Potential:** Unknown**Comments:** This category is for unknown spores that have at least some color and do not have enough distinctive characteristics to be identified as any particular type of spore that the laboratory recognizes.

There are a great many spore types that cannot be identified either because they are undescribed in the literature or new to science. Therefore, these types of spores are classified as "unclassified". There should not be an over abundance of this type of spore (or any spore) indoors. An large amount of this type of spore indoors would make this spore type as "water-indicating", but the origin and growth is not known.