

# Developments in Treatments Targeting Muscarinic Receptors for Alzheimer's Disease

## Abstract:

Alzheimer's Disease (AD) is the most common neurodegenerative disease to date and affects over 50 million patients worldwide, thus making it a major area of focus in drug discovery. Muscarinic Receptors, especially the M1 and M4 subtypes, are promising therapeutic targets due to their selectivity, high expression, and function in the central nervous system. Although the focus of research in AD has shifted away from muscarinic receptors in the past due to difficulties involving receptor selectivity, adverse effects, and failure in clinical trials, new discoveries in the functions of these receptors have opened up the possibility for the development of treatments with better efficacy and less risk, resulting in a resurgence of interest in them. This review aims to first examine previous research and data that have helped us recognize the functions of muscarinic receptors in AD, specifically where they are expressed in the brain, their intracellular mechanisms, and how these can help us find pathways to treat AD. Then, analyze the status of developments in different classes of treatments based on what stages in trial they were discontinued or are currently in, data regarding efficacy, pathways targeted, and adverse effects. Finally, we identify potential areas for further discovery and development of pharmaceuticals by highlighting the flaws of previous treatments, unknown aspects of AD pathology, and how more recent and less studied areas of research could guide future development and discovery.

## Introduction:

Alzheimer's Disease (AD) is a progressive neurodegenerative disorder characterized by memory loss, cognitive impairment, and behavioral changes. It is responsible for 60-70% of cases of dementia, which is a leading cause of death, disability, and dependency among the older population. (*Dementia*, n.d.) Despite its burden, current treatments are limited and not effective


enough to fully treat AD. Over the past few decades, various hypotheses for AD have developed. Today, widespread understanding of AD pathogenesis has stemmed from the cholinergic, amyloid cascade, and tau hypotheses, in addition to less recognized theories like the calcium homeostasis hypothesis. (Zhang et al., 2024)

First proposed in 1979, the cholinergic hypothesis is one of the earliest models of AD formation. This hypothesis linked a deficiency in acetylcholine (ACh) with the cognitive effects experienced by individuals with Alzheimer's. (Davies & Maloney, 1976) Acetylcholine is a neurotransmitter or chemical messenger that is synthesized in a neuron, released in the space between two neurons called a synaptic cleft, and then binds to a receptor. The hypothesis attributed the shortage in ACh to the loss of cholinergic neurons—neurons that use ACh to communicate—found in AD patients. These changes occur mostly in the hippocampus, cortex, and other cortical regions important for cognitive function. This discovery influenced the development of treatments that increased ACh levels in the brain, like acetylcholinesterase inhibitors (AChEIs). AChEIs work by inhibiting the enzyme acetylcholinesterase (AChE), which breaks down acetylcholine, therefore increasing acetylcholine levels in the brain and improving synaptic

function, specifically in the cholinergic system. Currently, Donepezil, Rivastigmine, and Galantamine are FDA-approved AChEIs in clinical use. (*Medications for Memory, Cognition & Dementia-Related Behaviors* | *Alz.Org*, n.d.) These drugs are often used in conjunction with an NMDA antagonist, Memantine. However, AChEIs plateau in their effectiveness as AD becomes more severe, calling for alternative means of increasing ACh levels. (Hogan, 2014)

Muscarinic receptors (mAChRs) and Nicotinic receptors (nAChRs)— the two types of acetylcholine receptors— have built on the cholinergic hypothesis, showing lots of potential as effective targets of therapeutics for AD. This review focuses on the developments in drugs targeting mAChRs. Muscarinic receptors are a family of class A G protein-coupled receptors (GPCRs) subdivided into 5 subtypes (M1-M5), which bind to and are activated by ACh in the central and peripheral nervous systems, a mechanism significant to the cholinergic system. Although no treatments targeting them have been approved yet, they have regained increasing interest in Alzheimer's research. Previously, research in the field shifted away from targeting muscarinic receptors for a period and instead focused on targeting biomarkers associated with AD. More specifically, on amyloid-beta (A $\beta$ ) plaques and neurofibrillary tangles (NFTs). A $\beta$  plaques are abnormal accumulations of A $\beta$  peptides, which then trigger a series of pathways, further progressing the disease. (Hardy & Higgins, 1992) One of these pathways is the hyperphosphorylation of tau proteins, leading to misfolded tau proteins that form NFTs, hindering synaptic function. (Frost et al., 2009) Understanding of these pathways has come mainly from the amyloid cascade and tau hypotheses. Treatments to come out of these discoveries include anti-amyloid therapies like Aducanumab, Lecanemab, and Donanemab. (*Medications for Memory, Cognition & Dementia-Related Behaviors* | *Alz.Org*, n.d.) However, these treatments also had limited efficacy. For instance, Lecanemab showed a 27% slower decline in Clinical Dementia Rating - Sum of Boxes scores over 18 months compared to placebo, which is statistically meaningful but modest, on top of having adverse effects like brain swelling or bleeding. (Hempel et al., 2023; Ren et al., 2025) Furthermore, by the time amyloid plaques accumulate, neurodegeneration has already advanced, limiting their ability to reverse damage. Nevertheless, the cholinergic, tau, and amyloid hypotheses have helped us gain a better idea of the mechanisms and pathways behind AD [Figure 1]. Most notably, the M1 and M4 receptors have frequently been researched for their role in cholinergic dysfunction and relationship with amyloid pathways. (Davis et al., 2010; Foster et al., 2014)

## Cholinergic Dysfunction in AD: Comparison



**Figure 1:** (Created with BioRender) In healthy conditions, acetylcholine (ACh) is efficiently released from presynaptic neurons and inhibited by M4 muscarinic receptors—particularly in forebrain regions such as the hippocampus and cortex—leading to substantial activation of M1 muscarinic receptors and regulation by acetylcholinesterase (AChE), allowing for normal cognitive function. In AD, degeneration of basal forebrain cholinergic neurons leads to reduced cholinergic neuron density, lower ACh levels, and ultimately less muscarinic activity. This dysfunction contributes to the development of Aβ plaques and NFTs, which collectively contribute to the progressive cognitive decline associated with Alzheimer’s disease. (BioRender, n.d.; Davis et al., 2010; Foster et al., 2014)

Given this current landscape of Alzheimer’s research and its treatments, this paper aims to synthesize past research to more clearly identify the potential roles muscarinic receptors can play in the progress of the field. It will first further examine the foundational studies that identified the functions, distributions, and intracellular mechanisms of mAChRs in the context of AD. Then, analyze the pathways that have been researched as a result and the status of drug developments and therapeutics that have attempted to target them. Finally, it will identify more recent and less studied discoveries that could hold solutions and discoveries in the field.

## Foundational Discoveries of mAChR Roles in AD

Many aspects of Alzheimer’s pathology are still unknown; however, their understanding has come a long way since the proposal of the cholinergic hypothesis. As more pathways and

locations of biomarkers involved in specific stages of AD have been identified, the roles mAChRs play have become clearer. While multiple factors impact their significance to AD pathology, their structures and distributions provided the fundamental rationale for their targeting in treatments.

### **Distribution and Concentration of mAChRs in Areas Impacted by Alzheimer's**

In its early stages, AD impacts areas significant to memory in the temporal lobe, specifically the hippocampus and entorhinal cortex, as well as the posterior cingulate gyrus, amygdala, and parahippocampal gyrus. Using MRIs, atrophy, brain shrinkage due to loss of neurons and synapses, has been found in these areas.(Schuff et al., 2009) As the disease progresses, it expands to other parts of the cerebral cortex, like the frontal and parietal lobes.(Ramos Bernardes da Silva Filho et al., 2017)


The M1 receptor, the most studied mAChR in AD research, is found throughout the central nervous system (CNS) in the cerebral cortex, hippocampus, and striatum, as well as the autonomic ganglia in the peripheral nervous system (PNS). The CNS consists of the brain and the spinal cord, acting as the control center. The PNS, made up of the nerves outside of the CNS, acts as the communication network that connects the CNS to the rest of the body. M1 is most densely distributed in layers II/III and V in the frontal, parietal, and temporal cortex and the hippocampus.(Levey et al., 1991) Due to its prominence in these memory circuits, it is the most commonly targeted muscarinic receptor in the CNS for AD research and treatment. While the number of M1 receptors has been found to be unaffected by AD, their signaling efficacy is impaired, most likely attributed to impaired coupling with G proteins. (Caccamo et al., 2006; Flynn et al., 1991)

Showing the highest overlap with the M1 receptor in the regions experiencing early neurodegeneration in AD, M4 receptors are mostly expressed in the caudate nucleus, putamen, and nucleus accumbens of the basal ganglia, as well as in the hippocampus and neocortex. Their high expression in the basal ganglia in particular is significant because they form part of the cortico–basal ganglia–thalamic loops, which integrate cognitive, motor, and reward-related information. In Alzheimer's, impairments in these circuits lead to cognitive impairment as well as neuropsychiatric symptoms such as apathy, agitation, and impaired decision-making. Similarly, M2 is an autoreceptor expressed in the hippocampus, cerebral cortex, and presynaptic terminals throughout the CNS, as well as in autonomic nuclei of the brainstem.(Levey et al., 1991) While not yet proved to be as effective a target due to its primary role being in the PNS, it provides major cholinergic pathways to the cortex and hippocampus.(Smiley et al., 1999) Acting as inhibitory autoreceptors regulating ACh release, its loss in Alzheimer's patients contributes significantly to the cholinergic deficits associated with AD.

Finally, while the M3 and M5 receptors are mostly distributed in the PNS, they have displayed some therapeutic potential as targets. In neurons, M3 enables excitatory cholinergic signaling and plays a role in neurovascular coupling, mediating vasodilation–the widening of blood vessels– through Gq-protein pathways in endothelial cells of the cerebral vascular

system.(Kruse et al., 2012) The M5 receptor, while the least abundant muscarinic subtype in the CNS, also regulates cerebral blood flow. Hence, the M3 and M5 receptors' ability to manage cerebral blood flow could make them valuable targets in early disease stages.

## Distribution of mAChR Subtypes in the Human Brain




**Figure 2. Distribution of muscarinic acetylcholine receptor (mAChR) subtypes in the human brain:** Coronal, axial, sagittal, and 3D projections from the Allen Human Brain Atlas showing gene expression patterns of mAChR subtypes in the areas they are most heavily distributed in (The brighter the red, the more concentrated).

- A. M1 expression in the cerebral cortex
- B. M2 expression in the pons
- C. M3 expression in the hypothalamus
- D. M4 expression in the striatum
- E. M5 expression in the pons

### **Structure of mAChRs**

Since mAChRs are GPCRs, all 5 subtypes have seven transmembrane (7TM) helices connected by extracellular and intracellular loops with an extracellular N-terminus and intracellular C-terminus. The orthosteric, or primary, binding site for ACh is located deep inside the 7TM bundle, which allows for precise ligand binding, conformational change—change in receptor shape after binding—to initiate intracellular signaling through heterotrimeric G proteins. The G protein that each mAChR couples with influences the pathways they signal. The M1, M3, and M5 receptors couple to Gq/11, which activates the phospholipase C (PLC)—IP<sub>3</sub>—Ca<sup>2+</sup> signaling cascade that promotes neuronal excitability and synaptic plasticity. On the other hand, the M2 and M4 receptors couple to Gi/o, which inhibit adenylyl cyclase and modulate ion channel activity to regulate neurotransmitter release. GPCR structure makes mAChRs a more ideal target for therapeutics than the other family of acetylcholine receptors, nAChRs. Unlike nAChRs, which are ligand-gated ion channels with rapid but less modulatable responses, mAChRs transduce signals through G protein-mediated cascades. This allows for a broader range of pharmacological modulation, like biased agonism, allosteric regulation, and subtype selectivity. (Wess, 1996) Furthermore, research using high-resolution crystallography and cryo-electron microscopy has discovered pocket structures and loop conformations that are

unique to each mAChR subtype.



**Figure 3:** (Created with BioRender) M1, M3, and M5 receptors couple to Gq/11 proteins, activating phospholipase C (PLC) and leading to production of IP<sub>3</sub> and second messenger Diacylglycerol (DAG), which increase intracellular Ca<sup>2+</sup> and activate protein kinase C (PKC). M2 and M4 receptors couple to Gi/o proteins, which inhibit adenylate cyclase, decrease cAMP levels, and activate G protein-coupled receptor kinase (GIRK) channels, resulting in membrane hyperpolarization.

## Connections to Other Pathways and Diseases

In addition to being central to cholinergic neurotransmission, mAChRs also intersect with multiple pathological mechanisms relevant to AD and other neurological disorders. Understanding these cross-pathway interactions is critical, as it helps create the bigger picture, allowing for the development of therapies with greater efficacy and less adverse effects.

### Amyloid and Tau Pathologies

As mentioned before, muscarinic receptors influence the development of amyloid and tau pathologies [Figure 1]. Early research *in vitro* demonstrated that muscarinic stimulation increases sAPP $\alpha$  release, which inhibits and suppresses A $\beta$  generation, an effect blocked by non-selective muscarinic antagonism and PKC inhibition. (Lin et al., 1999) The M1 receptor, however, has a direct effect as it modulates enzymes involved with the amyloid precursor protein (APP) processing activation. More specifically, it promotes non-amyloidogenic processing of amyloid precursor protein (APP) through stimulation of alpha-secretase, reducing the production of pathogenic A $\beta$  peptides. (Cisse et al., 2011; Fisher, 2008) *In vivo*, the 3xTg-AD mouse model, chronic treatment with the M1 agonist AF267B increased cortical and hippocampal sAPP $\alpha$  through ADAM17 activation, reduced A $\beta$ 42 levels and plaque burden, and

improved memory.(Caccamo et al., 2006) In addition to rodent models, AF267B also lowered CSF A $\beta$  in rabbits.(Beach et al., 2001)

In parallel, M1 receptor stimulation also helps decrease the tau pathology associated with AD. Through PLC/Ca<sup>2+</sup>-dependent protein kinase C signaling and downstream inhibition of GSK3 $\beta$ , M1 activation reduces tau phosphorylation in vitro and in vivo. In vitro, muscarinic activation modulated tau kinase cascades that included PKC and mitogen-activated protein kinase(MAPK), impacting sites relevant to tangle formation.(Forlenza et al., 2000) Additionally, in the same study, which found that M1 activation by AF267B in 3xTg-AD mice decreased amyloid pathology, AF267B treatment also decreased hyperphosphorylated tau. (Caccamo et al., 2006)

### **Links to Dopaminergic and Glutamatergic Systems**

In addition to cholinergic dysfunction, imbalances in the dopaminergic and glutamatergic systems significantly contribute to AD pathogenesis and progression. The dopaminergic system is primarily involved with brain functions like motivation, reward, and motor control, all of which are impacted by Alzheimer's. Muscarinic receptors, especially the M4 subtype, play essential roles in modulating dopaminergic transmissions. More specifically, activation of M4 limits dopamine release, making it a useful target for countering the dopamine imbalances associated with AD. Nicotinic receptors, which, unlike mAChRs, promote dopamine release through a quicker and more direct pathway, are also significant to dopamine balance. However, targeting nAChRs has been more challenging due to obstacles like downregulation in AD, receptor selectivity, receptor desensitization, and subtype selectivity. More specifically, in AD, nicotinic receptor expression of subtypes  $\alpha$ 4 $\beta$ 2 and  $\alpha$ 7 significantly diminished, reducing their efficacy as therapeutic targets.(Hoskin et al., 2018; Pimlott et al., 2004) Furthermore, issues like receptor desensitization with chronic activation and lack of subtype selectivity can lead to off-target effects.(Choi, 1988)

On the other hand, dysfunction in the glutaminergic system– the brain's primary excitatory network with glutamate as a neurotransmitter– is directly linked to the excitotoxicity and neuronal death found in AD. Among glutamate receptors, N-methyl-D-aspartate (NMDA) receptors are especially important because they are highly permeable to calcium and directly mediate synaptic plasticity, long-term potentiation (LTP), and memory encoding.(Malenka & Bear, 2004) In Alzheimer's, NMDARs are overactivated and disrupted by A $\beta$  plaques, which is why antagonists like memantine are used clinically to reduce excitotoxicity. However, memantine only dampens overactivation and does not restore the physiological pro-plasticity role of NMDA receptors. M1 receptors modulate NMDAR currents heavily. In particular, M1 activation has been shown to enhance NMDA receptor-mediated excitatory postsynaptic currents in hippocampal pyramidal neurons.(Marino et al., 1998) Additionally, M1 increases neuronal excitability and calcium entry through NMDA receptors by inhibiting small-conductance calcium-activated potassium (SK) channels.(Conn et al., 2009) Together, these mechanisms potentiate NMDAR currents and facilitate LTP, making M1 a very useful target in countering this aspect of AD progression.

## Connection to Other Diseases

In addition to AD, mAChRs have been proven to interact with many of the pathways in other neuropsychiatric and neurodegenerative disorders, such as Schizophrenia (SZ), Parkinson's, and Huntington's diseases. This has led to the testing of the same drugs for both SZ and AD, most notably xanomeline. Parkinson's, which is mainly characterized by the loss of dopaminergic neurons, has been associated with the loss of M4 receptors in the striatum due to their role in modulation.(Nielsen & Ford, 2024) Similarly, a decrease in the density and total of M2 has been found in post-mortem Huntington patients.(Lange et al., 1992) Although SZ is mostly classified as a neuropsychiatric disorder, multiple studies have demonstrated that M1 and M4 interact with many of the same neural circuits as AD. This has led to the testing of the same drugs for both SZ and AD, most notably xanomeline.

## The Progress and Current Status of mAChR-Targeted Therapeutics

Despite the advantageous aspects of muscarinic acetylcholine receptors identified previously, mAChRs, early attempts to develop effective therapeutics faced major obstacles. These limitations revolved mainly around the lack of available compounds that were selective enough to only target one mAChR subtype, peripheral cholinergic side effects, and, while broader than nicotinic modulation, the still narrow window for modulation of mAChRs. Fortunately, recent research has shifted to improving subtype selectivity, blood–brain barrier (BBB) penetration, and utilizing novel mechanisms that modulate receptors to achieve cognitive benefits that couldn't have been achieved before. Due to their role in AD pathology, M1 and M4 are usually the targets of these compounds.

### Subtype Selective Orthosteric Ligands

Orthosteric ligands target the acetylcholine binding site, deep inside the 7TM bundle. This site is common among all AChRs, making subtype selectivity challenging. For instance, Cevimeline (AF102B), developed as an orthosteric M1 agonist for AD, could cross the BBB and showed improved memory and reduced A $\beta$  pathology in rodents.(Mitoh et al., 2017) In human trials, the drug exhibited the same benefits but also showed peripheral adverse effects, causing it to fail FDA approval.(Oleksak et al., 2021) An older agent, which revealed that nonselective orthosteric ligands caused peripheral adverse effects, was the M1 and M4 agonist, xanomeline. Although now recognized as a dual orthosteric and allosteric M1 agonist, xanomeline exhibited similar adverse effects associated with its orthosteric off-target activations of M2 and M3 subtypes, such as nausea, vomiting, excessive salivation, gastrointestinal disturbances, and bradycardia.(Burger et al., 2023; Noetzel et al., 2009)

An emerging agent known as Kar-XT has combined xanomeline with tropium chloride, a peripherally restricted muscarinic antagonist that does not cross the BBB. This design preserves xanomeline's central M1/M4 activity while significantly reducing activity in the PNS. In a phase 1

healthy volunteer study (NCT02831231), the presence of trospium chloride reduced composite cholinergic adverse events by 46% compared to xanomeline alone. Adverse effects such as nausea, vomiting, diarrhea, sweating, and salivary hypersecretion were also reduced by at least 29%. (Breier et al., 2023) Additionally, in prior phase 2 schizophrenia trials, KAR-XT demonstrated good efficacy and tolerability. (Correll et al., 2022) While these clinical trials were conducted for schizophrenia, they target the same mechanism—muscarinic activation in the CNS protected from peripheral side effects—holding relevance for Alzheimer’s disease, where similar challenges of peripheral toxicity have long limited the development of orthosteric mAChR agonists.

Beyond Kar-XT, recent research using high-resolution crystallography and cryo-electron microscopy has discovered pocket geometries and loop conformations that are unique to each mAChR subtype, allowing for the design of ligands with improved bias. (Thal et al., 2016) For instance, VU0357017 and AC-260584 are M1-preferring agonists that have shown high efficacy while having little to no activation of the M2-M5 subtypes. (S. R. Bradley et al., 2010; Digby et al., 2012) AC-260584 exhibited potent functional activity ( $pEC_{50}$  of  $7.7 \pm 0.1$ ) and high efficacy with reduced activity at M2-M5. VU0357017 activated calcium signaling and ERK1/2 phosphorylation—pathways downstream of G-protein-coupled activation—while showing minimal  $\beta$ -arrestin recruitment, indicating a bias toward cognitive-relevant signaling cascades. While orthosteric agents offer a direct mechanism of action, even highly selective compounds can have adverse effects due to powerful signaling even at low levels of off-target activation. This risk has driven interest in allosteric modulators and bitopic ligands, which offer refined, tissue-specific modulation with potentially wider therapeutic scopes.

## **Allosteric Modulators**

Allosteric modulators bind to sites on the muscarinic receptor other than the orthosteric site. Unlike orthosteric ligands, which compete directly with endogenous ACh, allosteric modulators change receptor activity by influencing the receptor’s reaction to naturally released ACh. Positive Allosteric Modulators (PAMs) do this by increasing the response to acetylcholine, while Negative Allosteric Modulators (NAMs) regulate the response. Subtype selectivity is more achievable since allosteric sites are less conserved and more structurally unique among mAChR subtypes. Furthermore, endogenous ACh is released in specific locations and at precise times to regulate physiological functions. Allosteric modulation only occurs where endogenous ACh is present, so the risk of unwanted targeting in the PNS is reduced.

The most studied allosteric modulator has been M1-selective PAMs. The prototype M1 PAM, benzyl quinolone carboxylic acid (BQCA), has been developed as a preclinical research tool. It substantially promoted downstream signaling in medial prefrontal cortex (mPFC) pyramidal neurons, with effects absent in M1 knockout mice. (Shirey et al., 2009) Most notably, BQCA restored discrimination reversal learning, often lost in AD patients, in a transgenic AD mouse model and regulated non-amyloidogenic APP processing in vitro. This suggested that they could

possess symptomatic and disease-modifying potential. Additionally, in prion-diseased mice—an in vivo model with cholinergic dysfunction similar to AD—daily BQCA doses restored cognitive function and significantly increased survival, with minimal side effects compared to treatment with orthosteric agonists or acetylcholinesterase inhibitors. (S. J. Bradley et al., 2017) Despite this, BQCA displayed poor pharmacokinetic properties. More specifically, although it reached peak brain concentrations around 1.5 hours post-administration, systemic and tissue distribution favored organs like the kidney and lung over the brain, limiting its translational potential. (Dwomoh et al., 2022) Building on BQCA's model, MK-7622 became the first human-tested M1 PAM for AD. The phase II trial evaluated it as an adjunct to standard AChEI therapy but was terminated early due to it having no significant cognitive or functional benefit compared to placebo. Importantly, cholinergic-like adverse events such as diarrhea and nausea were more frequent than in control groups. (Voss et al., 2018) Displaying similar reactions on rodents, PF-06767832 and PF-06764427 helped identify intrinsic agonism or activation of the receptor as a cause for these effects. (Davoren et al., 2016; Moran et al., 2018) As a result, M1-PAMs like VU319, which minimize agonist activity, have shown a lack of adverse effects and are being further researched. (Poslunsey et al., 2024)

Given that M4 receptors act as inhibitory autoreceptors on cholinergic interneurons and as modulators of dopaminergic signaling, recent research has also been studying how they could potentially be targeted using PAMs to establish an excitatory–inhibitory balance in the networks affected by AD. Due to their recent rise to attention, most data on M4 PAMs for Alzheimer's treatment is preclinical and builds off of research on Schizophrenia, which involves some common pathways. A compound that has shown promise in this class of treatments is VU0152100, which is being developed at Vanderbilt University. It reversed amphetamine-induced hyperlocomotion in rodent models but not M4 knockout(KO) mice—demonstrating high selectivity—, lacked intrinsic agonist activity, and did not produce the adverse effects associated with non-selective agonists. (Byun et al., 2014) Although initially researched for schizophrenia, the mechanism of reducing excessive excitability through M4 signaling is relevant to Alzheimer's pathology, where similar dysregulation occurs. Among the next generation of M4 PAMs are VU0467154, CVL-231, and NMRA-86,1, which have shown better potency as well as many of the same effects seen in VU015210,0, showing progress and potential in this type of treatment. (Berezovskaia et al., 2025; Butler et al., 2024) Additionally, in 2018, it was revealed that Xanomeline could bind to the allosteric site on M4, allowing for further understanding of the mechanisms at work behind Kar-XT, and the potential for Xanomeline to be used as a PAM. (Burger et al., 2023)

Although less popular, negative allosteric modulators have also been explored for their roles in AD. Their purpose, unlike positive allosteric modulators, is to reduce the potency of endogenous ACh. Therefore, they target areas where mAChRs overactivation could be problematic, like the M2 receptor, which limits ACh in the cortex and hippocampus. (Zuchner et al., 2005) It has been suggested that selective M2 antagonism can increase synaptic ACh and potentially improve cognition in AD by complementing AChEIs. The M2-selective NAM, Gallamine— a neuromuscular blocker— demonstrated negative cooperativity at muscarinic receptors, guiding research into M2 NAMs. (Mash et al., 1985) Since then, there has been some research in

developing more subtype-selective and CNS-suitable M2-selective NAM candidates for AD. Despite some early compounds having demonstrated potential, factors like the possibility for increased A $\beta$  formation, less distribution in the CNS, and the lack of a complete understanding of the M2 receptor and its mechanisms, focus has remained on the more promising M1 and M4 receptors. While allosteric modulators solve many of the issues regarding subtype selectivity that orthosteric ligands couldn't, intrinsic agonism and their efficacy in general have driven focus to another type of agent, bitopic ligands.


## **Bitopic Ligands**

Bitopic or dualsteric ligands are a more recent method of targeting GPCRs that often have greater affinity and selectivity than orthosteric ligands or allosteric modulators alone. Engineered to engage the orthosteric site and a more structurally unique allosteric site at the same time, they can activate only designated downstream pathways through a process known as biased signaling. This is useful for targeting mAChRs in the CNS, where peripheral side effects limit traditional agonists. Therefore, bitopic ligands can potentially deliver CNS-selective effects with fewer off-target actions.


Like PAMs, bitopic ligands are a promising drug class for targeting M1. One of the earliest M1 targeting bitopic ligands, TBPB(tert-butyl peroxybenzoate) was initially thought to achieve its selectivity only through allosteric binding. However, later studies showed that mutations in both the orthosteric and allosteric binding sites affected the binding and signaling of TBPB.(Keov et al., 2014) In rodent models, TBPB enhanced cortical acetylcholine release, improved performance in object recognition tasks, and promoted non-amyloidogenic APP processing through increased amyloid-secretase activity while still not inducing gastrointestinal side effects.(Jones et al., 2008) Due to pharmacokinetic limitations, including poor solubility and modest brain penetration, development of TBPB ended; nevertheless, it serves as proof that M1-selective bitopic ligands could modulate pathways significant to Alzheimer's. 77-LH-28-1, another M1-directed bitopic ligand, improved spatial and working memory, in part by modulating calcium homeostasis and synaptic plasticity in hippocampal neurons without peripheral adverse effects.(Langmead et al., 2008) Bitopic ligands targeting M4 for AD are less developed than those for M1, and none have yet emerged as especially promising.

Binding Structures of Different Agent Types


a)



b)



c)



**Figure 4:** Chymol structure visualizations of class A GPCRs bound to orthosteric, allosteric, and bitopic ligands found on the RCSB Protein Data Bank. A2A and beta2 adrenergic receptors were used due to limited access to PDBs of mAChR binding.

4A: The beta2 adrenergic receptor bound to a bitopic ligand

4B: The A2A adrenergic receptor bound to an allosteric ligand

4C: The A2A adrenergic receptor bound to an orthosteric ligand

## Limitations of Agents In Current Trials and Future Directions

In addition to struggles relating to tolerability, pharmacokinetics, and target selectivity, recent trials of muscarinic-based therapies [Table 1] for AD are encountering a new set of challenges. A persistent translational gap remains, with many compounds showing efficacy in preclinical animal models but limited or inconsistent benefits in humans. This is often due to issues with species differences in receptor mechanisms and disease progression. Furthermore, the lack of known biomarkers for target engagement makes it difficult to determine whether disappointing outcomes reflect insufficient receptor modulation or true therapeutic inefficacy. Safety concerns also continue to emerge. For example, the selective M1 agonist HTL0018318 was associated with transient increases in blood pressure, posing a risk for elderly patients with cardiovascular comorbidities. (Nathan et al., 2022) Broader concerns about recruitment strategies in trials failing to take into account all groups of individuals, as well as unethical practices, remain. Recently, a prominent paper on amyloid pathology in AD published in 2006 by a team led by Sylvain Lesné was found to have manipulated images and data, misleading the direction of Alzheimer's research for over 15 years.

Given the aforementioned obstacles, muscarinic research for AD treatments is moving toward precision medicine approaches that find therapeutic strategies depending on patient-specific biology. The development of Positron Emission Tomography (PET) tracers for muscarinic receptors, such as [11C]VC-002 for M1, provides a tool to measure receptor availability in vivo. (Haense et al., 2012) This technology could allow for the grouping of patients based on receptor density and track target engagement during clinical trials. As a result, researchers can confirm with more confidence the efficacy of a drug. Genetic insights are also guiding muscarinic research. For instance, individuals carrying the APOE4 gene showed significant vulnerability to cholinergic dysfunction. (Hill et al., 2007) Hence, they may particularly benefit from muscarinic-targeted therapies early on. Additionally, mutations and polymorphisms in the CHRM1 and CHRM4 genes have been linked to altered receptor expression or signaling, influencing the individual responses of drugs being tested. (Dean & Scarr, 2021) The integration of genomic profiling into trials could reduce discrepancies like these. Combination therapy could also be helpful, given the complex nature of AD. More specifically, M4 PAMs could enhance cognition while anti-amyloid drugs like Aducanumab, Lecanemab, and Donanemab slow progression. Similarly, given that Memantine is taken in conjunction with AChEIs, pairing them with mAChR targeting therapeutics could improve balance in the glutamatergic system.

In the future, large molecule treatments may complement or even surpass small molecules in muscarinic receptor drug discovery. Allosteric antibodies, while not yet applied clinically to

muscarinic receptors, have already been developed for other GPCRs such as CCR5 and GLP-1R, for which they have stabilized receptor conformations. (Cong et al., 2021; Olson & Jacobson, 2009) Applying this strategy to M1 or M4 could lead to the development of therapeutics with improved selectivity, pharmacokinetics and tissue specificity, potentially overcoming the adverse effects seen with small-molecule drugs. Nanobodies (VHHs) are another promising platform as they have already been used to stabilize muscarinic receptor conformations. For instance, nanobodies Nb9-8 and Nb10-6 locked the M2 receptor in active states. (Kruse et al., 2013) With further engineering, nanobodies could be developed as therapeutic allosteric modulators of M1 or M4. Finally, chemogenetic approaches offer proof-of-concept for circuit-specific cholinergic modulation. Designer Receptors Exclusively Activated by Designer Drugs (DREADDs), which were originally engineered from muscarinic receptors, have demonstrated this. The hM3Dq receptor, derived from M3, produces excitatory Gq signaling, while hM4Di, engineered from M4, produces inhibitory Gi signaling. (Smith et al., 2016) Although DREADDs are primarily research tools, they have shown how muscarinic receptor scaffolds can be modified to enable precise spatiotemporal control of signaling, showing the potential for future gene-therapy-based interventions in Alzheimer’s disease.

**Table 1: Prominent Muscarinic Agents Currently Under Research for Relevance to AD**

Agent	Class / Mechanism	Development Stage & Context	Relevance to AD
KarXT (xanomeline + trospium chloride)	Dual Orthosteric and Allosteric M1/M4 agonist + peripheral orthosteric antagonist	Phase 3 trials for psychosis and agitation in AD; FDA-approved for schizophrenia as of 2024  Invented at PureTech Health; advanced by Karuna Therapeutics; acquired by Bristol Myers Squibb (BMS)	Central activation of M1/M4 while minimizing peripheral side effects; cognitive and behavioral relevance in AD.
HTL0018318	Orthosteric M1-selective agonist	Phase 1b/2a adjunct to donepezil in mild–moderate AD  Developed by Sosei Heptares in collaboration with Allergan	Designed for enhanced cognitive benefit with improved receptor specificity.
HTL0016878	Orthosteric M4-selective agonist	Phase 1 in healthy volunteers	Targeting neurobehavioral

		Originated at Heptares Therapeutics; Advanced by Neurocrine Biosciences	symptoms via M4 pathways.
HTL9936	Orthosteric M1-selective agonist	Phase 1 in healthy (elderly inclusive) subjects  Developed by Heptares Therapeutics	High selectivity addressing cognitive impairment with safety focus.
VU319	M1-positive allosteric modulator (PAM) with minimal intrinsic agonism	Phase 1 completed; proof-of-concept planned  Developed within Vanderbilt University's Warren Center for Neuroscience Drug Discovery (WCNDD)	Reduced side effects compared to earlier PAMs like MK-7622.
CVL-231 (Emraclidine)	M4 PAM	Phase 1 multiple-dose; includes AD dementia cohort  Developed by Cerevel Therapeutics; now under AbbVie	Balances excitatory/inhibitory signaling relevant in AD.
NS-136	M4 PAM	Phase 1 for schizophrenia; preclinical AD development  Developed by NeuShen Therapeutics	Forward-looking candidate, bridging schizophrenia and AD.
Compound 24	M4 PAM (tool compound)	Preclinical in rodents and rhesus monkeys  N/A (research tool)	Demonstrated procognitive and antipsychotic-like effects in primate models.

VU0467154	M4 PAM	Preclinical (rodent) Originated at Vanderbilt University	Enhanced learning and memory, strengthening case for cognitive benefit.
VU0152100 / VU0152099 / LY2033298	M4-selective PAMs (tool compounds)	Preclinical (rodents) N/A	Validated M4 as an attractive target for cognition and behavior modulation.
AF267B (also AF102B / AF150(S))	Orthosteric M1-selective agonist	Preclinical (various animal models) N/A	Reduced A $\beta$ and tau pathology; rescued memory in transgenic AD mice.
AF710B	M1-selective agonist with $\sigma$ 1 activity (bitopic-like)	Preclinical (rodents) N/A	Improved cognition and APP processing with high selectivity.
GSK1034702	Bitopic M1 agonist	Clinical testing in humans but problematic due to side effects (not discontinued yet)  Developed by GlaxoSmithKline (GSK)	Demonstrated bitopic engagement and cognitive effects, but poor safety highlights need for refined ligands.

## Conclusion

Research in Alzheimer's pathogenesis, specifically with regards to the mAChRs, has certainly undergone significant progress since the proposition of the cholinergic hypothesis in 1976. Moving beyond the initial setbacks of earlier orthosteric agonists, more refined approaches with subtype selectivity, allosteric modulation, and biased signaling have been discovered. The M1 and M4 subtypes remain the most prominent targets due to their dense expression in memory-related and dopaminergic circuits and their direct influence on amyloid and tau pathology. Understanding of the overlap between these systems and other neurological diseases has contributed to the development of treatments like Kar-XT which were successful in trials for these related disorders, like SZ, and have since shown promise in AD trials.

Nevertheless, much remains unknown and preclinical. Translation from animal models to humans continues to be inconsistent, trial design is slowed by the lack of biomarkers that are distinguishable, and the heterogeneity of Alzheimer's demands more precision clinical trials. Muscarinic-targeted therapies with genetic profiling, PET tracers, and potentially combinatorial strategies with anti-amyloid or anti-tau drugs could help overcome these challenges. Furthermore, beyond small molecules, new techniques such as allosteric antibodies,

nanobodies, and receptor-based gene therapies are beginning to gain attention. Together, these developments have improved understanding as well as served as a guide for the potential for innovation in treatments targeting mAChRs.

## **Acknowledgements**

I would like to thank [REDACTED] for their guidance and feedback throughout the research process.

## **Author**

Arnav Chatterjee is a senior at Saint John's High School in Shrewsbury, Massachusetts. Having done research on whether ATP synthase could lead to increased energy production, epilepsy, and developments in muscarinic receptor targeting treatments for Alzheimer's, Arnav has a deep interest and passion for the life sciences.

## References

- Beach, T. G., Walker, D. G., Potter, P. E., Sue, L. I., & Fisher, A. (2001). Reduction of cerebrospinal fluid amyloid  $\beta$  after systemic administration of M1 muscarinic agonists. *Brain Research*, *905*(1), 220–223. [https://doi.org/10.1016/S0006-8993\(01\)02484-2](https://doi.org/10.1016/S0006-8993(01)02484-2)
- Berezovskaia, A., Lindsley, C., Fink-Jensen, A., & Wörtwein, G. (2025). M4 Positive Allosteric Modulator VU0467154 Impacts Amphetamine Sensitization and Spontaneous Locomotion in Male Mice. *ACS Chemical Neuroscience*, *16*(5), 868–879. <https://doi.org/10.1021/acscemneuro.4c00795>
- BioRender*. (n.d.). Retrieved August 5, 2025, from <https://app.biorender.com/illustrations/6891dfba4e0509a1dbb919db>
- Bradley, S. J., Bourgoignon, J.-M., Sanger, H. E., Verity, N., Mogg, A. J., White, D. J., Butcher, A. J., Moreno, J. A., Molloy, C., Macedo-Hatch, T., Edwards, J. M., Wess, J., Pawlak, R., Read, D. J., Sexton, P. M., Broad, L. M., Steinert, J. R., Mallucci, G. R., Christopoulos, A., ... Tobin, A. B. (2017). M1 muscarinic allosteric modulators slow prion neurodegeneration and restore memory loss. *The Journal of Clinical Investigation*, *127*(2), 487–499. <https://doi.org/10.1172/JCI87526>
- Bradley, S. R., Lamah, J., Ohrmund, L., Son, T., Bajpai, A., Nguyen, D., Friberg, M., Burstein, E. S., Spalding, T. A., Ott, T. R., Schiffer, H. H., Tabatabaei, A., McFarland, K., Davis, R. E., & Bonhaus, D. W. (2010). AC-260584, an orally bioavailable M(1) muscarinic receptor allosteric agonist, improves cognitive performance in an animal model. *Neuropharmacology*, *58*(2), 365–373. <https://doi.org/10.1016/j.neuropharm.2009.10.003>
- Breier, A., Brannan, S. K., Paul, S. M., & Miller, A. C. (2023). Evidence of trospium's ability to mitigate cholinergic adverse events related to xanomeline: Phase 1 study results. *Psychopharmacology*, *240*(5), 1191–1198. <https://doi.org/10.1007/s00213-023-06362-2>
- Burger, W. A. C., Pham, V., Vuckovic, Z., Powers, A. S., Mobbs, J. I., Laloudakis, Y., Glukhova, A., Wootten, D., Tobin, A. B., Sexton, P. M., Paul, S. M., Felder, C. C., Danev, R., Dror,

- R. O., Christopoulos, A., Valant, C., & Thal, D. M. (2023). Xanomeline displays concomitant orthosteric and allosteric binding modes at the M4 mAChR. *Nature Communications*, 14(1), 5440. <https://doi.org/10.1038/s41467-023-41199-5>
- Butler, C. R., Popiolek, M., McAllister, L. A., LaChapelle, E. A., Kramer, M., Beck, E. M., Mente, S., Brodney, M. A., Brown, M., Gilbert, A., Helal, C., Ogilvie, K., Starr, J., Uccello, D., Grimwood, S., Edgerton, J., Garst-Orozco, J., Kozak, R., Lotarski, S., ... Steyn, S. J. (2024). Design and Synthesis of Clinical Candidate PF-06852231 (CVL-231): A Brain Penetrant, Selective, Positive Allosteric Modulator of the M4 Muscarinic Acetylcholine Receptor. *Journal of Medicinal Chemistry*, 67(13), 10831–10847. <https://doi.org/10.1021/acs.jmedchem.4c00293>
- Byun, N. E., Grannan, M., Bubser, M., Barry, R. L., Thompson, A., Rosanelli, J., Gowrishankar, R., Kelm, N. D., Damon, S., Bridges, T. M., Melancon, B. J., Tarr, J. C., Brogan, J. T., Avison, M. J., Deutch, A. Y., Wess, J., Wood, M. R., Lindsley, C. W., Gore, J. C., ... Jones, C. K. (2014). Antipsychotic drug-like effects of the selective M4 muscarinic acetylcholine receptor positive allosteric modulator VU0152100. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, 39(7), 1578–1593. <https://doi.org/10.1038/npp.2014.2>
- Caccamo, A., Oddo, S., Billings, L. M., Green, K. N., Martinez-Coria, H., Fisher, A., & LaFerla, F. M. (2006). M1 Receptors Play a Central Role in Modulating AD-like Pathology in Transgenic Mice. *Neuron*, 49(5), 671–682. <https://doi.org/10.1016/j.neuron.2006.01.020>
- Choi, D. W. (1988). Glutamate neurotoxicity and diseases of the nervous system. *Neuron*, 1(8), 623–634. [https://doi.org/10.1016/0896-6273\(88\)90162-6](https://doi.org/10.1016/0896-6273(88)90162-6)
- Cisse, M., Braun, U., Leitges, M., Fisher, A., Pages, G., Checler, F., & Vincent, B. (2011). ERK1-independent  $\alpha$ -secretase cut of  $\beta$ -amyloid precursor protein via M1 muscarinic receptors and PKC $\alpha/\epsilon$ . *Molecular and Cellular Neuroscience*, 47(3), 223–232. <https://doi.org/10.1016/j.mcn.2011.04.008>

- Cong, Z., Chen, L.-N., Ma, H., Zhou, Q., Zou, X., Ye, C., Dai, A., Liu, Q., Huang, W., Sun, X., Wang, X., Xu, P., Zhao, L., Xia, T., Zhong, W., Yang, D., Eric Xu, H., Zhang, Y., & Wang, M.-W. (2021). Molecular insights into ago-allosteric modulation of the human glucagon-like peptide-1 receptor. *Nature Communications*, *12*, 3763.  
<https://doi.org/10.1038/s41467-021-24058-z>
- Conn, P. J., Jones, C. K., & Lindsley, C. W. (2009). Subtype-selective allosteric modulators of muscarinic receptors for the treatment of CNS disorders. *Trends in Pharmacological Sciences*, *30*(3), 148–155. <https://doi.org/10.1016/j.tips.2008.12.002>
- Correll, C. U., Angelov, A. S., Miller, A. C., Weiden, P. J., & Brannan, S. K. (2022). Safety and tolerability of KarXT (xanomeline–trospium) in a phase 2, randomized, double-blind, placebo-controlled study in patients with schizophrenia. *Schizophrenia*, *8*(1), 109.  
<https://doi.org/10.1038/s41537-022-00320-1>
- Davies, P., & Maloney, A. J. F. (1976). SELECTIVE LOSS OF CENTRAL CHOLINERGIC NEURONS IN ALZHEIMER'S DISEASE. *The Lancet*, *308*(8000), 1403.  
[https://doi.org/10.1016/S0140-6736\(76\)91936-X](https://doi.org/10.1016/S0140-6736(76)91936-X)
- Davis, A. A., Fritz, J. J., Wess, J., Lah, J. J., & Levey, A. I. (2010). Deletion of M1 Muscarinic Acetylcholine Receptors Increases Amyloid Pathology In Vitro and In Vivo. *Journal of Neuroscience*, *30*(12), 4190–4196. <https://doi.org/10.1523/JNEUROSCI.6393-09.2010>
- Davoren, J. E., Lee, C.-W., Garnsey, M., Brodney, M. A., Cordes, J., Dlugolenski, K., Edgerton, J. R., Harris, A. R., Helal, C. J., Jenkinson, S., Kauffman, G. W., Kenakin, T. P., Lazzaro, J. T., Lotarski, S. M., Mao, Y., Nason, D. M., Northcott, C., Nottebaum, L., O'Neil, S. V., ... Grimwood, S. (2016). Discovery of the Potent and Selective M1 PAM-Agonist N-[(3R,4S)-3-Hydroxytetrahydro-2H-pyran-4-yl]-5-methyl-4-[4-(1,3-thiazol-4-yl)benzyl]pyridine-2-carboxamide (PF-06767832): Evaluation of Efficacy and Cholinergic Side Effects. *Journal of Medicinal Chemistry*, *59*(13), 6313–6328.  
<https://doi.org/10.1021/acs.jmedchem.6b00544>

- Dean, B., & Scarr, E. (2021). Changes in cortical gene expression in the muscarinic M1 receptor knockout mouse: Potential relevance to schizophrenia, Alzheimer's disease and cognition. *Npj Schizophrenia*, 7(1), 44. <https://doi.org/10.1038/s41537-021-00174-z>
- Dementia*. (n.d.). Retrieved July 29, 2025, from <https://www.who.int/news-room/fact-sheets/detail/dementia>
- Digby, G. J., Noetzel, M. J., Bubser, M., Utle, T. J., Walker, A. G., Byun, N. E., Lebois, E. P., Xiang, Z., Sheffler, D. J., Cho, H. P., Davis, A. A., Nemirovsky, N. E., Mennenga, S. E., Camp, B. W., Bimonte-Nelson, H. A., Bode, J., Italiano, K., Morrison, R., Daniels, J. S., ... Conn, P. J. (2012). Novel Allosteric Agonists of M1 Muscarinic Acetylcholine Receptors Induce Brain Region-Specific Responses That Correspond with Behavioral Effects in Animal Models. *The Journal of Neuroscience*, 32(25), 8532–8544. <https://doi.org/10.1523/JNEUROSCI.0337-12.2012>
- Dwomoh, L., Tejada, G. S., & Tobin, A. B. (2022). Targeting the M1 muscarinic acetylcholine receptor in Alzheimer's disease. *Neuronal Signaling*, 6(1), NS20210004. <https://doi.org/10.1042/NS20210004>
- Fisher, A. (2008). Cholinergic treatments with emphasis on m1 muscarinic agonists as potential disease-modifying agents for Alzheimer's disease. *Neurotherapeutics: The Journal of the American Society for Experimental NeuroTherapeutics*, 5(3), 433–442. <https://doi.org/10.1016/j.nurt.2008.05.002>
- Flynn, D. D., Weinstein, D. A., & Mash, D. C. (1991). Loss of high-affinity agonist binding to M1 muscarinic receptors in Alzheimer's disease: Implications for the failure of cholinergic replacement therapies. *Annals of Neurology*, 29(3), 256–262. <https://doi.org/10.1002/ana.410290305>
- Forlenza, O. V., Spink, J. M., Dayanandan, R., Anderton, B. H., Olesen, O. F., & Lovestone, S. (2000). Muscarinic agonists reduce tau phosphorylation in non-neuronal cells via

- GSK-3 $\beta$  inhibition and in neurons. *Journal of Neural Transmission*, 107(10), 1201–1212.  
<https://doi.org/10.1007/s007020070034>
- Foster, D. J., Choi, D. L., Conn, P. J., & Rook, J. M. (2014). Activation of M1 and M4 muscarinic receptors as potential treatments for Alzheimer's disease and schizophrenia. *Neuropsychiatric Disease and Treatment*, 10, 183–191.  
<https://doi.org/10.2147/NDT.S55104>
- Frost, B., Jacks, R. L., & Diamond, M. I. (2009). Propagation of tau misfolding from the outside to the inside of a cell. *The Journal of Biological Chemistry*, 284(19), 12845–12852.  
<https://doi.org/10.1074/jbc.M808759200>
- Haense, C., Kalbe, E., Herholz, K., Hohmann, C., Neumaier, B., Kraiss, R., & Heiss, W.-D. (2012). Cholinergic system function and cognition in mild cognitive impairment. *Neurobiology of Aging*, 33(5), 867–877.  
<https://doi.org/10.1016/j.neurobiolaging.2010.08.015>
- Hempel, H., Elhage, A., Cho, M., Apostolova, L. G., Nicoll, J. A. R., & Atri, A. (2023). Amyloid-related imaging abnormalities (ARIA): Radiological, biological and clinical characteristics. *Brain*, 146(11), 4414–4424. <https://doi.org/10.1093/brain/awad188>
- Hardy, J. A., & Higgins, G. A. (1992). Alzheimer's Disease: The Amyloid Cascade Hypothesis. *Science*, 256(5054), 184–185. <https://doi.org/10.1126/science.1566067>
- Hill, J. M., Bhattacharjee, P. S., & Neumann, D. M. (2007). Apolipoprotein E alleles can contribute to the pathogenesis of numerous clinical conditions including HSV-1 corneal disease. *Experimental Eye Research*, 84(5), 801–811.  
<https://doi.org/10.1016/j.exer.2006.08.001>
- Hogan, D. B. (2014). Long-Term Efficacy and Toxicity of Cholinesterase Inhibitors in the Treatment of Alzheimer Disease. *Canadian Journal of Psychiatry. Revue Canadienne de Psychiatrie*, 59(12), 618–623. <https://doi.org/10.1177/070674371405901202>

- Hoskin, J. L., Al-Hasan, Y., & Sabbagh, M. N. (2018). Nicotinic Acetylcholine Receptor Agonists for the Treatment of Alzheimer's Dementia: An Update. *Nicotine & Tobacco Research*, 21(3), 370–376. <https://doi.org/10.1093/ntr/nty116>
- Jones, C. K., Brady, A. E., Davis, A. A., Xiang, Z., Bubser, M., Tantawy, M. N., Kane, A. S., Bridges, T. M., Kennedy, J. P., Bradley, S. R., Peterson, T. E., Ansari, M. S., Baldwin, R. M., Kessler, R. M., Deutch, A. Y., Lah, J. J., Levey, A. I., Lindsley, C. W., & Conn, P. J. (2008). Novel Selective Allosteric Activator of the M1 Muscarinic Acetylcholine Receptor Regulates Amyloid Processing and Produces Antipsychotic-Like Activity in Rats. *The Journal of Neuroscience*, 28(41), 10422–10433. <https://doi.org/10.1523/JNEUROSCI.1850-08.2008>
- Keov, P., López, L., Devine, S. M., Valant, C., Lane, J. R., Scammells, P. J., Sexton, P. M., & Christopoulos, A. (2014). Molecular Mechanisms of Bitopic Ligand Engagement with the M1 Muscarinic Acetylcholine Receptor. *The Journal of Biological Chemistry*, 289(34), 23817–23837. <https://doi.org/10.1074/jbc.M114.582874>
- Kruse, A. C., Hu, J., Pan, A. C., Arlow, D. H., Rosenbaum, D. M., Rosemond, E., Green, H. F., Liu, T., Chae, P. S., Dror, R. O., Shaw, D. E., Weis, W. I., Wess, J., & Kobilka, B. K. (2012). Structure and dynamics of the M3 muscarinic acetylcholine receptor. *Nature*, 482(7386), 552–556. <https://doi.org/10.1038/nature10867>
- Kruse, A. C., Ring, A. M., Manglik, A., Hu, J., Hu, K., Eitel, K., Hübner, H., Pardon, E., Valant, C., Sexton, P. M., Christopoulos, A., Felder, C. C., Gmeiner, P., Steyaert, J., Weis, W. I., Garcia, K. C., Wess, J., & Kobilka, B. K. (2013). Activation and allosteric modulation of a muscarinic acetylcholine receptor. *Nature*, 504(7478), 101–106. <https://doi.org/10.1038/nature12735>
- Lange, K. W., Javoy-Agid, F., Agid, Y., Jenner, P., & Marsden, C. D. (1992). Brain muscarinic cholinergic receptors in Huntington's disease. *Journal of Neurology*, 239(2), 103–104. <https://doi.org/10.1007/BF00862983>

- Langmead, C. J., Austin, N. E., Branch, C. L., Brown, J. T., Buchanan, K. A., Davies, C. H., Forbes, I. T., Fry, V. a. H., Hagan, J. J., Herdon, H. J., Jones, G. A., Jeggo, R., Kew, J. N. C., Mazzali, A., Melarange, R., Patel, N., Pardoe, J., Randall, A. D., Roberts, C., ... Watson, J. (2008). Characterization of a CNS penetrant, selective M1 muscarinic receptor agonist, 77-LH-28-1. *British Journal of Pharmacology*, *154*(5), 1104–1115. <https://doi.org/10.1038/bjp.2008.152>
- Levey, A., Kitt, C., Simonds, W., Price, D., & Brann, M. (1991). Identification and localization of muscarinic acetylcholine receptor proteins in brain with subtype-specific antibodies. *The Journal of Neuroscience*, *11*(10), 3218–3226. <https://doi.org/10.1523/JNEUROSCI.11-10-03218.1991>
- Lin, L., Georgievska, B., Mattsson, A., & Isacson, O. (1999). Cognitive changes and modified processing of amyloid precursor protein in the cortical and hippocampal system after cholinergic synapse loss and muscarinic receptor activation. *Proceedings of the National Academy of Sciences*, *96*(21), 12108–12113. <https://doi.org/10.1073/pnas.96.21.12108>
- Malenka, R. C., & Bear, M. F. (2004). LTP and LTD: An embarrassment of riches. *Neuron*, *44*(1), 5–21. <https://doi.org/10.1016/j.neuron.2004.09.012>
- Marino, M. J., Rouse, S. T., Levey, A. I., Potter, L. T., & Conn, P. J. (1998). Activation of the genetically defined m1 muscarinic receptor potentiates N-methyl-D-aspartate (NMDA) receptor currents in hippocampal pyramidal cells. *Proceedings of the National Academy of Sciences of the United States of America*, *95*(19), 11465–11470. <https://doi.org/10.1073/pnas.95.19.11465>
- Mash, D. C., Flynn, D. D., & Potter, L. T. (1985). Loss of M2 muscarine receptors in the cerebral cortex in Alzheimer's disease and experimental cholinergic denervation. *Science (New York, N.Y.)*, *228*(4703), 1115–1117. <https://doi.org/10.1126/science.3992249>

*Medications for Memory, Cognition & Dementia-Related Behaviors* | *alz.org*. (n.d.). Alzheimer's Association. Retrieved July 30, 2025, from

<https://www.alz.org/alzheimers-dementia/treatments/medications-for-memory>

Mitoh, Y., Ueda, H., Ichikawa, H., Fujita, M., Kobashi, M., & Matsuo, R. (2017). Effects of cevimeline on excitability of parasympathetic preganglionic neurons in the superior salivatory nucleus of rats. *Autonomic Neuroscience*, *206*, 1–7.

<https://doi.org/10.1016/j.autneu.2017.05.010>

Moran, S. P., Dickerson, J. W., Cho, H. P., Xiang, Z., Maksymetz, J., Remke, D. H., Lv, X., Doyle, C. A., Rajan, D. H., Niswender, C. M., Engers, D. W., Lindsley, C. W., Rook, J. M., & Conn, P. J. (2018). M1-positive allosteric modulators lacking agonist activity provide the optimal profile for enhancing cognition. *Neuropsychopharmacology*, *43*(8),

1763–1771. <https://doi.org/10.1038/s41386-018-0033-9>

Nathan, P. J., Millais, S. B., Godwood, A., Dewit, O., Cross, D. M., Liptrot, J., Ruparelia, B., Jones, S. P., Bakker, G., Maruff, P. T., Light, G. A., Brown, A. J. H., Weir, M. P., Congreve, M., & Tasker, T. (2022). A phase 1b/2a multicenter study of the safety and preliminary pharmacodynamic effects of selective muscarinic M1 receptor agonist HTL0018318 in patients with mild-to-moderate Alzheimer's disease. *Alzheimer's & Dementia : Translational Research & Clinical Interventions*, *8*(1), e12273.

<https://doi.org/10.1002/trc2.12273>

Nielsen, B. E., & Ford, C. P. (2024). Reduced striatal M4-cholinergic signaling following dopamine loss contributes to parkinsonian and L-DOPA-induced dyskinetic behaviors. *Science Advances*, *10*(47), eadp6301. <https://doi.org/10.1126/sciadv.adp6301>

Noetzel, M. J., Grant, M. K. O., & El-Fakahany, E. E. (2009). Immediate and Delayed Consequences of Xanomeline Wash-Resistant Binding at the M3 Muscarinic Receptor. *Neurochemical Research*, *34*(6), 1138–1149. <https://doi.org/10.1007/s11064-008-9886-3>

- Oleksak, P., Novotny, M., Patocka, J., Nepovimova, E., Hort, J., Pavlik, J., Klimova, B., Valis, M., & Kuca, K. (2021). Neuropharmacology of Cevimeline and Muscarinic Drugs—Focus on Cognition and Neurodegeneration. *International Journal of Molecular Sciences*, *22*(16), 8908. <https://doi.org/10.3390/ijms22168908>
- Olson, W. C., & Jacobson, J. M. (2009). CCR5 Monoclonal Antibodies for HIV-1 Therapy. *Current Opinion in HIV and AIDS*, *4*(2), 104–111. <https://doi.org/10.1097/COH.0b013e3283224015>
- Pimlott, S. L., Piggott, M., Owens, J., Grealley, E., Court, J. A., Jaros, E., Perry, R. H., Perry, E. K., & Wyper, D. (2004). Nicotinic Acetylcholine Receptor Distribution in Alzheimer's Disease, Dementia with Lewy Bodies, Parkinson's Disease, and Vascular Dementia: In Vitro Binding Study Using 5-[125I]-A-85380. *Neuropsychopharmacology*, *29*(1), 108–116. <https://doi.org/10.1038/sj.npp.1300302>
- Poslunsey, M. S., Wood, M. R., Han, C., Stauffer, S. R., Panarese, J. D., Melancon, B. J., Engers, J. L., Dickerson, J. W., Peng, W., Noetzel, M. J., Cho, H. P., Rodriguez, A. L., Hopkins, C. R., Morrison, R., Crouch, R. D., Bridges, T. M., Blobaum, A. L., Boutaud, O., Daniels, J. S., ... Lindsley, C. W. (2024). Discovery of VU0467319: An M1 Positive Allosteric Modulator Candidate That Advanced into Clinical Trials. *ACS Chemical Neuroscience*, *16*(1), 95–107. <https://doi.org/10.1021/acscemneuro.4c00769>
- Ramos Bernardes da Silva Filho, S., Oliveira Barbosa, J. H., Rondinoni, C., dos Santos, A. C., Garrido Salmon, C. E., da Costa Lima, N. K., Ferriolli, E., & Moriguti, J. C. (2017). Neuro-degeneration profile of Alzheimer's patients: A brain morphometry study. *NeuroImage: Clinical*, *15*, 15–24. <https://doi.org/10.1016/j.nicl.2017.04.001>
- Ren, S., Singh, J., Gsteiger, S., Cogley, C., Reed, B., Abrams, K. R., Dawoud, D., Owen, R. K., Tappenden, P., Quinn, T. J., & Bujkiewicz, S. (2025). *Evaluating amyloid-beta as a surrogate endpoint in trials of anti-amyloid drugs in Alzheimer's disease: A Bayesian meta-analysis* (No. arXiv:2504.06807). arXiv. <https://doi.org/10.48550/arXiv.2504.06807>

- Schuff, N., Woerner, N., Boreta, L., Kornfield, T., Shaw, L. M., Trojanowski, J. Q., Thompson, P. M., Jack, C. R., Weiner, M. W., & Alzheimer's Disease Neuroimaging Initiative. (2009). MRI of hippocampal volume loss in early Alzheimer's disease in relation to ApoE genotype and biomarkers. *Brain: A Journal of Neurology*, *132*(Pt 4), 1067–1077. <https://doi.org/10.1093/brain/awp007>
- Shirey, J. K., Brady, A. E., Jones, P. J., Davis, A. A., Bridges, T. M., Kennedy, J. P., Jadhav, S. B., Menon, U. N., Xiang, Z., Watson, M. L., Christian, E. P., Doherty, J. J., Quirk, M. C., Snyder, D. H., Lah, J. J., Levey, A. I., Nicolle, M. M., Lindsley, C. W., & Conn, P. J. (2009). A Selective Allosteric Potentiator of the M1 Muscarinic Acetylcholine Receptor Increases Activity of Medial Prefrontal Cortical Neurons and Restores Impairments in Reversal Learning. *The Journal of Neuroscience*, *29*(45), 14271–14286. <https://doi.org/10.1523/JNEUROSCI.3930-09.2009>
- Smiley, J. F., Levey, A. I., & Mesulam, M.-M. (1999). M2 Muscarinic receptor immunolocalization in cholinergic cells of the monkey basal forebrain and striatum. *Neuroscience*, *90*(3), 803–814. [https://doi.org/10.1016/S0306-4522\(98\)00527-2](https://doi.org/10.1016/S0306-4522(98)00527-2)
- Smith, K. S., Bucci, D. J., Luikart, B. W., & Mahler, S. V. (2016). DREADDs: Use and Application in Behavioral Neuroscience. *Behavioral Neuroscience*, *130*(2), 137–155. <https://doi.org/10.1037/bne0000135>
- Thal, D. M., Sun, B., Feng, D., Nawaratne, V., Leach, K., Felder, C. C., Bures, M. G., Evans, D. A., Weis, W. I., Bachhawat, P., Kobilka, T. S., Sexton, P. M., Kobilka, B. K., & Christopoulos, A. (2016). Crystal structures of the M1 and M4 muscarinic acetylcholine receptors. *Nature*, *531*(7594), 335–340. <https://doi.org/10.1038/nature17188>
- Voss, T., Li, J., Cummings, J., Farlow, M., Assaid, C., Froman, S., Leibensperger, H., Snow-Adami, L., McMahon, K. B., Egan, M., & Michelson, D. (2018). Randomized, controlled, proof-of-concept trial of MK-7622 in Alzheimer's disease. *Alzheimer's &*

*Dementia: Translational Research & Clinical Interventions*, 4(1), 173–181.

<https://doi.org/10.1016/j.trci.2018.03.004>

Wess, J. (1996). Molecular biology of muscarinic acetylcholine receptors. *Critical Reviews in Neurobiology*, 10(1), 69–99. <https://doi.org/10.1615/critrevneurobiol.v10.i1.40>

Zhang, J., Zhang, Y., Wang, J., Xia, Y., Zhang, J., & Chen, L. (2024). Recent advances in Alzheimer's disease: Mechanisms, clinical trials and new drug development strategies. *Signal Transduction and Targeted Therapy*, 9(1), 211.

<https://doi.org/10.1038/s41392-024-01911-3>

Zuchner, T., Schliebs, R., & Perez-Polo, J. R. (2005). Down-regulation of muscarinic acetylcholine receptor M2 adversely affects the expression of Alzheimer's disease-relevant genes and proteins. *Journal of Neurochemistry*, 95(1), 20–32.

<https://doi.org/10.1111/j.1471-4159.2005.03335.x>

## **EDITOR COMMENTS AND RECOMMENDATION:**

The author presents a thorough review of targeting muscarinic receptors for Alzheimer's Disease, including the use of four figures which aid in the presentation of the data and analysis. Reviewer 2 specifically highlights several points which require revision prior to acceptance of the manuscript. First, the reviewer identifies several inaccuracies in Figure 4 that require correction. Additionally, the reviewer suggests that the author elaborate further in the Limitations section to include a discussion of the use of animal models as proxies for human neurodegenerative disease, with the reviewer noting that this critical analysis will demonstrate the author's ability to bridge the basic and translational studies described with clinical applicability and relevance. Given these critiques, the editor suggests that the manuscript undergo a major revision prior to publication.

## **REVIEWER 1**

### **Final Recommendation: Accept with Minor Revisions**

#### 1. Originality & Significance

This is a highly impressive and ambitious review for a high school journal. The author provides a comprehensive and nuanced synthesis of research developments targeting muscarinic acetylcholine receptors (mAChRs) in Alzheimer's disease (AD), with a clear emphasis on translational relevance and therapeutic mechanisms.

The manuscript makes a valuable contribution by:

- Framing the history of cholinergic hypotheses and shifting focus toward mAChR subtypes.
- Describing orthosteric ligands, allosteric modulators, and bitopic ligands with clarity.
- Connecting receptor pharmacology to broader neurobiological pathways, including glutamatergic, dopaminergic, and tau/amyloid systems.
- Including a well-structured summary of ongoing trials and future directions, such as nanobody, DREADD, and PET tracer applications.

This is one of the strongest papers I've seen at the high school level in terms of biological depth, scope, and writing.

Suggestions:

- Consider adding keywords to the abstract, per standard academic formatting.

#### 2. Clarity & Structure

The paper is exceptionally well-structured, particularly for a student submission. Section transitions are smooth, paragraph development is logical, and technical vocabulary is clearly explained. There is a thoughtful narrative flow from foundational research through therapeutic development and into translational applications.

#### Minor Formatting Suggestions:

- The first paragraph of the Introduction needs slight formatting cleanup (possibly a missing line break between lines 5–6).
- Consider simplifying the phrase “Clinical Dementia Rating – Sum of Boxes scores” in the Introduction.
- Consider adding brief clarifying sentences that connect receptor-level mechanisms to patient-level outcomes (e.g., how M1 modulation might improve memory or reduce behavioral symptoms). These would help reinforce the paper’s translational relevance and strengthen the connection between scientific detail and therapeutic goals.
- Table 1 formatting: The drug development table is dense and helpful, but formatting consistency could be improved. You should follow the specific APA guidelines. Consider:
  - Aligning columns visually
  - Using abbreviations more sparingly (e.g. define “KO” again if used after a long gap),
  - Possibly moving preclinical compounds into a supplementary table to make the main one more digestible
  -

### 3. Use of Evidence & Research Methods

As this is a literature review rather than a primary research paper, the focus is appropriately on summarizing prior studies. The citations are extensive and well-chosen, and figures are clearly described.

#### Citation Formatting Suggestions:

- Ensure all citations are placed before punctuation marks, as required by APA. • Ensure figures created in BioRender are compliant with their open-access license. • Ensure Figure 2 is compliant with the Allen Institute’s citation guidelines • Ensure Figure 4 is compliant with the RCSB Protein Data Bank citation policies • *Dementia (n.d.)* reference from WHO should be cited correctly as a fact sheet. Note also that the publication year is available (2025). Same is true for “*Medications for Memory, Cognition & Dementia-Related Behaviors*” reference. Refer to these guidelines for both and update the citations according to APA guidelines: <https://apastyle.apa.org/style-grammar-guidelines/references/examples/webpage-website-references#4>

### 4. Engagement with Literature

The student demonstrates exceptional engagement with the relevant literature. Key concepts are defined and placed in historical context. The discussion of:

- mAChR structure and GPCR signalling
- Subtype selectivity,
- Limitations of prior trials, and
- Advanced therapeutic approaches (e.g., biased signalling, nanobodies), ...reflects a high level of understanding and critical integration.
-

Suggestions:

- Add brief mention of how the review expands upon or contrasts with earlier reviews on similar topics.

## 5. Grammar & Language

The writing is clean, precise, and mature. Sentences are well-formed, definitions are appropriately placed, and clarity is consistent throughout.

Minor Suggestions:

- Standardize formatting for abbreviations.
- Correct minor typo on page 9, second paragraph, last sentence

Concluding Remarks:

This paper is among the most impressive high school-level reviews I've read. It synthesizes complex biochemical, pharmacological, and clinical content with clarity and precision. The author demonstrates both intellectual maturity and strong science communication skills. Only minor revisions are needed, mainly stylistic consistency and formatting corrections.

## REVIEWER 2

Overall: Author has written a very comprehensive coverage of muscarinic receptors, and the drugs/compounds targeting these receptors in the context of Alzheimer's disease. Furthermore, the paper is strengthened by the addition of 4 figures that further illustrate the points made in the review. In addition, the author has discussed in detail the background and structure of the receptors, demonstrating an excellent familiarity with literature of the field.

However, there is some inaccuracies with the Figure 4 that the author needs to clear up before the paper should be accepted.

There is also concern that AI was used in the writing of this paper. This reviewer would like to suggest that the editor do an independent AI check on the writing.

Points to address:

1. Abstract: "Then, analyze the status of developments in different classes of treatments based on what stages in trial they were discontinued or are currently in, data regarding efficacy, pathways targeted, and adverse effects." This sentence is grammatically incomplete.
2. The first paragraph of introduction has unusual formatting and margins, but this could possibly be due to the conversion to a pdf document.

3. "Given this current landscape of Alzheimer's research and its treatments, this paper aims to synthesize past research to more clearly identify the potential roles muscarinic receptors can play in the progress of the field." I do not think "synthesize" is the appropriate word to use here; in fact, it muddies the intent of this sentence.

4. "Then, analyze the pathways that have been researched as a result and the status of drug developments and therapeutics that have attempted to target them." Again, this sentence is grammatically incomplete.

5. "Togethe,r these mechanisms potentiate NMDAR currents and facilitate LTP, making M1 a very useful target in countering this aspect of AD progression." There is a typing error in this sentence.

6. Figure 4, major point. This figure is meant to illustrate the protein interactions with the ligand (in this case a lipid), but when I looked it up on RCSB, the detergent is also in the structure. The author needs to label clearly in the figure.

7. Figure 4, major point. More clarifications needed. The author needs to label clearly the intracellular and extracellular regions.

7. Figure 4, minor point. Please specific the what the red and green models are.

8. Figure 4: Also, what is Chymol? Do you mean PyMol?

10. Figure 4, if these are PDB structures from the database, the PDB ID needs to be explicitly cited, so the reader can look up the structures as well.

11. The author has written extensively on the mode of action of different compounds/drugs. Since muscarinic receptors are found throughout the body, it would enrich the paper further if the author can elaborate on side effects from various compounds, especially those with potential human clinical usage. There was some mention of increase blood pressure from selective M1 agonist HTL0018318, but more information on other compounds would increase the depth of this review.

12. Under Limitations, this reviewer would like to see the author elaborate on inconsistencies or pitfalls of using animal models to model a human neurodegenerative disease, as basic research uses mouse models of AD. Even a short paragraph will go a long way to demonstrate the author's ability to bridge experimental models and clinical studies.

13. Throughout the paper, there are a few instances of grammatical inconsistencies. Please amend before final submission.

# Developments in Treatments Targeting Muscarinic Receptors for Alzheimer's Disease

## Abstract:

Alzheimer's Disease (AD) is the most common neurodegenerative disease to date and affects over 50 million patients worldwide, thus making it a major area of focus in drug discovery. Muscarinic Receptors, especially the M1 and M4 subtypes, are promising therapeutic targets due to their selectivity, high expression, and function in the central nervous system. Although the focus of research in AD has shifted away from muscarinic receptors in the past due to difficulties involving receptor selectivity, adverse effects, and failure in clinical trials, new discoveries in the functions of these receptors have opened up the possibility for the development of treatments with better efficacy and less risk, resulting in a resurgence of interest in them. This review aims to first examine previous research and data that have helped us recognize the functions of muscarinic receptors in AD, specifically where they are expressed in the brain, their intracellular mechanisms, and how these can help us find pathways to treat AD. It then analyzes the status of developments in different classes of treatments based on performance in trials, efficacy data, pathways targeted, and adverse effects. Finally, we identify potential areas for further discovery and development of pharmaceuticals by highlighting the flaws of previous treatments, unknown aspects of AD pathology, and how more recent and less studied areas of research could guide future development and discovery.

**Keywords:** Muscarinic Receptors, Alzheimer's Disease, Cholinergic Dysfunction, Amyloid Pathology, Tau Pathology, Literature Review

## Introduction:


Alzheimer's Disease (AD) is a progressive neurodegenerative disorder characterized by memory loss, cognitive impairment, and behavioral changes. It is responsible for 60-70% of cases of dementia, which is a leading cause of death, disability, and dependency among the older population (World Health Organization, 2025). Despite its burden, current treatments are limited and not effective enough to fully treat AD. Over the past few decades, various hypotheses for AD have developed. Today, widespread understanding of AD pathogenesis has stemmed from the cholinergic, amyloid cascade, and tau hypotheses, in addition to less recognized theories like the calcium homeostasis hypothesis (Zhang et al., 2024).

First proposed in 1979, the cholinergic hypothesis is one of the earliest models of AD formation. This hypothesis linked a deficiency in acetylcholine (ACh) with the cognitive effects experienced by individuals with Alzheimer's (Davies & Maloney, 1976). Acetylcholine is a neurotransmitter or chemical messenger that is synthesized in a neuron, released in the space between two neurons called a synaptic cleft, and then binds to a receptor. The hypothesis attributed the shortage in ACh to the loss of cholinergic neurons—neurons that use ACh to communicate—found in AD patients. These changes occur mostly in the hippocampus, cortex, and other cortical regions important for cognitive function. This discovery influenced the development of

treatments that increased ACh levels in the brain, like acetylcholinesterase inhibitors (AChEIs). AChEIs work by inhibiting the enzyme acetylcholinesterase (AChE), which breaks down acetylcholine, therefore increasing acetylcholine levels in the brain and improving synaptic function, specifically in the cholinergic system. Currently, Donepezil, Rivastigmine, and Galantamine are FDA-approved AChEIs in clinical use (Alzheimer's Association, 2025). These drugs are often used in conjunction with an NMDA antagonist, Memantine. However, AChEIs plateau in their effectiveness as AD becomes more severe, calling for alternative means of increasing ACh levels (Hogan, 2014).

Muscarinic receptors (mAChRs) and Nicotinic receptors (nAChRs)— the two types of acetylcholine receptors— have built on the cholinergic hypothesis, showing lots of potential as effective targets of therapeutics for AD. This review focuses on the developments in drugs targeting mAChRs. Muscarinic receptors are a family of class A G protein-coupled receptors (GPCRs) subdivided into 5 subtypes (M1-M5), which bind to and are activated by ACh in the central and peripheral nervous systems, a mechanism significant to the cholinergic system. Although no treatments targeting them have been approved yet, they have regained increasing interest in Alzheimer's research. Previously, research in the field shifted away from targeting muscarinic receptors for a period and instead focused on targeting biomarkers associated with AD. More specifically, on amyloid-beta ( $A\beta$ ) plaques and neurofibrillary tangles (NFTs).  $A\beta$  plaques are abnormal accumulations of  $A\beta$  peptides, which then trigger a series of pathways, further progressing the disease (Hardy & Higgins, 1992). One of these pathways is the hyperphosphorylation of tau proteins, leading to misfolded tau proteins that form NFTs, hindering synaptic function (Frost et al., 2009). Understanding of these pathways has come mainly from the amyloid cascade and tau hypotheses. Treatments to come out of these discoveries include anti-amyloid therapies like Aducanumab, Lecanemab, and Donanemab (Alzheimer's Association, 2025). However, these treatments also had limited efficacy. For instance, Lecanemab showed a 27% slower decline in CDR-SB (Clinical Dementia Rating – Sum of Boxes) scores over 18 months compared to placebo, which is statistically meaningful but modest, on top of having adverse effects like brain swelling or bleeding (Hempel et al., 2023; Ren et al., 2025). Furthermore, by the time amyloid plaques accumulate, neurodegeneration has already advanced, limiting their ability to reverse damage. Nevertheless, the cholinergic, tau, and amyloid hypotheses have helped us gain a better idea of the mechanisms and pathways behind AD [Figure 1]. Most notably, the M1 and M4 receptors have frequently been researched for their role in cholinergic dysfunction and relationship with amyloid pathways (Davis et al., 2010; Foster et al., 2014).

## Cholinergic Dysfunction in AD: Comparison



**Figure 1:** (Created with BioRender) In healthy conditions, acetylcholine (ACh) is efficiently released from presynaptic neurons and inhibited by M4 muscarinic receptors—particularly in forebrain regions such as the hippocampus and cortex—leading to substantial activation of M1 muscarinic receptors and regulation by acetylcholinesterase (AChE), allowing for normal cognitive function. In AD, degeneration of basal forebrain cholinergic neurons leads to reduced cholinergic neuron density, lower ACh levels, and ultimately less muscarinic activity. This dysfunction contributes to the development of Aβ plaques and NFTs, which collectively contribute to the progressive cognitive decline associated with Alzheimer’s disease. (BioRender, n.d.; Davis et al., 2010; Foster et al., 2014)

Given this current landscape of Alzheimer’s research and its treatments, this literature review aims to analyze past research to more clearly identify the potential roles muscarinic receptors can play in the progress of the field. It will first further examine the foundational studies that identified the functions, distributions, and intracellular mechanisms of mAChRs in the context of AD. The paper will then examine both the pathways that have been researched as a result, and the status of drug developments and therapeutics that have attempted to target them. Finally, it will identify more recent and less studied discoveries that could hold solutions and discoveries in the field.

## Foundational Discoveries of mAChR Roles in AD

Many aspects of Alzheimer's pathology are still unknown; however, their understanding has come a long way since the proposal of the cholinergic hypothesis. As more pathways and locations of biomarkers involved in specific stages of AD have been identified, the roles mAChRs play have become clearer. While multiple factors impact their significance to AD pathology, their structures and distributions provided the fundamental rationale for their targeting in treatments.

### **Distribution and Concentration of mAChRs in Areas Impacted by Alzheimer's**

In its early stages, AD impacts areas significant to memory in the temporal lobe, specifically the hippocampus and entorhinal cortex, as well as the posterior cingulate gyrus, amygdala, and parahippocampal gyrus. Using MRIs, atrophy, brain shrinkage due to loss of neurons and synapses, has been found in these areas (Schuff et al., 2009). As the disease progresses, it expands to other parts of the cerebral cortex, like the frontal and parietal lobes (Ramos Bernardes da Silva Filho et al., 2017).


The M1 receptor, the most studied mAChR in AD research, is found throughout the central nervous system (CNS) in the cerebral cortex, hippocampus, and striatum, as well as the autonomic ganglia in the peripheral nervous system (PNS). The CNS consists of the brain and the spinal cord, acting as the control center. The PNS, made up of the nerves outside of the CNS, acts as the communication network that connects the CNS to the rest of the body. M1 is most densely distributed in layers II/III and V in the frontal, parietal, and temporal cortex and the hippocampus (Levey et al., 1991). Due to its prominence in these memory circuits, it is the most commonly targeted muscarinic receptor in the CNS for AD research and treatment. While the number of M1 receptors has been found to be unaffected by AD, their signaling efficacy is impaired, most likely attributed to impaired coupling with G proteins (Caccamo et al., 2006; Flynn et al., 1991).

Showing the highest overlap with the M1 receptor in the regions experiencing early neurodegeneration in AD, M4 receptors are mostly expressed in the caudate nucleus, putamen, and nucleus accumbens of the basal ganglia, as well as in the hippocampus and neocortex. Their high expression in the basal ganglia in particular is significant because they form part of the cortico–basal ganglia–thalamic loops, which integrate cognitive, motor, and reward-related information. In Alzheimer's, impairments in these circuits lead to cognitive impairment as well as neuropsychiatric symptoms such as apathy, agitation, and impaired decision-making. Similarly, M2 is an autoreceptor expressed in the hippocampus, cerebral cortex, and presynaptic terminals throughout the CNS, as well as in autonomic nuclei of the brainstem (Levey et al., 1991). While not yet proved to be as effective a target due to its primary role being in the PNS, it provides major cholinergic pathways to the cortex and hippocampus (Smiley et al., 1999). Acting as inhibitory autoreceptors regulating ACh release, its loss in Alzheimer's patients contributes significantly to the cholinergic deficits associated with AD.

Finally, while the M3 and M5 receptors are mostly distributed in the PNS, they have displayed some therapeutic potential as targets. In neurons, M3 enables excitatory cholinergic signaling

and plays a role in neurovascular coupling, mediating vasodilation—the widening of blood vessels— through Gq-protein pathways in endothelial cells of the cerebral vascular system (Kruse et al., 2012). The M5 receptor, while the least abundant muscarinic subtype in the CNS, also regulates cerebral blood flow. Hence, the M3 and M5 receptors' ability to manage cerebral blood flow could make them valuable targets in early disease stages.

## Distribution of mAChR Subtypes in the Human Brain




**Figure 2. Distribution of muscarinic acetylcholine receptor (mAChR) subtypes in the human brain:** Coronal, axial, sagittal, and 3D projections from the Allen Human Brain Atlas showing gene expression patterns of mAChR subtypes in the areas they are most heavily distributed in (The brighter the red, the more concentrated). Allen Human Brain Atlas, <https://human.brain-map.org> and <https://atlas.brain-map.org>.

- A. M1 expression in the cerebral cortex
  - B. M2 expression in the pons
  - C. M3 expression in the hypothalamus
  - D. M4 expression in the striatum
  - E. M5 expression in the pons
- (Allen Institute for Science, n.d.)

### Structure of mAChRs

Since mAChRs are GPCRs, all 5 subtypes have seven transmembrane (7TM) helices connected by extracellular and intracellular loops with an extracellular N-terminus and intracellular C-terminus. The orthosteric, or primary, binding site for ACh is located deep inside the 7TM bundle, which allows for precise ligand binding, conformational change—change in receptor shape after binding—to initiate intracellular signaling through heterotrimeric G proteins. The G protein that each mAChR couples with influences the pathways they signal. The M1, M3, and M5 receptors couple to Gq/11, which activates the phospholipase C (PLC)—IP<sub>3</sub>—Ca<sup>2+</sup> signaling cascade that promotes neuronal excitability and synaptic plasticity. On the other hand, the M2 and M4 receptors couple to Gi/o, which inhibit adenylyl cyclase and modulate ion channel activity to regulate neurotransmitter release. GPCR structure makes mAChRs a more ideal target for therapeutics than the other family of acetylcholine receptors, nAChRs. Unlike nAChRs, which are ligand-gated ion channels with rapid but less modulatable responses, mAChRs transduce signals through G protein-mediated cascades. This allows for a broader range of pharmacological modulation, like biased agonism, allosteric regulation, and subtype selectivity (Wess, 1996). Furthermore, research using high-resolution crystallography and cryo-electron microscopy has discovered pocket structures and loop conformations that are

unique to each mAChR subtype.



**Figure 3:** (Created with BioRender) M1, M3, and M5 receptors couple to Gq/11 proteins, activating phospholipase C (PLC) and leading to production of IP<sub>3</sub> and second messenger Diacylglycerol (DAG), which increase intracellular Ca<sup>2+</sup> and activate protein kinase C (PKC). M2 and M4 receptors couple to Gi/o proteins, which inhibit adenylate cyclase, decrease cAMP levels, and activate G protein-coupled receptor kinase (GIRK) channels, resulting in membrane hyperpolarization. (*BioRender*, n.d.)

## Connections to Other Pathways and Diseases

In addition to being central to cholinergic neurotransmission, mAChRs also intersect with multiple pathological mechanisms relevant to AD and other neurological disorders. Understanding these cross-pathway interactions is critical, as it helps create the bigger picture, allowing for the development of therapies with greater efficacy and less adverse effects.

### Amyloid and Tau Pathologies

As mentioned before, muscarinic receptors influence the development of amyloid and tau pathologies [Figure 1]. Early research *in vitro* demonstrated that muscarinic stimulation increases sAPP $\alpha$  release, which inhibits and suppresses A $\beta$  generation, an effect blocked by non-selective muscarinic antagonism and PKC inhibition (Lin et al., 1999). The M1 receptor, however, has a direct effect as it modulates enzymes involved with the amyloid precursor protein (APP) processing activation. More specifically, it promotes non-amyloidogenic processing of amyloid precursor protein (APP) through stimulation of alpha-secretase, reducing the production of pathogenic A $\beta$  peptides (Cisse et al., 2011; Fisher, 2008). *In vivo*, the 3xTg-AD mouse model, chronic treatment with the M1 agonist AF267B increased cortical and hippocampal sAPP $\alpha$  through ADAM17 activation, reduced A $\beta$ 42 levels and plaque burden, and

improved memory (Caccamo et al., 2006). In addition to rodent models, AF267B also lowered CSF A $\beta$  in rabbits (Beach et al., 2001).

In parallel, M1 receptor stimulation also helps decrease the tau pathology associated with AD. Through PLC/Ca<sup>2+</sup>-dependent protein kinase C signaling and downstream inhibition of GSK3 $\beta$ , M1 activation reduces tau phosphorylation in vitro and in vivo. In vitro, muscarinic activation modulated tau kinase cascades that included PKC and mitogen-activated protein kinase (MAPK), impacting sites relevant to tangle formation (Forlenza et al., 2000). Additionally, in the same study, which found that M1 activation by AF267B in 3xTg-AD mice decreased amyloid pathology, AF267B treatment also decreased hyperphosphorylated tau. (Caccamo et al., 2006)

### **Links to Dopaminergic and Glutamatergic Systems**

In addition to cholinergic dysfunction, imbalances in the dopaminergic and glutamatergic systems significantly contribute to AD pathogenesis and progression. The dopaminergic system is primarily involved with brain functions like motivation, reward, and motor control, all of which are impacted by Alzheimer's. Muscarinic receptors, especially the M4 subtype, play essential roles in modulating dopaminergic transmissions. More specifically, activation of M4 limits dopamine release, making it a useful target for countering the dopamine imbalances associated with AD. Nicotinic receptors, which, unlike mAChRs, promote dopamine release through a quicker and more direct pathway, are also significant to dopamine balance. However, targeting nAChRs has been more challenging due to obstacles like downregulation in AD, receptor selectivity, receptor desensitization, and subtype selectivity. More specifically, in AD, nicotinic receptor expression of subtypes  $\alpha$ 4 $\beta$ 2 and  $\alpha$ 7 significantly diminished, reducing their efficacy as therapeutic targets (Hoskin et al., 2018; Pimlott et al., 2004). Furthermore, issues like receptor desensitization with chronic activation and lack of subtype selectivity can lead to off-target effects (Choi, 1988).

On the other hand, dysfunction in the glutamatergic system— the brain's primary excitatory network with glutamate as a neurotransmitter— is directly linked to the excitotoxicity and neuronal death found in AD. Among glutamate receptors, N-methyl-D-aspartate (NMDA) receptors are especially important because they are highly permeable to calcium and directly mediate synaptic plasticity, long-term potentiation (LTP), and memory encoding (Malenka & Bear, 2004). In Alzheimer's, NMDARs are overactivated and disrupted by A $\beta$  plaques, which is why antagonists like memantine are used clinically to reduce excitotoxicity. However, memantine only dampens overactivation and does not restore the physiological pro-plasticity role of NMDA receptors. M1 receptors modulate NMDAR currents heavily. In particular, M1 activation has been shown to enhance NMDA receptor-mediated excitatory postsynaptic currents in hippocampal pyramidal neurons (Marino et al., 1998). Additionally, M1 increases neuronal excitability and calcium entry through NMDA receptors by inhibiting small-conductance calcium-activated potassium (SK) channels (Conn et al., 2009). Together these mechanisms potentiate NMDAR currents and facilitate LTP, making M1 a very useful target in countering this aspect of AD progression.

## Connection to Other Diseases

In addition to AD, mAChRs have been proven to interact with many of the pathways in other neuropsychiatric and neurodegenerative disorders, such as Schizophrenia (SZ), Parkinson's, and Huntington's diseases. This has led to the testing of the same drugs for both SZ and AD, most notably xanomeline. Parkinson's, which is mainly characterized by the loss of dopaminergic neurons, has been associated with the loss of M4 receptors in the striatum due to their role in modulation (Nielsen & Ford, 2024). Similarly, a decrease in the density and total of M2 has been found in post-mortem Huntington patients (Lange et al., 1992). Although SZ is mostly classified as a neuropsychiatric disorder, multiple studies have demonstrated that M1 and M4 interact with many of the same neural circuits as AD. This has led to the testing of the same drugs for both SZ and AD, most notably xanomeline.

## The Progress and Current Status of mAChR-Targeted Therapeutics

Despite the advantageous aspects of muscarinic acetylcholine receptors identified previously, mAChRs, early attempts to develop effective therapeutics faced major obstacles. These limitations revolved mainly around the lack of available compounds that were selective enough to only target one mAChR subtype, peripheral cholinergic side effects, and, while broader than nicotinic modulation, the still narrow window for modulation of mAChRs. Fortunately, recent research has shifted to improving subtype selectivity, blood–brain barrier (BBB) penetration, and utilizing novel mechanisms that modulate receptors to achieve cognitive benefits that couldn't have been achieved before. Due to their role in AD pathology, M1 and M4 are usually the targets of these compounds.

### Subtype Selective Orthosteric Ligands

Orthosteric ligands target the acetylcholine binding site, deep inside the 7TM bundle. This site is common among all AChRs, making subtype selectivity challenging. For instance, Cevimeline (AF102B), developed as an orthosteric M1 agonist for AD, could cross the BBB and showed improved memory and reduced A $\beta$  pathology in rodents (Mitoh et al., 2017). In human trials, the drug exhibited the same benefits but also showed peripheral adverse effects, causing it to fail FDA approval (Oleksak et al., 2021). An older agent, which revealed that nonselective orthosteric ligands caused peripheral adverse effects, was the M1 and M4 agonist, xanomeline. Although now recognized as a dual orthosteric and allosteric M1 agonist, xanomeline exhibited similar adverse effects associated with its orthosteric off-target activations of M2 and M3 subtypes, such as nausea, vomiting, excessive salivation, gastrointestinal disturbances, and bradycardia (Burger et al., 2023; Noetzel et al., 2009).

An emerging agent known as Kar-XT has combined xanomeline with tropium chloride, a peripherally restricted muscarinic antagonist that does not cross the BBB. This design preserves xanomeline's central M1/M4 activity while significantly reducing activity in the PNS. In a phase 1

healthy volunteer study (NCT02831231), the presence of trospium chloride reduced composite cholinergic adverse events by 46% compared to xanomeline alone. Adverse effects such as nausea, vomiting, diarrhea, sweating, and salivary hypersecretion were also reduced by at least 29% (Breier et al., 2023). Additionally, in prior phase 2 schizophrenia trials, KAR-XT demonstrated good efficacy and tolerability (Correll et al., 2022). While these clinical trials were conducted for schizophrenia, they target the same mechanism—muscarinic activation in the CNS protected from peripheral side effects—holding relevance for Alzheimer’s disease, where similar challenges of peripheral toxicity have long limited the development of orthosteric mAChR agonists.

Beyond Kar-XT, recent research using high-resolution crystallography and cryo-electron microscopy has discovered pocket geometries and loop conformations that are unique to each mAChR subtype, allowing for the design of ligands with improved bias (Thal et al., 2016). For instance, VU0357017 and AC-260584 are M1-preferring agonists that have shown high efficacy while having little to no activation of the M2-M5 subtypes (S. R. Bradley et al., 2010; Digby et al., 2012). AC-260584 exhibited potent functional activity (pEC<sub>50</sub> of 7.7 ± 0.1) and high efficacy with reduced activity at M2-M5. VU0357017 activated calcium signaling and ERK1/2 phosphorylation—pathways downstream of G-protein-coupled activation—while showing minimal β-arrestin recruitment, indicating a bias toward cognitive-relevant signaling cascades. While orthosteric agents offer a direct mechanism of action, even highly selective compounds can have adverse effects due to powerful signaling even at low levels of off-target activation. This risk has driven interest in allosteric modulators and bitopic ligands, which offer refined, tissue-specific modulation with potentially wider therapeutic scopes.

## **Allosteric Modulators**

Allosteric modulators bind to sites on the muscarinic receptor other than the orthosteric site. Unlike orthosteric ligands, which compete directly with endogenous ACh, allosteric modulators change receptor activity by influencing the receptor’s reaction to naturally released ACh. Positive Allosteric Modulators (PAMs) do this by increasing the response to acetylcholine, while Negative Allosteric Modulators (NAMs) regulate the response. Subtype selectivity is more achievable since allosteric sites are less conserved and more structurally unique among mAChR subtypes. Furthermore, endogenous ACh is released in specific locations and at precise times to regulate physiological functions. Allosteric modulation only occurs where endogenous ACh is present, so the risk of unwanted targeting in the PNS is reduced.

The most studied allosteric modulator has been M1-selective PAMs. The prototype M1 PAM, benzyl quinolone carboxylic acid (BQCA), has been developed as a preclinical research tool. It substantially promoted downstream signaling in medial prefrontal cortex (mPFC) pyramidal neurons, with effects absent in M1 knockout (KO) mice (Shirey et al., 2009). Most notably, BQCA restored discrimination reversal learning, often lost in AD patients, in a transgenic AD mouse model and regulated non-amyloidogenic APP processing in vitro. This suggested that

they could possess symptomatic and disease-modifying potential. Additionally, in prion-diseased mice—an in vivo model with cholinergic dysfunction similar to AD—daily BQCA doses restored cognitive function and significantly increased survival, with minimal side effects compared to treatment with orthosteric agonists or acetylcholinesterase inhibitors (S. J. Bradley et al., 2017). Despite this, BQCA displayed poor pharmacokinetic properties. More specifically, although it reached peak brain concentrations around 1.5 hours post-administration, systemic and tissue distribution favored organs like the kidney and lung over the brain, limiting its translational potential (Dwomoh et al., 2022). Building on BQCA's model, MK-7622 became the first human-tested M1 PAM for AD. The phase II trial evaluated it as an adjunct to standard AChEI therapy but was terminated early due to it having no significant cognitive or functional benefit compared to placebo. Importantly, cholinergic-like adverse events such as diarrhea and nausea were more frequent than in control groups (Voss et al., 2018). Displaying similar reactions on rodents, PF-06767832 and PF-06764427 helped identify intrinsic agonism or activation of the receptor as a cause for these effects (Davoren et al., 2016; Moran et al., 2018). As a result, M1-PAMs like VU319, which minimize agonist activity, have shown a lack of adverse effects and are being further researched (Poslunsey et al., 2024).

Given that M4 receptors act as inhibitory autoreceptors on cholinergic interneurons and as modulators of dopaminergic signaling, recent research has also been studying how they could potentially be targeted using PAMs to establish an excitatory–inhibitory balance in the networks affected by AD. Due to their recent rise to attention, most data on M4 PAMs for Alzheimer's treatment is preclinical and builds off of research on Schizophrenia, which involves some common pathways. A compound that has shown promise in this class of treatments is VU0152100, which is being developed at Vanderbilt University. It reversed amphetamine-induced hyperlocomotion in rodent models but not M4 knockout (KO) mice—demonstrating high selectivity—, lacked intrinsic agonist activity, and did not produce the adverse effects associated with non-selective agonists (Byun et al., 2014). Although initially researched for schizophrenia, the mechanism of reducing excessive excitability through M4 signaling is relevant to Alzheimer's pathology, where similar dysregulation occurs. Among the next generation of M4 PAMs are VU0467154, CVL-231, and NMRA-861, which have shown better potency as well as many of the same effects seen in VU0152100, showing progress and potential in this type of treatment (Berezovskaia et al., 2025; Butler et al., 2024). Additionally, in 2018, it was revealed that Xanomeline could bind to the allosteric site on M4, allowing for further understanding of the mechanisms at work behind Kar-Xt, and the potential for Xanomeline to be used as a PAM. (Burger et al., 2023)

Although less popular, negative allosteric modulators have also been explored for their roles in AD. Their purpose, unlike positive allosteric modulators, is to reduce the potency of endogenous ACh. Therefore, they target areas where mAChRs overactivation could be problematic, like the M2 receptor, which limits ACh in the cortex and hippocampus (Zuchner et al., 2005). It has been suggested that selective M2 antagonism can increase synaptic ACh and potentially improve cognition in AD by complementing AChEIs. The M2-selective NAM, Gallamine— a neuromuscular blocker— demonstrated negative cooperativity at muscarinic receptors, guiding research into M2 NAMs (Mash et al., 1985). Since then, there has been some research in

developing more subtype-selective and CNS-suitable M2-selective NAM candidates for AD. Despite some early compounds having demonstrated potential, factors like the possibility for increased A $\beta$  formation, less distribution in the CNS, and the lack of a complete understanding of the M2 receptor and its mechanisms, focus has remained on the more promising M1 and M4 receptors. While allosteric modulators solve many of the issues regarding subtype selectivity that orthosteric ligands couldn't, intrinsic agonism and their efficacy in general have driven focus to another type of agent, bitopic ligands.


## **Bitopic Ligands**

Bitopic or dualsteric ligands are a more recent method of targeting GPCRs that often have greater affinity and selectivity than orthosteric ligands or allosteric modulators alone. Engineered to engage the orthosteric site and a more structurally unique allosteric site at the same time, they can activate only designated downstream pathways through a process known as biased signaling. This is useful for targeting mAChRs in the CNS, where peripheral side effects limit traditional agonists. Therefore, bitopic ligands can potentially deliver CNS-selective effects with fewer off-target actions.


Like PAMs, bitopic ligands are a promising drug class for targeting M1. One of the earliest M1 targeting bitopic ligands, TBPB(tert-butyl peroxybenzoate) was initially thought to achieve its selectivity only through allosteric binding. However, later studies showed that mutations in both the orthosteric and allosteric binding sites affected the binding and signaling of TBPB (Keov et al., 2014). In rodent models, TBPB enhanced cortical acetylcholine release, improved performance in object recognition tasks, and promoted non-amyloidogenic APP processing through increased amyloid-secretase activity while still not inducing gastrointestinal side effects (Jones et al., 2008). Due to pharmacokinetic limitations, including poor solubility and modest brain penetration, development of TBPB ended; nevertheless, it serves as proof that M1-selective bitopic ligands could modulate pathways significant to Alzheimer's. 77-LH-28-1, another M1-directed bitopic ligand, improved spatial and working memory, in part by modulating calcium homeostasis and synaptic plasticity in hippocampal neurons without peripheral adverse effects (Langmead et al., 2008). Bitopic ligands targeting M4 for AD are less developed than those for M1, and none have yet emerged as especially promising.

## Binding Structures of Different Agent Types


1)



2)



3)







**Figure 4:** UCSF Chimera structure visualizations of class A GPCRs(green) bound to orthosteric, allosteric, and bitopic ligands(red) found on the RCSB Protein Data Bank. A2A and beta2 adrenergic receptors were used due to limited access to PDBs of mAChR binding.

4A: The A2A adrenergic receptor bound to an orthosteric ligand (PDB ID: 4EIY)

4B: The A2A adrenergic receptor bound to an allosteric ligand (PDB ID: 3VG9)

4C: The beta2 adrenergic receptor bound to an bitopic ligand (PDB ID: 8W1V)

*(Bitopic Ligands Support the Presence of a Metastable Binding Site at the B2 Adrenergic Receptor | Journal of Medicinal Chemistry, n.d.; G-Protein-Coupled Receptor Inactivation by an Allosteric Inverse-Agonist Antibody | Nature, n.d.; Structural Basis for Allosteric Regulation of GPCRs by Sodium Ions | Science, n.d.; Ef et al., n.d.-a, n.d.-b)*



### **Figure 5: Chemical Structures of Prominent mAChR Ligands Investigated in AD:**

Primary aromatic scaffolds (blue) provide the conserved features necessary for receptor interaction, linkers (red) influence binding orientation within the orthosteric pocket, hydrophobic or bulky substituents (green) improve bitopic interactions and selectivity. The structures were obtained from PubChem and highlighted using ChemDraw.

5A: Xanomeline, a Orthosteric Agonist with Secondary Allosteric Engagement

5B: MK-7622, a Selective M1-PAM

5C: VU0467319, a Bitopic PAM

5D: CVL-231, a Recent Allosteric PAM

(PubChem, n.d.-d, n.d.-c, n.d.-a, n.d.-b)

### **Limitations of Agents In Current Trials and Future Directions**

In addition to struggles relating to tolerability, pharmacokinetics, and target selectivity, recent trials of muscarinic-based therapies for AD are encountering a new set of challenges. A persistent translational gap remains, with many compounds showing efficacy in preclinical animal models [Table 2] but limited or inconsistent benefits in humans. This is frequently associated with species differences in receptor mechanisms and disease progression. For instance, although mouse models are able to emulate the basic aspects of cholinergic dysfunction such as ACh release and amyloid pathology, they lack the degeneration found in the basal forebrain. This is often due to the inability of mouse models to exhibit the same characteristics found in aging humans, who suffer the most from AD (R & T, 2011). Additionally, data collected from mouse models may not be able to indicate the effects of treatments on the more complex cognitive functions of humans like language and executive function. Furthermore, the lack of known biomarkers for target engagement makes it difficult to determine whether disappointing outcomes reflect insufficient receptor modulation or true therapeutic inefficacy.

Safety concerns also continue to emerge. Although recently developed ligands in clinical trials [Table 1] have shown improved receptor selectivity they have not fully eliminated the adverse effects associated with PNS activation. Both the selective M1 agonist HTL0018318 and M4 PAM CVL-231 were associated with transient increases in blood pressure, posing a risk for elderly patients with cardiovascular comorbidities (Jh et al., 2022; Nathan et al., 2022). Additionally, broader concerns about recruitment strategies in trials failing to take into account all groups of individuals, as well as unethical practices, remain. Recently, a prominent paper on amyloid pathology in AD published in 2006 by a team led by Sylvain Lesné was found to have manipulated images and data, misleading the direction of Alzheimer's research for over 15 years.

Given the aforementioned obstacles, muscarinic research for AD treatments is moving toward precision medicine approaches that find therapeutic strategies depending on patient-specific biology. The development of Positron Emission Tomography (PET) tracers for muscarinic receptors, such as [<sup>11</sup>C]VC-002 for M1, provides a tool to measure receptor availability in vivo (Haense et al., 2012). This technology could allow for the grouping of patients based on receptor density and track target engagement during clinical trials. As a result, researchers can

confirm with more confidence the efficacy of a drug. Genetic insights are also guiding muscarinic research. For instance, individuals carrying the APOE4 gene showed significant vulnerability to cholinergic dysfunction (Hill et al., 2007). Hence, they may particularly benefit from muscarinic-targeted therapies early on. Additionally, mutations and polymorphisms in the CHRM1 and CHRM4 genes have been linked to altered receptor expression or signaling, influencing the individual responses of drugs being tested (Dean & Scarr, 2021). The integration of genomic profiling into trials could reduce discrepancies like these. Combination therapy could also be helpful, given the complex nature of AD. More specifically, M4 PAMs could enhance cognition while anti-amyloid drugs like Aducanumab, Lecanemab, and Donanemab slow progression. Similarly, given that Memantine is taken in conjunction with AChEIs, pairing them with mAChR targeting therapeutics could improve balance in the glutamatergic system.

In the future, large molecule treatments may complement or even surpass small molecules in muscarinic receptor drug discovery. Allosteric antibodies, while not yet applied clinically to muscarinic receptors, have already been developed for other GPCRs such as CCR5 and GLP-1R, for which they have stabilized receptor conformations (Cong et al., 2021; Olson & Jacobson, 2009). Applying this strategy to M1 or M4 could lead to the development of therapeutics with improved selectivity, pharmacokinetics and tissue specificity, potentially overcoming the adverse effects seen with small-molecule drugs. Nanobodies (VHHs) are another promising platform as they have already been used to stabilize muscarinic receptor conformations. For instance, nanobodies Nb9-8 and Nb10-6 locked the M2 receptor in active states (Kruse et al., 2013). With further engineering, nanobodies could be developed as therapeutic allosteric modulators of M1 or M4. Finally, chemogenetic approaches offer proof-of-concept for circuit-specific cholinergic modulation. Designer Receptors Exclusively Activated by Designer Drugs (DREADDs), which were originally engineered from muscarinic receptors, have demonstrated this. The hM3Dq receptor, derived from M3, produces excitatory Gq signaling, while hM4Di, engineered from M4, produces inhibitory Gi signaling (Smith et al., 2016). Although DREADDs are primarily research tools, they have shown how muscarinic receptor scaffolds can be modified to enable precise spatiotemporal control of signaling, showing the potential for future gene-therapy-based interventions in Alzheimer’s disease.

**Table 1: Prominent Muscarinic Agents Currently Under Research for Relevance to AD in Clinical Trials**

Agent	Class / Mechanism	Development Stage & Context	Relevance to AD
KarXT (xanomeline + trospium chloride)	Dual Orthosteric and Allosteric M1/M4 agonist + peripheral orthosteric antagonist	Phase 3 trials for psychosis and agitation in AD; FDA-approved for schizophrenia as of 2024	Central activation of M1/M4 while minimizing peripheral side effects; cognitive and behavioral relevance in AD.

		Invented at PureTech Health; advanced by Karuna Therapeutics; acquired by Bristol Myers Squibb (BMS)	
HTL0018318	Orthosteric M1-selective agonist	Phase 1b/2a adjunct to donepezil in mild–moderate AD  Developed by Sosei Heptares in collaboration with Allergan	Designed for enhanced cognitive benefit with improved receptor specificity.
HTL0016878	Orthosteric M4-selective agonist	Phase 1 in healthy volunteers  Originated at Heptares Therapeutics; Advanced by Neurocrine Biosciences	Targeting neurobehavioral symptoms via M4 pathways.
HTL9936	Orthosteric M1-selective agonist	Phase 1 in healthy (elderly inclusive) subjects  Developed by Heptares Therapeutics	High selectivity addressing cognitive impairment with safety focus.
VU319	M1-positive allosteric modulator (PAM) with minimal intrinsic agonism	Phase 1 completed; proof-of-concept planned  Developed within Vanderbilt University's Warren Center for Neuroscience Drug Discovery (WCNDD)	Reduced side effects compared to earlier PAMs like MK-7622.
CVL-231 (Emraclidine)	M4 PAM	Phase 1 multiple-dose; includes AD dementia cohort	Balances excitatory/inhibitory signaling relevant in AD.

		Developed by Cerevel Therapeutics; now under AbbVie	
NS-136	M4 PAM	Phase 1 for schizophrenia; preclinical AD development  Developed by NeuShen Therapeutics	Forward-looking candidate, bridging schizophrenia and AD.

**Table 2: Prominent Muscarinic Agents Currently Under Research for Relevance to AD in Preclinical Trials**

Compound 24	M4 PAM (tool compound)	Preclinical in rodents and rhesus monkeys  N/A (research tool)	Demonstrated procognitive and antipsychotic-like effects in primate models.
VU0467154	M4 PAM	Preclinical (rodent)  Originated at Vanderbilt University	Enhanced learning and memory, strengthening case for cognitive benefit.
VU0152100 / VU0152099 / LY2033298	M4-selective PAMs (tool compounds)	Preclinical (rodents)  N/A	Validated M4 as an attractive target for cognition and behavior modulation.
AF267B (also AF102B / AF150(S))	Orthosteric M1-selective agonist	Preclinical (various animal models)  N/A	Reduced A $\beta$ and tau pathology; rescued memory in transgenic AD mice.
AF710B	M1-selective agonist with $\sigma$ 1 activity (bitopic-like)	Preclinical (rodents)  N/A	Improved cognition and APP processing with high selectivity.
GSK1034702	Bitopic M1 agonist	Clinical testing in humans but problematic due to side effects (not discontinued yet)	Demonstrated bitopic engagement and cognitive effects, but poor safety highlights need for refined ligands.

		Developed by GlaxoSmithKline (GSK)	
--	--	------------------------------------------	--

## Conclusion

Research in Alzheimer's pathogenesis, specifically with regards to the mAChRs, has certainly undergone significant progress since the proposition of the cholinergic hypothesis in 1976. Moving beyond the initial setbacks of earlier orthosteric agonists, more refined approaches with subtype selectivity, allosteric modulation, and biased signaling have been discovered. The M1 and M4 subtypes remain the most prominent targets due to their dense expression in memory-related and dopaminergic circuits and their direct influence on amyloid and tau pathology. Understanding of the overlap between these systems and other neurological diseases has contributed to the development of treatments like Kar-XT which were successful in trials for these related disorders, like SZ, and have since shown promise in AD trials. Nevertheless, much remains unknown and preclinical. Translation from animal models to humans continues to be inconsistent, trial design is slowed by the lack of biomarkers that are distinguishable, and the heterogeneity of Alzheimer's demands more precision in clinical trials.

Muscarinic-targeted therapies with genetic profiling, PET tracers, and potentially combined strategies with anti-amyloid or anti-tau drugs could help overcome these challenges. Furthermore, beyond small molecules, new techniques such as allosteric antibodies, nanobodies, and receptor-based gene therapies are beginning to gain attention. Together, these developments have improved understanding as well as served as a guide for the potential for innovation in treatments targeting mAChRs.

## Acknowledgements

I would like to thank [REDACTED] for their guidance and feedback throughout the research process.

## Author

[REDACTED] Having done research on whether ATP synthase could lead to increased energy production, epilepsy, and developments in muscarinic receptor targeting treatments for Alzheimer's, [REDACTED] has a deep interest and passion for the life sciences.

## References

Allen Institute for Science. (n.d.). *Allen Brain Atlas* [Database]. Allen Brain Map.

<https://human.brain-map.org>

Beach, T. G., Walker, D. G., Potter, P. E., Sue, L. I., & Fisher, A. (2001). Reduction of cerebrospinal fluid amyloid  $\beta$  after systemic administration of M1 muscarinic agonists.

*Brain Research*, 905(1), 220–223. [https://doi.org/10.1016/S0006-8993\(01\)02484-2](https://doi.org/10.1016/S0006-8993(01)02484-2)

Berezovskaia, A., Lindsley, C., Fink-Jensen, A., & Wörtwein, G. (2025). M4 Positive Allosteric Modulator VU0467154 Impacts Amphetamine Sensitization and Spontaneous Locomotion in Male Mice. *ACS Chemical Neuroscience*, 16(5), 868–879.

<https://doi.org/10.1021/acscemneuro.4c00795>

*BioRender*. (n.d.). Retrieved August 5, 2025, from

<https://app.biorender.com/illustrations/6891dfba4e0509a1dbb919db>

*Bitopic Ligands Support the Presence of a Metastable Binding Site at the  $\beta$ 2 Adrenergic*

*Receptor* | *Journal of Medicinal Chemistry*. (n.d.). Retrieved October 5, 2025, from

<https://pubs.acs.org/doi/10.1021/acs.jmedchem.4c00578>

Bradley, S. J., Bourgoignon, J.-M., Sanger, H. E., Verity, N., Mogg, A. J., White, D. J., Butcher, A. J., Moreno, J. A., Molloy, C., Macedo-Hatch, T., Edwards, J. M., Wess, J., Pawlak, R., Read, D. J., Sexton, P. M., Broad, L. M., Steinert, J. R., Mallucci, G. R., Christopoulos, A., ... Tobin, A. B. (2017). M1 muscarinic allosteric modulators slow prion neurodegeneration and restore memory loss. *The Journal of Clinical Investigation*, 127(2), 487–499. <https://doi.org/10.1172/JCI87526>

Bradley, S. R., Lamah, J., Ohrmund, L., Son, T., Bajpai, A., Nguyen, D., Friberg, M., Burstein, E. S., Spalding, T. A., Ott, T. R., Schiffer, H. H., Tabatabaei, A., McFarland, K., Davis, R. E., & Bonhaus, D. W. (2010). AC-260584, an orally bioavailable M(1) muscarinic receptor allosteric agonist, improves cognitive performance in an animal model.

*Neuropharmacology*, 58(2), 365–373. <https://doi.org/10.1016/j.neuropharm.2009.10.003>

- Breier, A., Brannan, S. K., Paul, S. M., & Miller, A. C. (2023). Evidence of trospium's ability to mitigate cholinergic adverse events related to xanomeline: Phase 1 study results. *Psychopharmacology*, *240*(5), 1191–1198. <https://doi.org/10.1007/s00213-023-06362-2>
- Burger, W. A. C., Pham, V., Vuckovic, Z., Powers, A. S., Mobbs, J. I., Laloudakis, Y., Glukhova, A., Wootten, D., Tobin, A. B., Sexton, P. M., Paul, S. M., Felder, C. C., Danev, R., Dror, R. O., Christopoulos, A., Valant, C., & Thal, D. M. (2023). Xanomeline displays concomitant orthosteric and allosteric binding modes at the M4 mAChR. *Nature Communications*, *14*(1), 5440. <https://doi.org/10.1038/s41467-023-41199-5>
- Butler, C. R., Popiolek, M., McAllister, L. A., LaChapelle, E. A., Kramer, M., Beck, E. M., Mente, S., Brodney, M. A., Brown, M., Gilbert, A., Helal, C., Ogilvie, K., Starr, J., Uccello, D., Grimwood, S., Edgerton, J., Garst-Orozco, J., Kozak, R., Lotarski, S., ... Steyn, S. J. (2024). Design and Synthesis of Clinical Candidate PF-06852231 (CVL-231): A Brain Penetrant, Selective, Positive Allosteric Modulator of the M4 Muscarinic Acetylcholine Receptor. *Journal of Medicinal Chemistry*, *67*(13), 10831–10847. <https://doi.org/10.1021/acs.jmedchem.4c00293>
- Byun, N. E., Grannan, M., Bubser, M., Barry, R. L., Thompson, A., Rosanelli, J., Gowrishankar, R., Kelm, N. D., Damon, S., Bridges, T. M., Melancon, B. J., Tarr, J. C., Brogan, J. T., Avison, M. J., Deutch, A. Y., Wess, J., Wood, M. R., Lindsley, C. W., Gore, J. C., ... Jones, C. K. (2014). Antipsychotic drug-like effects of the selective M4 muscarinic acetylcholine receptor positive allosteric modulator VU0152100. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *39*(7), 1578–1593. <https://doi.org/10.1038/npp.2014.2>
- Caccamo, A., Oddo, S., Billings, L. M., Green, K. N., Martinez-Coria, H., Fisher, A., & LaFerla, F. M. (2006). M1 Receptors Play a Central Role in Modulating AD-like Pathology in Transgenic Mice. *Neuron*, *49*(5), 671–682. <https://doi.org/10.1016/j.neuron.2006.01.020>

- Choi, D. W. (1988). Glutamate neurotoxicity and diseases of the nervous system. *Neuron*, 1(8), 623–634. [https://doi.org/10.1016/0896-6273\(88\)90162-6](https://doi.org/10.1016/0896-6273(88)90162-6)
- Cisse, M., Braun, U., Leitges, M., Fisher, A., Pages, G., Checler, F., & Vincent, B. (2011). ERK1-independent  $\alpha$ -secretase cut of  $\beta$ -amyloid precursor protein via M1 muscarinic receptors and PKC $\alpha/\epsilon$ . *Molecular and Cellular Neuroscience*, 47(3), 223–232. <https://doi.org/10.1016/j.mcn.2011.04.008>
- Cong, Z., Chen, L.-N., Ma, H., Zhou, Q., Zou, X., Ye, C., Dai, A., Liu, Q., Huang, W., Sun, X., Wang, X., Xu, P., Zhao, L., Xia, T., Zhong, W., Yang, D., Eric Xu, H., Zhang, Y., & Wang, M.-W. (2021). Molecular insights into ago-allosteric modulation of the human glucagon-like peptide-1 receptor. *Nature Communications*, 12, 3763. <https://doi.org/10.1038/s41467-021-24058-z>
- Conn, P. J., Jones, C. K., & Lindsley, C. W. (2009). Subtype-selective allosteric modulators of muscarinic receptors for the treatment of CNS disorders. *Trends in Pharmacological Sciences*, 30(3), 148–155. <https://doi.org/10.1016/j.tips.2008.12.002>
- Correll, C. U., Angelov, A. S., Miller, A. C., Weiden, P. J., & Brannan, S. K. (2022). Safety and tolerability of KarXT (xanomeline–trospium) in a phase 2, randomized, double-blind, placebo-controlled study in patients with schizophrenia. *Schizophrenia*, 8(1), 109. <https://doi.org/10.1038/s41537-022-00320-1>
- Davies, P., & Maloney, A. J. F. (1976). SELECTIVE LOSS OF CENTRAL CHOLINERGIC NEURONS IN ALZHEIMER'S DISEASE. *The Lancet*, 308(8000), 1403. [https://doi.org/10.1016/S0140-6736\(76\)91936-X](https://doi.org/10.1016/S0140-6736(76)91936-X)
- Davis, A. A., Fritz, J. J., Wess, J., Lah, J. J., & Levey, A. I. (2010). Deletion of M1 Muscarinic Acetylcholine Receptors Increases Amyloid Pathology In Vitro and In Vivo. *Journal of Neuroscience*, 30(12), 4190–4196. <https://doi.org/10.1523/JNEUROSCI.6393-09.2010>
- Davoren, J. E., Lee, C.-W., Garnsey, M., Brodney, M. A., Cordes, J., Dlugolenski, K., Edgerton, J. R., Harris, A. R., Helal, C. J., Jenkinson, S., Kauffman, G. W., Kenakin, T. P., Lazzaro,

- J. T., Lotarski, S. M., Mao, Y., Nason, D. M., Northcott, C., Nottebaum, L., O'Neil, S. V., ... Grimwood, S. (2016). Discovery of the Potent and Selective M1 PAM-Agonist N-[(3R,4S)-3-Hydroxytetrahydro-2H-pyran-4-yl]-5-methyl-4-[4-(1,3-thiazol-4-yl)benzyl]pyridine-2-carboxamide (PF-06767832): Evaluation of Efficacy and Cholinergic Side Effects. *Journal of Medicinal Chemistry*, 59(13), 6313–6328.  
<https://doi.org/10.1021/acs.jmedchem.6b00544>
- Dean, B., & Scarr, E. (2021). Changes in cortical gene expression in the muscarinic M1 receptor knockout mouse: Potential relevance to schizophrenia, Alzheimer's disease and cognition. *Npj Schizophrenia*, 7(1), 44. <https://doi.org/10.1038/s41537-021-00174-z>
- Dementia*. (n.d.). Retrieved July 29, 2025, from <https://www.who.int/news-room/fact-sheets/detail/dementia>
- Digby, G. J., Noetzel, M. J., Bubser, M., Utley, T. J., Walker, A. G., Byun, N. E., Lebois, E. P., Xiang, Z., Sheffler, D. J., Cho, H. P., Davis, A. A., Nemirovsky, N. E., Mennenga, S. E., Camp, B. W., Bimonte-Nelson, H. A., Bode, J., Italiano, K., Morrison, R., Daniels, J. S., ... Conn, P. J. (2012). Novel Allosteric Agonists of M1 Muscarinic Acetylcholine Receptors Induce Brain Region-Specific Responses That Correspond with Behavioral Effects in Animal Models. *The Journal of Neuroscience*, 32(25), 8532–8544.  
<https://doi.org/10.1523/JNEUROSCI.0337-12.2012>
- Dwomoh, L., Tejada, G. S., & Tobin, A. B. (2022). Targeting the M1 muscarinic acetylcholine receptor in Alzheimer's disease. *Neuronal Signaling*, 6(1), NS20210004.  
<https://doi.org/10.1042/NS20210004>
- Ef, P., Td, G., Cc, H., Gs, C., Dm, G., Ec, M., & Te, F. (n.d.-a). UCSF Chimera—A visualization system for exploratory research and analysis. *PubMed*. Retrieved October 5, 2025, from <https://pubmed.ncbi.nlm.nih.gov/15264254/>

Ef, P., Td, G., Cc, H., Gs, C., Dm, G., Ec, M., & Te, F. (n.d.-b). UCSF Chimera—A visualization system for exploratory research and analysis. *PubMed*. Retrieved October 6, 2025, from <https://pubmed.ncbi.nlm.nih.gov/15264254/>

Fisher, A. (2008). Cholinergic treatments with emphasis on m1 muscarinic agonists as potential disease-modifying agents for Alzheimer's disease. *Neurotherapeutics: The Journal of the American Society for Experimental NeuroTherapeutics*, 5(3), 433–442.  
<https://doi.org/10.1016/j.nurt.2008.05.002>

Flynn, D. D., Weinstein, D. A., & Mash, D. C. (1991). Loss of high-affinity agonist binding to M1 muscarinic receptors in Alzheimer's disease: Implications for the failure of cholinergic replacement therapies. *Annals of Neurology*, 29(3), 256–262.  
<https://doi.org/10.1002/ana.410290305>

Forlenza, O. V., Spink, J. M., Dayanandan, R., Anderton, B. H., Olesen, O. F., & Lovestone, S. (2000). Muscarinic agonists reduce tau phosphorylation in non-neuronal cells via GSK-3 $\beta$  inhibition and in neurons. *Journal of Neural Transmission*, 107(10), 1201–1212.  
<https://doi.org/10.1007/s007020070034>

Foster, D. J., Choi, D. L., Conn, P. J., & Rook, J. M. (2014). Activation of M1 and M4 muscarinic receptors as potential treatments for Alzheimer's disease and schizophrenia. *Neuropsychiatric Disease and Treatment*, 10, 183–191.  
<https://doi.org/10.2147/NDT.S55104>

Frost, B., Jacks, R. L., & Diamond, M. I. (2009). Propagation of tau misfolding from the outside to the inside of a cell. *The Journal of Biological Chemistry*, 284(19), 12845–12852.  
<https://doi.org/10.1074/jbc.M808759200>

*G-protein-coupled receptor inactivation by an allosteric inverse-agonist antibody* | *Nature*. (n.d.). Retrieved October 5, 2025, from <https://www.nature.com/articles/nature10750>

Haense, C., Kalbe, E., Herholz, K., Hohmann, C., Neumaier, B., Kraus, R., & Heiss, W.-D. (2012). Cholinergic system function and cognition in mild cognitive impairment.

*Neurobiology of Aging*, 33(5), 867–877.

<https://doi.org/10.1016/j.neurobiolaging.2010.08.015>

Hampel, H., Elhage, A., Cho, M., Apostolova, L. G., Nicoll, J. A. R., & Atri, A. (2023).

Amyloid-related imaging abnormalities (ARIA): Radiological, biological and clinical characteristics. *Brain*, 146(11), 4414–4424. <https://doi.org/10.1093/brain/awad188>

Hardy, J. A., & Higgins, G. A. (1992). Alzheimer's Disease: The Amyloid Cascade Hypothesis.

*Science*, 256(5054), 184–185. <https://doi.org/10.1126/science.1566067>

Hill, J. M., Bhattacharjee, P. S., & Neumann, D. M. (2007). Apolipoprotein E alleles can

contribute to the pathogenesis of numerous clinical conditions including HSV-1 corneal disease. *Experimental Eye Research*, 84(5), 801–811.

<https://doi.org/10.1016/j.exer.2006.08.001>

Hogan, D. B. (2014). Long-Term Efficacy and Toxicity of Cholinesterase Inhibitors in the

Treatment of Alzheimer Disease. *Canadian Journal of Psychiatry. Revue Canadienne de Psychiatrie*, 59(12), 618–623. <https://doi.org/10.1177/070674371405901202>

Hoskin, J. L., Al-Hasan, Y., & Sabbagh, M. N. (2018). Nicotinic Acetylcholine Receptor Agonists

for the Treatment of Alzheimer's Dementia: An Update. *Nicotine & Tobacco Research*, 21(3), 370–376. <https://doi.org/10.1093/ntr/nty116>

Jh, K., Jm, K., Cu, C., Dp, W., M, L., S, D., S, P., I, C., P, I., L, F., S, V., P, P., R, S., & J, R.

(2022). Emraclidine, a novel positive allosteric modulator of cholinergic M4 receptors, for the treatment of schizophrenia: A two-part, randomised, double-blind, placebo-controlled, phase 1b trial. *PubMed*.

[https://pubmed.ncbi.nlm.nih.gov/36528376/?utm\\_source=chatgpt.com](https://pubmed.ncbi.nlm.nih.gov/36528376/?utm_source=chatgpt.com)

Jones, C. K., Brady, A. E., Davis, A. A., Xiang, Z., Bubser, M., Tantawy, M. N., Kane, A. S.,

Bridges, T. M., Kennedy, J. P., Bradley, S. R., Peterson, T. E., Ansari, M. S., Baldwin, R. M., Kessler, R. M., Deutch, A. Y., Lah, J. J., Levey, A. I., Lindsley, C. W., & Conn, P. J.

(2008). Novel Selective Allosteric Activator of the M1 Muscarinic Acetylcholine Receptor

Regulates Amyloid Processing and Produces Antipsychotic-Like Activity in Rats. *The Journal of Neuroscience*, 28(41), 10422–10433.

<https://doi.org/10.1523/JNEUROSCI.1850-08.2008>

Keov, P., López, L., Devine, S. M., Valant, C., Lane, J. R., Scammells, P. J., Sexton, P. M., & Christopoulos, A. (2014). Molecular Mechanisms of Bitopic Ligand Engagement with the M1 Muscarinic Acetylcholine Receptor. *The Journal of Biological Chemistry*, 289(34), 23817–23837. <https://doi.org/10.1074/jbc.M114.582874>

Kruse, A. C., Hu, J., Pan, A. C., Arlow, D. H., Rosenbaum, D. M., Rosemond, E., Green, H. F., Liu, T., Chae, P. S., Dror, R. O., Shaw, D. E., Weis, W. I., Wess, J., & Kobilka, B. K. (2012). Structure and dynamics of the M3 muscarinic acetylcholine receptor. *Nature*, 482(7386), 552–556. <https://doi.org/10.1038/nature10867>

Kruse, A. C., Ring, A. M., Manglik, A., Hu, J., Hu, K., Eitel, K., Hübner, H., Pardon, E., Valant, C., Sexton, P. M., Christopoulos, A., Felder, C. C., Gmeiner, P., Steyaert, J., Weis, W. I., Garcia, K. C., Wess, J., & Kobilka, B. K. (2013). Activation and allosteric modulation of a muscarinic acetylcholine receptor. *Nature*, 504(7478), 101–106.

<https://doi.org/10.1038/nature12735>

Lange, K. W., Javoy-Agid, F., Agid, Y., Jenner, P., & Marsden, C. D. (1992). Brain muscarinic cholinergic receptors in Huntington's disease. *Journal of Neurology*, 239(2), 103–104.

<https://doi.org/10.1007/BF00862983>

Langmead, C. J., Austin, N. E., Branch, C. L., Brown, J. T., Buchanan, K. A., Davies, C. H., Forbes, I. T., Fry, V. a. H., Hagan, J. J., Herdon, H. J., Jones, G. A., Jeggo, R., Kew, J. N. C., Mazzali, A., Melarange, R., Patel, N., Pardoe, J., Randall, A. D., Roberts, C., ... Watson, J. (2008). Characterization of a CNS penetrant, selective M1 muscarinic receptor agonist, 77-LH-28-1. *British Journal of Pharmacology*, 154(5), 1104–1115.

<https://doi.org/10.1038/bjp.2008.152>

- Levey, A., Kitt, C., Simonds, W., Price, D., & Brann, M. (1991). Identification and localization of muscarinic acetylcholine receptor proteins in brain with subtype-specific antibodies. *The Journal of Neuroscience*, *11*(10), 3218–3226.  
<https://doi.org/10.1523/JNEUROSCI.11-10-03218.1991>
- Lin, L., Georgievska, B., Mattsson, A., & Isacson, O. (1999). Cognitive changes and modified processing of amyloid precursor protein in the cortical and hippocampal system after cholinergic synapse loss and muscarinic receptor activation. *Proceedings of the National Academy of Sciences*, *96*(21), 12108–12113. <https://doi.org/10.1073/pnas.96.21.12108>
- Malenka, R. C., & Bear, M. F. (2004). LTP and LTD: An embarrassment of riches. *Neuron*, *44*(1), 5–21. <https://doi.org/10.1016/j.neuron.2004.09.012>
- Marino, M. J., Rouse, S. T., Levey, A. I., Potter, L. T., & Conn, P. J. (1998). Activation of the genetically defined m1 muscarinic receptor potentiates N-methyl-D-aspartate (NMDA) receptor currents in hippocampal pyramidal cells. *Proceedings of the National Academy of Sciences of the United States of America*, *95*(19), 11465–11470.  
<https://doi.org/10.1073/pnas.95.19.11465>
- Mash, D. C., Flynn, D. D., & Potter, L. T. (1985). Loss of M2 muscarine receptors in the cerebral cortex in Alzheimer's disease and experimental cholinergic denervation. *Science (New York, N.Y.)*, *228*(4703), 1115–1117. <https://doi.org/10.1126/science.3992249>
- Medications for Memory, Cognition & Dementia-Related Behaviors* | *alz.org*. (n.d.). Alzheimer's Association. Retrieved July 30, 2025, from  
<https://www.alz.org/alzheimers-dementia/treatments/medications-for-memory>
- Mitoh, Y., Ueda, H., Ichikawa, H., Fujita, M., Kobashi, M., & Matsuo, R. (2017). Effects of cevimeline on excitability of parasympathetic preganglionic neurons in the superior salivatory nucleus of rats. *Autonomic Neuroscience*, *206*, 1–7.  
<https://doi.org/10.1016/j.autneu.2017.05.010>

- Moran, S. P., Dickerson, J. W., Cho, H. P., Xiang, Z., Maksymetz, J., Remke, D. H., Lv, X., Doyle, C. A., Rajan, D. H., Niswender, C. M., Engers, D. W., Lindsley, C. W., Rook, J. M., & Conn, P. J. (2018). M1-positive allosteric modulators lacking agonist activity provide the optimal profile for enhancing cognition. *Neuropsychopharmacology*, *43*(8), 1763–1771. <https://doi.org/10.1038/s41386-018-0033-9>
- Nathan, P. J., Millais, S. B., Godwood, A., Dewit, O., Cross, D. M., Liptrot, J., Ruparelia, B., Jones, S. P., Bakker, G., Maruff, P. T., Light, G. A., Brown, A. J. H., Weir, M. P., Congreve, M., & Tasker, T. (2022). A phase 1b/2a multicenter study of the safety and preliminary pharmacodynamic effects of selective muscarinic M1 receptor agonist HTL0018318 in patients with mild-to-moderate Alzheimer's disease. *Alzheimer's & Dementia : Translational Research & Clinical Interventions*, *8*(1), e12273. <https://doi.org/10.1002/trc2.12273>
- Nielsen, B. E., & Ford, C. P. (2024). Reduced striatal M4-cholinergic signaling following dopamine loss contributes to parkinsonian and L-DOPA-induced dyskinetic behaviors. *Science Advances*, *10*(47), eadp6301. <https://doi.org/10.1126/sciadv.adp6301>
- Noetzel, M. J., Grant, M. K. O., & El-Fakahany, E. E. (2009). Immediate and Delayed Consequences of Xanomeline Wash-Resistant Binding at the M3 Muscarinic Receptor. *Neurochemical Research*, *34*(6), 1138–1149. <https://doi.org/10.1007/s11064-008-9886-3>
- Oleksak, P., Novotny, M., Patocka, J., Nepovimova, E., Hort, J., Pavlik, J., Klimova, B., Valis, M., & Kuca, K. (2021). Neuropharmacology of Cevimeline and Muscarinic Drugs—Focus on Cognition and Neurodegeneration. *International Journal of Molecular Sciences*, *22*(16), 8908. <https://doi.org/10.3390/ijms22168908>
- Olson, W. C., & Jacobson, J. M. (2009). CCR5 Monoclonal Antibodies for HIV-1 Therapy. *Current Opinion in HIV and AIDS*, *4*(2), 104–111. <https://doi.org/10.1097/COH.0b013e3283224015>

Pimlott, S. L., Piggott, M., Owens, J., Greally, E., Court, J. A., Jaros, E., Perry, R. H., Perry, E. K., & Wyper, D. (2004). Nicotinic Acetylcholine Receptor Distribution in Alzheimer's Disease, Dementia with Lewy Bodies, Parkinson's Disease, and Vascular Dementia: In Vitro Binding Study Using 5-[125I]-A-85380. *Neuropsychopharmacology*, 29(1), 108–116. <https://doi.org/10.1038/sj.npp.1300302>

Poslunsey, M. S., Wood, M. R., Han, C., Stauffer, S. R., Panarese, J. D., Melancon, B. J., Engers, J. L., Dickerson, J. W., Peng, W., Noetzel, M. J., Cho, H. P., Rodriguez, A. L., Hopkins, C. R., Morrison, R., Crouch, R. D., Bridges, T. M., Blobaum, A. L., Boutaud, O., Daniels, J. S., ... Lindsley, C. W. (2024). Discovery of VU0467319: An M1 Positive Allosteric Modulator Candidate That Advanced into Clinical Trials. *ACS Chemical Neuroscience*, 16(1), 95–107. <https://doi.org/10.1021/acschemneuro.4c00769>

PubChem. (n.d.-a).

6-[[2,6-difluoro-4-(2-methyl-2H-indazol-4-yl)phenyl]methyl]-6,7-dihydro-5H-Pyrrolo[3,4-b]pyridin-5-one. Retrieved October 5, 2025, from <https://pubchem.ncbi.nlm.nih.gov/compound/71555143>

PubChem. (n.d.-b). *Emraclidine*. Retrieved October 5, 2025, from <https://pubchem.ncbi.nlm.nih.gov/compound/140830653>

PubChem. (n.d.-c). *Mk-7622*. Retrieved October 6, 2025, from <https://pubchem.ncbi.nlm.nih.gov/compound/46207733>

PubChem. (n.d.-d). *Xanomeline*. Retrieved October 5, 2025, from <https://pubchem.ncbi.nlm.nih.gov/compound/60809>

R, S., & T, A. (2011). The cholinergic system in aging and neuronal degeneration. *PubMed*. <https://pubmed.ncbi.nlm.nih.gov/21145918/>

Ramos Bernardes da Silva Filho, S., Oliveira Barbosa, J. H., Rondinoni, C., dos Santos, A. C., Garrido Salmon, C. E., da Costa Lima, N. K., Ferriolli, E., & Moriguti, J. C. (2017).

Neuro-degeneration profile of Alzheimer's patients: A brain morphometry study.

*NeuroImage: Clinical*, 15, 15–24. <https://doi.org/10.1016/j.nicl.2017.04.001>

Ren, S., Singh, J., Gsteiger, S., Cogley, C., Reed, B., Abrams, K. R., Dawoud, D., Owen, R. K., Tappenden, P., Quinn, T. J., & Bujkiewicz, S. (2025). *Evaluating amyloid-beta as a surrogate endpoint in trials of anti-amyloid drugs in Alzheimer's disease: A Bayesian meta-analysis* (No. arXiv:2504.06807). arXiv. <https://doi.org/10.48550/arXiv.2504.06807>

Schuff, N., Woerner, N., Boreta, L., Kornfield, T., Shaw, L. M., Trojanowski, J. Q., Thompson, P. M., Jack, C. R., Weiner, M. W., & Alzheimer's Disease Neuroimaging Initiative. (2009). MRI of hippocampal volume loss in early Alzheimer's disease in relation to ApoE genotype and biomarkers. *Brain: A Journal of Neurology*, 132(Pt 4), 1067–1077. <https://doi.org/10.1093/brain/awp007>

Shirey, J. K., Brady, A. E., Jones, P. J., Davis, A. A., Bridges, T. M., Kennedy, J. P., Jadhav, S. B., Menon, U. N., Xiang, Z., Watson, M. L., Christian, E. P., Doherty, J. J., Quirk, M. C., Snyder, D. H., Lah, J. J., Levey, A. I., Nicolle, M. M., Lindsley, C. W., & Conn, P. J. (2009). A Selective Allosteric Potentiator of the M1 Muscarinic Acetylcholine Receptor Increases Activity of Medial Prefrontal Cortical Neurons and Restores Impairments in Reversal Learning. *The Journal of Neuroscience*, 29(45), 14271–14286. <https://doi.org/10.1523/JNEUROSCI.3930-09.2009>

Smiley, J. F., Levey, A. I., & Mesulam, M.-M. (1999). M2 Muscarinic receptor immunolocalization in cholinergic cells of the monkey basal forebrain and striatum. *Neuroscience*, 90(3), 803–814. [https://doi.org/10.1016/S0306-4522\(98\)00527-2](https://doi.org/10.1016/S0306-4522(98)00527-2)

Smith, K. S., Bucci, D. J., Luikart, B. W., & Mahler, S. V. (2016). DREADDs: Use and Application in Behavioral Neuroscience. *Behavioral Neuroscience*, 130(2), 137–155. <https://doi.org/10.1037/bne0000135>

*Structural Basis for Allosteric Regulation of GPCRs by Sodium Ions* | *Science*. (n.d.). Retrieved October 5, 2025, from <https://www.science.org/doi/10.1126/science.1219218>

- Thal, D. M., Sun, B., Feng, D., Nawaratne, V., Leach, K., Felder, C. C., Bures, M. G., Evans, D. A., Weis, W. I., Bachhawat, P., Kobilka, T. S., Sexton, P. M., Kobilka, B. K., & Christopoulos, A. (2016). Crystal structures of the M1 and M4 muscarinic acetylcholine receptors. *Nature*, *531*(7594), 335–340. <https://doi.org/10.1038/nature17188>
- Voss, T., Li, J., Cummings, J., Farlow, M., Assaid, C., Froman, S., Leibensperger, H., Snow-Adami, L., McMahon, K. B., Egan, M., & Michelson, D. (2018). Randomized, controlled, proof-of-concept trial of MK-7622 in Alzheimer's disease. *Alzheimer's & Dementia: Translational Research & Clinical Interventions*, *4*(1), 173–181. <https://doi.org/10.1016/j.trci.2018.03.004>
- Wess, J. (1996). Molecular biology of muscarinic acetylcholine receptors. *Critical Reviews in Neurobiology*, *10*(1), 69–99. <https://doi.org/10.1615/critrevneurobiol.v10.i1.40>
- Zhang, J., Zhang, Y., Wang, J., Xia, Y., Zhang, J., & Chen, L. (2024). Recent advances in Alzheimer's disease: Mechanisms, clinical trials and new drug development strategies. *Signal Transduction and Targeted Therapy*, *9*(1), 211. <https://doi.org/10.1038/s41392-024-01911-3>
- Zuchner, T., Schliebs, R., & Perez-Polo, J. R. (2005). Down-regulation of muscarinic acetylcholine receptor M2 adversely affects the expression of Alzheimer's disease-relevant genes and proteins. *Journal of Neurochemistry*, *95*(1), 20–32. <https://doi.org/10.1111/j.1471-4159.2005.03335.x>

#### Suggestions:

- Consider adding keywords to the abstract, per standard academic formatting.
  - I added keywords below my abstract.

#### Minor Formatting Suggestions:

- The first paragraph of the Introduction needs slight formatting cleanup (possibly a missing line break between lines 5–6).
  - This indent happened when I converted my citations from manual numbered parenthesis to APA to using Zotero. I'm not sure how to remove it.
- Consider simplifying the phrase “Clinical Dementia Rating – Sum of Boxes scores” in the Introduction.
  - Revised: For instance, Lecanemab showed a 27% slower decline in CDR-SB (Clinical Dementia Rating – Sum of Boxes) scores over 18 months
- Consider adding brief clarifying sentences that connect receptor-level mechanisms to patient-level outcomes (e.g., how M1 modulation might improve memory or reduce behavioral symptoms). These would help reinforce the paper’s translational relevance and strengthen the connection between scientific detail and therapeutic goals.
  - I connected it to cognitive function. However, much remains unknown and I don't want to write anything misleading.
- Table 1 formatting: The drug development table is dense and helpful, but formatting consistency could be improved. You should follow the specific APA guidelines. Consider:  
Aligning columns visually  
Using abbreviations more sparingly (e.g. define “KO” again if used after a long gap)
  - I specified that KO meant knockout.

Possibly moving preclinical compounds into a supplementary table to make the main one more digestible

- I split the table in two (one for clinical and one for preclinical).

#### Citation Formatting Suggestions:

- Ensure all citations are placed before punctuation marks, as required by APA.
- Ensure figures created in BioRender are compliant with their open-access license.
- Ensure Figure 2 is compliant with the Allen Institute’s citation guidelines
- Ensure Figure 4 is compliant with the RCSB Protein Data Bank citation policies
- Dementia (n.d.) reference from WHO should be cited correctly as a fact sheet. Note also that the publication year is available (2025). Same is true for “Medications for Memory, Cognition & Dementia-Related Behaviors” reference. Refer to these guidelines for both and update the citations according to APA guidelines: <https://apastyle.apa.org/style-grammar-guidelines/references/examples/webpage-website-references#4>
  - I fixed my citations to follow APA guidelines.

#### 4. Engagement with Literature

##### Suggestions:

- Add brief mention of how the review expands upon or contrasts with earlier reviews on similar topics.

- I believe I already did this by mentioning how research on treatments is shifting away from solely focusing on amyloid and tau pathology, and that new technologies and discoveries have come about since, bringing perspectives on the topic.

#### 5. Grammar & Language

##### Minor Suggestions:

- Standardize formatting for abbreviations.
- Correct minor typo on page 9, second paragraph, last sentence
  - I fixed these issues.

- 1) Revised Sentence: "It then analyzes the status of developments in different classes of treatments based on performance in trials, efficacy data, pathways targeted, and adverse effects."  
I made the sentence grammatically correct and rephrased it to make it more clear and shorter.
- 2) This indent happened when I converted my citations from manual numbered parenthesis to APA to using Zotero. I'm not sure how to remove it.
- 3) I changed "synthesize" to "analyze" and "paper" to "literature review"
- 4) Previous Sentence: "Then, analyze the pathways that have been researched as a result and the status of drug developments and therapeutics that have attempted to target them."  
Revised: "The paper will then examine the both pathways that have been researched as a result, and the status of drug developments and therapeutics that have attempted to target them."
- 5) I got rid of the comma.
- 6) I removed the detergents in UCSF Chimera and labeled the GPCRs and ligands.
- 7) I labeled the regions to clearly depict where the cell membranes would be
- 8) I put parenthesis indicating the colors of the ligand and GPCR.
- 9) I used UCSF Chimera and fixed the mistake.
- 10) I stated the PDB IDs
- 11) I added that CVL-231 exhibited similar adverse effects, but most of the compounds in clinical trials are still in phase 1, therefore the adverse effects aren't yet known or public.
- 12) I wrote a paragraph on how struggles with translation from mouse models are relevant to the development of ligands targeting mAChRs for AD.
- 13) I fixed grammar errors.

Dear Convergence Editorial Board,

In the writing of my manuscript, I made sure all AI use was ethical and very limited. I ran my use by both my TA and advisor multiple times to make sure it was proper and in line with IRIS's academic integrity standards. At no point did it write any part of my manuscript.

Artificial Intelligence was used solely as an enhanced search engine to define terms and find information. When I looked up questions, Google automatically gave an AI generated response as well as some cited studies or articles. The AI summaries often weren't of much use to me, and I relied more on the sources it cited for information. I would then check myself to find the content for my paper. When reading the articles, if I came across any technical phrases or terms that I was unfamiliar with, I would sometimes ask Chat-GPT to define them in more detail if I didn't fully understand what they meant after looking them up. Any paraphrasing I did was influenced only by feedback in the IRIS Writing Workshops and from the advisors themselves. I had been warned against using AI for this purpose and didn't. I did, however, use Grammarly to check for grammatical errors, and even with that, there were still some left. I also used Chat-GPT to validate that the information I had on agents wasn't outdated and to clear up any questions I had when I couldn't speak with a person. Nevertheless, I made sure to check the sources it cited for any inaccuracies in its responses. Finally, I used AI to find and navigate databases and tools that I was unfamiliar with, like UCSF Chimera X and the Allen Brain Institute. To clarify, I did not use artificial intelligence to analyze and describe the information I found on those platforms.

Once again, all analyses, interpretations, and suggestions in the paper were my own work. Any AI use was used solely to clarify facts and navigate platforms, and not for ideas. I value academic integrity as much as Convergence does and cited any information that did not come directly from me.

Regards,



**Academic Integrity Guidelines:** [GPT Policy](#), [Student & Parent Honor Code](#), [Mentor Honor Code](#)  
**Indigo Contact Information:** [hello@indigoresearch.org](mailto:hello@indigoresearch.org)  
[Indigo Research IRIS Webpage](#)

Dear Colleague,

I write in reference to [REDACTED] whom I taught through Indigo Research's 2025 IRIS Summer Program.


[REDACTED] performance in my course [REDACTED] gave me every reason to believe that his final paper is an original work of scholarship. After watching [REDACTED] develop his paper over the course of our six-week seminar, meeting with them individually, and reading his final paper, it is my opinion that this paper should be allowed to continue through the peer-review process.

[REDACTED] attended nearly every office hours session with me. During these sessions, he shared multiple drafts of his paper. He showed deep knowledge of his topic, both verbally and in writing. He was committed to improving his writing skills throughout the process. [REDACTED] often checked in with me to confirm that he used AI appropriately and within the IRIS guidelines, which included using AI to define terms, and nothing more. Further, when I ran his paper through the Grammarly AI checker, the writing showed only 3% likely AI use. To the best of my knowledge, the paper was written and edited by [REDACTED], and he spent a significant amount of time researching the literature and processing the information required to understand the topic and write the review.

While it is impossible to confirm with absolute certainty that no AI was involved in the writing of this paper, I write as an instructor in support of my student's academic integrity. I am not agreeing to take on any personal liability for this paper.

Sincerely,

[REDACTED], PhD



[REDACTED]  
IRIS Program 2025

Oct 20, 2025

Recommendation: **Accept**

The initial submission was a very detailed study of muscarinic receptors, their role in Alzheimer's disease and the strategies of treatment by targeting muscarinic receptors. Here, the author has submitted a revised version of the manuscript that is much more polished in terms of structure and writing. The author has also satisfactorily addressed all previous concerns, specifically Figure 4. In particular, the author has indicated the ligands, protein structure, intracellular and extracellular portions of the receptor. These will go a long way in helping the readers understand the concepts. The author has also included all PDB IDs that can be referred to in the databases. Furthermore, the author has taken the initiative of adding one more figure in this version: Figure 5, which is an illustration of the chemical structures of the various ligands, and indicated the important aspects in the ligands for their role in binding to muscarinic receptors. While it was not requested, this reviewer has found Figure 5 extremely helpful in illustrating the points in the paper and appreciated this addition.

Suggestion: The paper is detailed and informative and the author obviously knows the topic well. This reviewer would like to suggest better integration and more references to figures through out the main body of the text. There are 5 figures and 2 tables, all of which contain key information for the paper. However, there are not many references to the figures in the text. There were 2 references to Figure 1 in the introduction part and in "Connections to Other pathways and diseases", and references to the 2 tables, but this reviewer did not see other references to other figures. This reviewer would like to suggest that the author make more references to the other figures in the text and this will strengthen the paper.