


# Bartlett Surgery & Specialty Clinic

3225 HOSPITAL DRIVE, SUITE 101A JUNEAU, AK 99801 • 907-796-8700 • FAX 907-796-8710

<b>Before Surgery Instructions:</b>		
<b>2 days prior to surgery</b>	<b>Start drops two days prior to surgery in:  Right eye / left eye</b>	<b>Moxifloxacin/ Ofloxacin</b>
		<b>1 drop 4 times a day  (breakfast, lunch, dinner, bedtime)</b>
		<b>Ketorolac</b>
		<b>1 drop 4 times a day  (breakfast, lunch, dinner, bedtime)</b>

**Install 1 drop. Wait 5 minutes then install next drop.**



<b>After Surgery Instructions:</b>			
<b>First week after surgery</b>	<b>Resume drops after surgery for 1 week</b>	<b>Moxifloxacin/ Ofloxacin</b>	<b>Ketorolac</b>
		<b>1 drop 4 times a day for 1 week (then stop)</b>	<b>1 drop 4 times a day (resume)</b>
			<b>Prednisolone</b>
			<b>1 drop 4 times a day – shake bottle (resume)</b>
<b>Resume for 5 weeks</b>	<b>Resume drops for 5 additional weeks</b>	<b>Ketorolac</b>	
		<b>1 drop 4 times a day for 5 weeks (then stop)</b>	
		<b>Prednisolone</b>	
		<b>1 drop 2 times a day for 5 weeks – shake bottle (then stop)</b>	