

INDIANA REGIONAL MEDICAL CENTER
Anesthesia Protocol ---- **TABLE 1**
Table 1 ALL patients except those on Table 2

INDICATION	EKG	CBC	BMP	PT/INR - PTT	T&S	XM	K+ DOS	PT Day of Surgery	PTT Day of Surgery	FBS Day of Surgery	HCG Day of Surgery
AGE > 60 yrs	✓	✓	✓								
CARDIO											
Hx Angina (chest pain)	✓	✓									
Hx Angioplasty	✓	✓									
Hx Aortic Aneurysm	✓	✓	✓								
Hx Arrhythmia	✓	✓	✓								
Hx Bypass surgery	✓	✓	✓								
Hx Congenital Heart Disease	✓	✓	✓								
Hx Valve Replacement	✓	✓	✓								
Hypertension, medicated	✓	✓	✓								
PVD	✓	✓									
Valvular Heart Disease	✓										
Hx CHF	✓	✓	✓								
Hx AICD, pacemaker	✓	✓	✓								
Hx MI	✓	✓	✓								
PULMONARY											
COPD/Pulmonary	✓	✓	✓								
O2 – supplemental use	✓	✓									
Smoking >20 pk/yr	✓	✓									
Hx Orthopnea	✓	✓	✓								
Hx PND	✓										
ENDOCRINE											
Diabetic, medicated	✓		✓							✓	
Hyperparathyroid disease			✓								
Renal Failure	✓	✓	✓	✓							
Hx Dialysis	✓	✓	✓				✓	✓	✓		
Chance of Pregnancy											✓

INDIANA REGIONAL MEDICAL CENTER

Anesthesia Protocol ---- **TABLE 1**

Table 1 ALL patients except those on Table 2

INDICATION	EKG	CBC	BMP	PT/INR - PTT	T&S	XM	K+ DOS	PT Day of Surgery	PTT Day of Surgery	FBS Day of Surgery	HCG Day of Surgery
NEURO											
CNS disease/CVA/TIA	✓	✓	✓								
BLOOD											
Bleeding disorders (patient or blood relative)		✓		✓							
Liver disease	✓	✓	✓	✓							
Hx Sickle Cell		✓									
OTHERS											
Hx Carotid Endarterectomy	✓	✓	✓								
Morbid Obesity – BMI >39	✓										
METS < 4 (determine only if Docusys completed)	✓										
MEDICATION USE:											
Ace Inhibitor	✓		✓								
Angiotensions Receptor Blocker			✓								
Anti Arrhythmics	✓		✓								
Argatroban								✓	✓		
Beta Blockers with Diuretic			✓								
Coumadin								✓			
Diuretic			✓								
Factor Preparations								✓	✓		
Florinef			✓								
Heparin								✓	✓		
Inj. Antidiabetic-non insulin	✓		✓							✓	
Lanoxin	✓		✓								
Nitrates	✓										

*If EKG is done within 6 months at IRMC, no need to repeat unless patient is symptomatic or change in condition.