

---

Mercyhealth Prehospital and Emergency Services Center  
January 2021 CEU Quiz (All Levels)

Name \_\_\_\_\_

Level \_\_\_\_\_

Agency \_\_\_\_\_

1. In order to properly maintain the patency of an airway and align the c-spine of infants and children less than 2 years old, you should place padding under the head.
  - A. True
  - B. False
  
2. What observations are included in the pediatric assessment triangle?
  - A. Airway, Breathing, Circulation
  - B. Appearance, Work of breathing, Level of consciousness
  - C. Appearance, Work of breathing, Circulation to the skin
  
3. Which of the following is a quick, easy, and reliable sign of shock in pediatric patients?
  - A. Tachycardia
  - B. Delayed capillary refill
  - C. Hypotension
  - D. Diarrhea
  
4. What works well as a vagal maneuver for infants?
  - A. Have them bear down
  - B. Have them blow into a syringe
  - C. Ice bath
  - D. Vagal maneuvers do not work for infants
  
5. What is the proper depth for chest compressions for an infant?
  - A. 2 inches
  - B. About 2 inches
  - C. About one half the depth of the chest
  - D. About one third the depth of the chest
  
6. What is the pediatric dose of dextrose 10%?
  - A. 250ml bolus
  - B. 5ml/kg
  - C. 125ml bolus
  - D. 10ml/kg
  
7. For children ages 1 to 10, what is considered systolic hypotension?
  - A. 90 plus 2 times their age
  - B. 80 plus 2 times their age

- C. Below 100
  - D. 70 plus 2 times their age
8. What is the proper ventilation rate for an infant in cardiac arrest with an advanced airway in place?
- A. 20-30 BPM
  - B. 6 BPM (1 every 10 sec.)
  - C. 10 BPM (1 every 6 sec)
  - D. 12-20 BPM
9. What is the pediatric dose of Amiodarone during cardiac arrest?
- A. 300mg IVP
  - B. 150mg IVP
  - C. 0.1mg/kg IVP
  - D. 5mg/kg IVP
10. What is the initial dose of Adenosine for the pediatric patient?
- A. 1.0mg/kg
  - B. 0.01mg/kg
  - C. 0.1mg/kg
  - D. 6mg