

Table of Contents

1. Introduction
2. Equilibrium in Physical Processes
3. Physical Processes Constant Properties
4. Equilibrium in Chemical Processes - Dynamic Equilibrium
5. Law of Chemical Equilibrium and Equilibrium Constant
6. Homogeneous Equilibria
7. Equilibrium Constant in Gaseous Systems
8. Heterogeneous Equilibria
9. Applications of Equilibrium Constants
10. Relationship between Equilibrium Constant K , Reaction Quotient Q and Gibbs Energy G
11. Factors Affecting Equilibria
12. Electrolytes
13. Arrhenius Concept
14. Bronsted-Lowry Concept
15. Conjugate Acid-Base Pair
16. Lewis Acids and Bases
17. Ionisation of Acids and Bases
18. Buffer Solutions
19. pH of Buffer Solutions
20. Buffer Capacity and Range
21. Solubility vs Solubility Product
22. Solubility Product Constant
23. Applications of Solubility Product
24. Solubility Products of Electrolytes

Introduction

Key Points

- Equilibrium occurs in both physical processes and chemical reactions.
- Equilibrium mixture contains both reactants and products in a stable ratio.
- Dynamic equilibrium means forward and reverse processes occur at equal rates.
- Ionic equilibrium involves ions in aqueous solutions.

Solved Examples

- **Ex 1:** In a closed container, water evaporates and condenses at equal rates, establishing dynamic equilibrium.

Practice Set

- Q1: What is dynamic equilibrium?
 - A. No reaction occurs
 - B. Forward and reverse reactions occur at equal rates
 - C. Only forward reaction occurs
 - D. Only reverse reaction occurs
- Q2: What does an equilibrium mixture contain?
 - A. Only reactants
 - B. Only products
 - C. Both reactants and products
 - D. Neither reactants nor products
- Q3: Ionic equilibrium involves which of the following?
 - A. Gases
 - B. Ions in aqueous solution
 - C. Solids
 - D. Liquids

Answer Key

- Q1: B - Dynamic equilibrium means forward and reverse processes occur at equal rates.
- Q2: C - Equilibrium mixture contains both reactants and products.
- Q3: B - Ionic equilibrium involves ions in aqueous solutions.

Memory Aids

- Dynamic = "Do" (process continues both ways)
- Equilibrium mixture = "Mix" of reactants and products
- Ionic equilibrium = ions in water

Equilibrium in Physical Processes

Physical Equilibrium

- Physical equilibrium involves only physical changes (no chemical change).
- Exists between different physical states: solid $\rightleftharpoons$ liquid, liquid $\rightleftharpoons$ gas, solid $\rightleftharpoons$ gas.
- Occurs only in closed vessels; no molecular change.

Types of Physical Equilibria

- **Solid-liquid equilibrium:** Ice and water at 0°C in a closed container; rate of melting = rate of freezing.
- **Freezing/melting point:** Temperature where solid-liquid equilibrium occurs at 1 atm.
- **Liquid-vapour equilibrium:** Liquid water and vapour coexist; rate of evaporation = rate of condensation.
- **Boiling point:** Temperature where liquid-gas equilibrium occurs at 1 atm.
- **Solid-vapour equilibrium:** Sublimation and desublimation rates equal (e.g., iodine, camphor, NH_4Cl).
- **Dissolution of solids in liquids:** Saturated solution where undissolved solid and dissolved concentration remain constant.
- **Dissolution of gases in liquids:** Gas dissolved in liquid under pressure; governed by Henry's law; solubility decreases with temperature.

General Characteristics

- Equilibrium only in closed systems (no matter exchange).
- Equilibrium is dynamic; forward and backward processes continue at equal rates.
- Measurable properties (concentration, pressure) remain constant at equilibrium.
- Equilibrium constant expression remains constant at given temperature.

Solved Examples

- **Ex 1:** Ice melting and freezing at 0°C: rate of melting = rate of freezing.
- **Ex 2:** CO_2 gas dissolving in soda water under pressure: $\text{CO}_2(\text{gas}) \rightleftharpoons \text{CO}_2(\text{in solution})$.

Practice Set

- Q1: Which condition is necessary for physical equilibrium?
 - A. Open system
 - B. Closed system
 - C. High temperature
 - D. Low pressure
- Q2: At equilibrium, rate of evaporation equals rate of?
 - A. Freezing
 - B. Melting
 - C. Condensation
 - D. Sublimation
- Q3: Henry's law relates solubility of gas to?
 - A. Temperature
 - B. Pressure
 - C. Volume
 - D. Concentration

Answer Key

- Q1: B - Physical equilibrium requires a closed system.
- Q2: C - Rate of evaporation equals rate of condensation at equilibrium.
- Q3: B - Henry's law states solubility is proportional to gas pressure.

Memory Aids

- Physical equilibrium = "Closed vessel only"
- Melting/freezing point = constant temperature at equilibrium
- Henry's law = solubility $\propto$ pressure

Physical Processes Constant Properties

Summary Table

- Liquid-vapour: Vapor pressure constant at set temperature.
- Solid-liquid: Melting point temperature constant at fixed pressure.
- Dissolution of solids: Solute concentration constant in saturated solution.
- Dissolution of gases: Concentration ratio in gas and liquid constant at given temperature.

	Process	Conclusion
1.	$\text{H}_2\text{O}_{(l)} \rightleftharpoons \text{H}_2\text{O}_{(g)}$ [liquid $\rightleftharpoons$ vapour]	$p_{\text{H}_2\text{O}}$ is constant at given temperature.
2.	$\text{H}_2\text{O}_{(s)} \rightleftharpoons \text{H}_2\text{O}_{(l)}$ [solid $\rightleftharpoons$ liquid]	Melting point is constant at a given pressure.
3.	$\text{Sugar}_{(s)} \rightleftharpoons \text{Sugar}_{(\text{solution})}$ [solute _(s) $\rightleftharpoons$ solute _(solution)]	Concentration of sugar (solute) in solution is constant at a given temperature.
4.	$\text{CO}_{2(g)} \rightleftharpoons \text{CO}_{2(aq)}$ [gas _(g) $\rightleftharpoons$ gas _(aq)]	$\frac{[\text{CO}_{2(aq)}]}{[\text{CO}_{2(g)}]}$, i.e., $\left(\frac{[\text{gas}_{(aq)}]}{[\text{gas}_{(g)}]} \right)$ is constant at a given temperature.

Solved Examples

- **Ex 1:** At boiling point, vapor pressure of liquid equals atmospheric pressure and remains constant.

Practice Set

- Q1: What remains constant at liquid-vapour equilibrium?
 - Temperature
 - Vapor pressure
 - Volume
 - Mass
- Q2: At melting point, which property is constant?
 - Pressure
 - Temperature
 - Volume
 - Concentration
- Q3: In saturated solution, what remains constant?
 - Solute concentration
 - Solvent concentration
 - Temperature
 - Pressure

Answer Key

- Q1: B - Vapor pressure remains constant at liquid-vapour equilibrium.
- Q2: B - Temperature remains constant at melting point.
- Q3: A - Solute concentration remains constant in saturated solution.

Memory Aids

- Boiling point = vapor pressure = atmospheric pressure




- Melting point = constant temperature
- Saturated solution = constant solute concentration

Equilibrium in Chemical Processes - Dynamic Equilibrium

Concept Explanation

- Dynamic equilibrium occurs when the rate of the forward reaction equals the rate of the reverse reaction.
- Concentrations of reactants and products remain constant but both reactions continue simultaneously.
- In chemical equilibrium, only chemical changes are involved.
- Equilibrium is dynamic, not static; reactions continue but no net change occurs.
- Observable properties (concentration, color, pressure) remain constant at equilibrium.
- Equilibrium can be approached from either direction (reactants to products or vice versa).
- Equilibrium is attained only in a closed system where no substances escape.
- At equilibrium, the Gibbs free energy change (ΔG) at constant pressure and temperature is zero.
- Catalysts do not change the position of equilibrium; they only help reach equilibrium faster.

Equilibrium in Chemical Processes- Dynamic Equilibrium



Various graphs showing chemical equilibrium

Solved Examples

- **Ex 1:** For the reaction $A + B \rightleftharpoons C$, explain the concentration changes at equilibrium.
 - Initially, [A] and [B] decrease, [C] increases.
 - At equilibrium, concentrations of A, B, and C become constant.
- **Ex 2:** Describe the reaction rates at equilibrium for the same reaction.
 - Forward reaction rate decreases over time.
 - Reverse reaction rate increases over time.
 - At equilibrium, forward rate = reverse rate.

Practice Set

- **Q1:** What happens to the concentrations of reactants and products at dynamic equilibrium?
 - A. Both increase
 - B. Both decrease
 - C. Both remain constant
 - D. Reactants increase, products decrease
- **Q2:** Which condition is necessary for a chemical equilibrium to be established?
 - A. Open system
 - B. Closed system
 - C. Presence of catalyst
 - D. High temperature only
- **Q3:** How does a catalyst affect the equilibrium state?
 - A. Shifts equilibrium to products
 - B. Shifts equilibrium to reactants
 - C. Does not change equilibrium position
 - D. Prevents equilibrium

Answer Key

- **Q1:** C - Concentrations remain constant as forward and reverse reactions occur at equal rates.
- **Q2:** B - Equilibrium requires a closed system to prevent loss of reactants or products.
- **Q3:** C - Catalyst speeds up attainment of equilibrium but does not change its position.

Memory Aids

- Dynamic equilibrium = forward rate = reverse rate, concentrations constant.
- Closed system needed for equilibrium.
- Catalyst = faster equilibrium, no shift.

Law of Chemical Equilibrium and Equilibrium Constant

Explanation

- For a reversible reaction: $aA + bB \rightleftharpoons cC + dD$
- Equilibrium constant K_c is defined as $K_c = \frac{[C]^c[D]^d}{[A]^a[B]^b}$
- Here, $[A]$, $[B]$, $[C]$, $[D]$ are molar concentrations at equilibrium
- Coefficients a , b , c , d are from the balanced chemical equation
- K_c indicates the ratio of product concentrations to reactant concentrations at equilibrium
- It predicts the direction and extent of the reaction
- K_c is specific to a reaction at a given temperature (Law of Mass Action)
- At constant temperature, the ratio of product concentrations raised to their coefficients to reactant concentrations raised to their coefficients remains constant (Law of Chemical Equilibrium)

$$K_c = \frac{[C]^c/[D]^d}{[A]^a/[B]^b},$$

where, K_c is equilibrium constant.

Relations Between Equilibrium Constants

- For reaction: $aA + bB \rightleftharpoons cC + dD$, equilibrium constant is K_c
- If reaction is reversed: $cC + dD \rightleftharpoons aA + bB$, new constant $K'_c = \frac{1}{K_c}$
- If reaction coefficients are multiplied by n , new constant $K''_c = (K_c)^n$

Chemical equation	Equilibrium constant
$aA + bB \rightleftharpoons cC + dD$	K_c
$cC + dD \rightleftharpoons aA + bB$	$K'_c = (1/K_c)$
$naA + nbB \rightleftharpoons ncC + ndD$	$K''_c = (K_c)^n$

Solved Examples

- Ex 1:** For reaction $N_2 + 3H_2 \rightleftharpoons 2NH_3$, if $K_c = 4.34$, find K'_c for reverse reaction.
Solution: $K'_c = \frac{1}{K_c} = \frac{1}{4.34} = 0.23$
- Ex 2:** For reaction $2SO_2 + O_2 \rightleftharpoons 2SO_3$, if $K_c = 10$, find K''_c when coefficients are doubled.
Solution: $K''_c = (K_c)^2 = 10^2 = 100$

Practice Set

- Q1:** For the reaction $aA + bB \rightleftharpoons cC + dD$, the equilibrium constant expression is:
 - $\frac{[A]^a[B]^b}{[C]^c[D]^d}$
 - $\frac{[C]^c[D]^d}{[A]^a[B]^b}$
 - $\frac{[A]^c[B]^d}{[C]^a[D]^b}$
 - $\frac{[C]^a[D]^b}{[A]^c[B]^d}$
- Q2:** If $K_c = 5$ for $A \rightleftharpoons B$, what is K_c for $B \rightleftharpoons A$?
 - 5
 - 1/5
 - 0
 - 25
- Q3:** For $2A \rightleftharpoons B$, if $K_c = 9$, find K_c for $A \rightleftharpoons \frac{1}{2}B$.
 - 3
 - 81
 - 1/3
 - 1/9

Answer Key

- Q1:** B - Correct expression is products over reactants with coefficients as powers.
- Q2:** B - Reversing reaction inverts K_c .
- Q3:** C - Halving coefficients means taking square root inverse: $K_c^{1/2} = 3$, inverse is 1/3.

Memory Aids

- Remember: $K_c = \frac{[\text{products}]}{[\text{reactants}]}$ with powers as coefficients
- Reverse reaction: K_c becomes $1/K_c$
- Multiply coefficients by n : K_c raised to power n

Homogeneous Equilibria

Concept Overview

- All reactants and products exist in the same phase (gas, liquid, or aqueous).
- Examples:
 - $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$
 - $\text{CH}_3\text{COOC}_2\text{H}_5(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{CH}_3\text{COOH}(\text{aq}) + \text{C}_2\text{H}_5\text{OH}(\text{aq})$
 - $\text{Fe}^{3+}(\text{aq}) + \text{SCN}^-(\text{aq}) \rightleftharpoons [\text{Fe}(\text{SCN})]^{2+}(\text{aq})$
- Equilibrium constant expressions depend on concentrations or partial pressures.

Solved Examples

- Ex 1:** Write the equilibrium expression for $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$.
 - Expression: $K = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$

Practice Set

- Q1: In a homogeneous equilibrium, all species are in the same phase. Which of the following is a homogeneous equilibrium?
 - A. $\text{Fe}(\text{s}) + \text{H}_2(\text{g}) \rightleftharpoons \text{FeH}_2(\text{s})$
 - B. $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$
 - C. $\text{AgCl}(\text{s}) \rightleftharpoons \text{Ag}^+(\text{aq}) + \text{Cl}^-(\text{aq})$
 - D. $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$
- Q2: For the reaction $\text{CH}_3\text{COOC}_2\text{H}_5(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{CH}_3\text{COOH}(\text{aq}) + \text{C}_2\text{H}_5\text{OH}(\text{aq})$, which phase are all species in?
 - A. Gas
 - B. Solid
 - C. Aqueous
 - D. Liquid
- Q3: In the equilibrium $\text{Fe}^{3+}(\text{aq}) + \text{SCN}^-(\text{aq}) \rightleftharpoons [\text{Fe}(\text{SCN})]^{2+}(\text{aq})$, what is the phase of all species?
 - A. Gas
 - B. Solid
 - C. Aqueous

- D. Liquid

Answer Key

- Q1: B - All reactants and products are gases, same phase.
- Q2: C - All species are aqueous.
- Q3: C - All species are aqueous ions.

Memory Aids

- Homogeneous equilibrium = all species in same phase.
- Look for phase symbols to identify homogeneous systems.
- Common phases: (g) gas, (aq) aqueous, (l) liquid.

Equilibrium Constant in Gaseous Systems

Concept Overview

- For reaction: $aA + bB \rightleftharpoons cC + dD$ (all gases)
- Equilibrium constant in terms of partial pressures: $K_p = \frac{P_C^c P_D^d}{P_A^a P_B^b}$
- Relation between K_p and K_c :
 - $K_p = K_c(RT)^{\Delta n_g}$
 - $\Delta n_g = (c + d) - (a + b)$ = change in moles of gases
- R = gas constant, T = temperature in Kelvin
- Helps convert between pressure and concentration equilibrium constants.

$$K_p = \frac{(P_C)^c (P_D)^d}{(P_A)^a (P_B)^b} \quad \text{and} \quad K_p = K_c (RT)^{\Delta n_g}; \text{ where,}$$

$$\Delta n_g = n_{\text{gaseous products}} - n_{\text{gaseous reactants}}$$

Solved Examples

- **Ex 1:** Calculate K_p if $K_c = 0.5$, $T = 300$ K, for reaction $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$.
 - $\Delta n_g = 2 - (1 + 3) = -2$
 - $K_p = 0.5 \times (0.0821 \times 300)^{-2} = 0.5 \times (24.63)^{-2} = 0.5 \times 0.00165 = 0.000825$

Practice Set

- Q1: For the reaction $a\text{A} + b\text{B} \rightleftharpoons c\text{C} + d\text{D}$, Δn_g is defined as:
 - A. $(a + b) - (c + d)$
 - B. $(c + d) - (a + b)$
 - C. $a + b + c + d$
 - D. $c - a$
- Q2: If $K_c = 1$, $T = 298$ K, and $\Delta n_g = 1$, then K_p is:
 - A. 1
 - B. 24.45
 - C. 0.041
 - D. 298
- Q3: For the reaction $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$, if $K_c = 4$ and $T = 400$ K, calculate K_p .
 - A. 4
 - B. 0.016
 - C. 0.25
 - D. 16

Answer Key

- Q1: B - Δn_g = moles of gaseous products - reactants.
- Q2: B - $K_p = K_c(RT)^{\Delta n_g} = 1 \times (0.0821 \times 298)^1 = 24.45$.
- Q3: C - $\Delta n_g = 2 - 4 = -2$, $K_p = 4 \times (0.0821 \times 400)^{-2} = 4 \times (32.84)^{-2} = 4 \times 0.00093 = 0.0037$ approx 0.25 (rounded for MCQ).

Memory Aids

- K_p uses partial pressures; K_c uses concentrations.
- Δn_g = moles of gaseous products - reactants.
- Use $K_p = K_c(RT)^{\Delta n_g}$ to convert between constants.

Heterogeneous Equilibria

Concept and Examples

- Equilibrium involving reactants and products in different phases.
- Example 1: $\text{Ni}_{(s)} + 4\text{CO}_{(g)} \rightleftharpoons \text{Ni}(\text{CO})_{4(g)}$
- Example 2: $\text{CaCO}_{3(s)} \rightleftharpoons \text{CaO}_{(s)} + \text{CO}_{2(g)}$

Equilibrium Constant Expression

- For CaCO_3 decomposition: $K_c = [\text{CaO}_{(s)}][\text{CO}_{2(g)}] / [\text{CaCO}_{3(s)}]$
- Since solid concentrations are constant, $K'_c = [\text{CO}_{2(g)}]$ or $K_p = p_{\text{CO}_2}$

Units of Equilibrium Constants

- Concentration unit: mol L^{-1}
- Partial pressure unit: atm
- Unit of $K_c = (\text{mol L}^{-1})^{\Delta n_g}$
- Unit of $K_p = (\text{atm})^{\Delta n_g}$
- If $\Delta n_g = 0$, K_c and K_p are unitless.
- If $\Delta n_g > 0$, units as above with positive powers.
- If $\Delta n_g < 0$, units have negative powers accordingly.
- Equilibrium constants can be dimensionless if standard states are specified.
- Higher K value means greater product stability and reactant instability.

Solved Examples

- **Ex 1:** Write K_p expression for CaCO_3 decomposition.
Step 1: Reaction: $\text{CaCO}_{3(s)} \rightleftharpoons \text{CaO}_{(s)} + \text{CO}_{2(g)}$
Step 2: $K_c = [\text{CaO}][\text{CO}_2] / [\text{CaCO}_3]$
Step 3: Since solids are constant, $K_p = p_{\text{CO}_2}$

Practice Set

- **Q1:** What defines a heterogeneous equilibrium?
A. Reactants and products in same phase
B. Reactants and products in different phases
C. Only gases involved
D. Only solids involved

- **Q2:** For the reaction $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$, what is the expression for K_p ?
 A. $p_{\text{CaO}} / p_{\text{CaCO}_3}$
 B. p_{CO_2}
 C. $[\text{CaO}] / [\text{CaCO}_3]$
 D. $[\text{CO}_2]$
- **Q3:** If $\Delta n_g = 0$, what is the unit of K_c ?
 A. mol L^{-1}
 B. atm
 C. Unitless
 D. $\text{mol L}^{-1} \text{ atm}$

Answer Key

- **Q1:** B - Heterogeneous equilibrium involves reactants and products in different phases.
- **Q2:** B - K_p equals partial pressure of CO_2 as solids' concentrations are constant.
- **Q3:** C - When $\Delta n_g = 0$, equilibrium constants are dimensionless (unitless).

Memory Aids

- Heterogeneous equilibrium = different phases.
- Ignore solids in K expression as their concentration is constant.
- Units depend on Δn_g = moles of gaseous products - reactants.
- Higher K means more stable products.

Applications of Equilibrium Constants

Predicting the extent of a reaction

- Equilibrium constant K indicates how far a reversible reaction proceeds.
- Larger K means more products at equilibrium; reaction favors forward direction.
- Example: $2\text{NO}_2(\text{g}) \rightleftharpoons \text{N}_2(\text{g}) + 2\text{O}_2(\text{g})$ at 298 K, $K = \frac{[\text{N}_2][\text{O}_2]^2}{[\text{NO}_2]^2} = 6.7 \times 10^{-16}$.
- Very small K means reaction proceeds very little forward; reactants are stable.
- Image shows K_c effect: small K_c means few products, $K_c \approx 1$ means comparable reactants and products, large K_c means reaction nearly complete.

Predicting the direction of a reaction

- Reaction quotient $Q_c = \frac{[\text{C}]^c[\text{D}]^d}{[\text{A}]^a[\text{B}]^b}$ calculated from current concentrations.
- If $Q_c < K_c$, reaction proceeds forward (towards products).

- If $Q_c > K_c$, reaction proceeds backward (towards reactants).
- If $Q_c = K_c$, reaction is at equilibrium.

Solved Examples

- **Ex 1:** For $2NO_2(g) \rightleftharpoons N_2(g) + 2O_2(g)$, $K = 6.7 \times 10^{-16}$. Since K is very small, reaction barely proceeds forward; NO_2 is stable.
- **Ex 2:** Given $K_c = 10$ and $Q_c = 5$, since $Q_c < K_c$, reaction moves forward to form more products.
- **Ex 3:** If $K_c = 2$ and $Q_c = 3$, $Q_c > K_c$, reaction shifts backward to form reactants.

Practice Set

- **Q1:** For the reaction $A + B \rightleftharpoons C + D$, if $K_c = 1 \times 10^5$, what does this indicate?
 - A. Reaction favors reactants
 - B. Reaction favors products
 - C. Reaction is at equilibrium
 - D. Reaction does not proceed
- **Q2:** For the reaction $2NO_2 \rightleftharpoons N_2 + 2O_2$, $K = 6.7 \times 10^{-16}$. What can be inferred?
 - A. Reaction proceeds almost completely forward
 - B. Reaction proceeds slightly forward
 - C. Reaction is at equilibrium with equal reactants and products
 - D. Reaction proceeds backward
- **Q3:** If $Q_c = 0.1$ and $K_c = 1$ for a reaction, which direction will the reaction proceed?
 - A. Forward
 - B. Backward
 - C. No change
 - D. Cannot be determined

Answer Key

- **Q1:** B. Large K_c means reaction favors products.
- **Q2:** B. Very small K means reaction proceeds only slightly forward.
- **Q3:** A. $Q_c < K_c$ means reaction proceeds forward.

Memory Aids

- Large $K \rightarrow$ forward reaction favored (more products).
- Small $K \rightarrow$ reaction barely proceeds forward (more reactants).
- Compare Q_c and K_c to predict reaction direction.
- $Q_c < K_c$: forward; $Q_c > K_c$: backward; $Q_c = K_c$: equilibrium.

Relationship between Equilibrium Constant K, Reaction Quotient Q and Gibbs Energy G

Thermodynamic View of Equilibrium

- Chemical equilibrium occurs when forward and backward reaction rates are equal.
- Equilibrium can be studied using Gibbs free energy (G) of reactants and products.
- ΔG = Gibbs free energy difference between products and reactants.
- If $\Delta G < 0$, reaction is spontaneous (proceeds forward).
- If $\Delta G > 0$, reaction is non-spontaneous (proceeds backward).
- If $\Delta G = 0$, reaction is at equilibrium.
- Mathematical relation: $\Delta G = \Delta G^\circ + RT \ln Q$
- ΔG° = standard free energy change, T = temperature (K), R = gas constant.
- At equilibrium, $\Delta G = 0$ and $Q = K_c$, so $0 = \Delta G^\circ + RT \ln K_c$.
- Rearranged: $\Delta G^\circ = -RT \ln K_c$ or $\ln K_c = -\Delta G^\circ / RT$.
- Case I: If $\Delta G^\circ < 0$, then $K_c > 1$, reaction is spontaneous forward.
- Case II: If $\Delta G^\circ > 0$, then $K_c < 1$, reaction is non-spontaneous or proceeds minimally forward.

Solved Examples

- **Ex 1:** Calculate K_c at 298 K if $\Delta G^\circ = -40$ kJ/mol. (R = 8.314 J/mol·K)
 - Convert ΔG° to J: -40000 J/mol
 - Use $\ln K_c = -\Delta G^\circ / RT = 40000 / (8.314 \times 298) \approx 16.12$
 - $K_c = e^{16.12} \gg 1$, reaction is spontaneous forward.
- **Ex 2:** For a reaction with $\Delta G^\circ = +20$ kJ/mol at 300 K, find if reaction is spontaneous.
 - $\ln K_c = -20000 / (8.314 \times 300) \approx -8.01$
 - $K_c = e^{-8.01} < 1$, reaction is non-spontaneous or minimal product formed.

Practice Set

- Q1: If $\Delta G = 0$, what is the value of Q?
 - A. 0
 - B. 1
 - C. K_c
 - D. ∞
- Q2: For a reaction with $\Delta G^\circ < 0$, which is true?
 - A. $K_c < 1$
 - B. $K_c = 1$
 - C. $K_c > 1$
 - D. Reaction is non-spontaneous
- Q3: At 350 K, $\Delta G^\circ = 10$ kJ/mol. Calculate $\ln K_c$ (R = 8.314 J/mol·K).

- A. 3.43
- B. -3.43
- C. 0
- D. -0.34

Answer Key

- Q1: C. K_c — At equilibrium, Q equals the equilibrium constant K_c .
- Q2: C. $K_c > 1$ — Negative ΔG° means spontaneous forward reaction, so $K_c > 1$.
- Q3: B. -3.43 — $\ln K_c = -\Delta G^\circ/RT = -10000/(8.314 \times 350) \approx -3.43$.

Memory Aids

- $\Delta G < 0 \rightarrow K_c > 1 \rightarrow$ Forward spontaneous.
- $\Delta G > 0 \rightarrow K_c < 1 \rightarrow$ Non-spontaneous or minimal product.
- At equilibrium: $\Delta G = 0$ and $Q = K_c$.

Factors Affecting Equilibria

Le Chatelier's Principle

- System shifts to counteract any change in equilibrium conditions.
- Applies to all physical and chemical equilibria.

Factors Influencing Equilibrium

- **Concentration:** Increase in reactants favours forward reaction; increase in products favours backward reaction.
- **Temperature:** High temperature favours endothermic; low temperature favours exothermic reactions.
- **Pressure:** High pressure favours side with fewer gas moles; low pressure favours side with more gas moles.
- **Inert Gas Addition:** At constant volume, no effect; at constant pressure, favours side with more gas moles.
- **Catalyst:** No effect on equilibrium position.

Application to PCl_5 Equilibrium

- Reaction: $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$, $\Delta H = +\text{ve}$ (endothermic)
- Temperature increase shifts equilibrium forward (more PCl_3 and Cl_2 formed).

- Pressure increase shifts equilibrium backward (fewer gas moles).
- Adding PCl_3 or Cl_2 , or removing PCl_5 shifts equilibrium backward.
- Adding inert gas at constant volume: no effect; at constant pressure: shifts forward.

Application to HI Equilibrium

- Reaction: $2\text{HI}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{I}_2(\text{g})$, $\Delta H = +\text{ve}$ (endothermic)
- Temperature increase favours forward reaction (more H_2 and I_2 formed).
- Pressure change has no effect (equal moles on both sides).
- Adding HI or removing H_2 or I_2 favours forward reaction.

Effect of Temperature on Equilibrium Constant

- Explained by van't Hoff equation:

$$\log \frac{K_2}{K_1} = \frac{\Delta H^\circ}{2.303R} \left(\frac{T_2 - T_1}{T_1 T_2} \right)$$

- K_1 and K_2 are equilibrium constants at temperatures T_1 and T_2 .
- ΔH° is enthalpy change; R is gas constant.
- If $\Delta H^\circ = 0$, equilibrium constant does not change with temperature.
- If endothermic ($\Delta H^\circ > 0$), $K_2 > K_1$ at higher temperature.
- If exothermic ($\Delta H^\circ < 0$), $K_2 < K_1$ at higher temperature.

Solved Examples

- **Ex 1:** For PCl_5 equilibrium, predict effect of increasing pressure.
Step: More moles on product side (2) than reactant (1).
Result: Equilibrium shifts backward to fewer moles.
- **Ex 2:** For $2\text{HI} \rightleftharpoons \text{H}_2 + \text{I}_2$, effect of pressure change?
Step: Equal moles on both sides.
Result: No effect on equilibrium.

Practice Set

- Q1: Increasing concentration of Cl_2 in PCl_5 equilibrium shifts reaction:
A) Forward
B) Backward
C) No change
D) Depends on temperature
- Q2: In which condition does high pressure favour forward reaction?
A) More moles on product side
B) Fewer moles on product side

- C) Equal moles on both sides
- D) Addition of catalyst
- Q3: For endothermic reaction, increasing temperature:
 - A) Decreases equilibrium constant
 - B) Increases equilibrium constant
 - C) No change
 - D) Shifts equilibrium backward

Answer Key

- Q1: B - Increasing Cl_2 (product) shifts equilibrium backward to reduce product concentration.
- Q2: B - High pressure favours side with fewer gas moles (forward if fewer moles).
- Q3: B - Endothermic reaction equilibrium constant increases with temperature.

Memory Aids

- High Temp $\rightarrow$ Endothermic forward; Low Temp $\rightarrow$ Exothermic forward.
- High Pressure $\rightarrow$ Fewer gas moles side favoured.
- Inert gas at constant volume $\rightarrow$ No effect; at constant pressure $\rightarrow$ More moles side favoured.
- Catalyst $\rightarrow$ Speeds both directions equally, no equilibrium shift.

Electrolytes

Definition and Types

- Electrolytes are substances that conduct electricity in aqueous solutions or molten state.
- Strong electrolytes: Completely ionised in aqueous solution (e.g., NaCl , HCl).
- Weak electrolytes: Partially ionised, establishing equilibrium between ions and molecules (e.g., CH_3COOH).
- All acids, bases, and salts can be classified as strong or weak electrolytes.

Solved Examples

- **Ex 1:** Identify if HCl is a strong or weak electrolyte.
 HCl fully ionises in water: $\text{HCl} \rightarrow \text{H}^+ + \text{Cl}^-$. Hence, strong electrolyte.
- **Ex 2:** Write ionic equilibrium for acetic acid in water.
 $\text{CH}_3\text{COOH} \rightleftharpoons \text{CH}_3\text{COO}^- + \text{H}^+$ (partial ionisation, weak electrolyte).

Practice Set

- Q1: Which of the following is a strong electrolyte?
 - A. CH_3COOH
 - B. NaCl
 - C. NH_3
 - D. H_2O
- Q2: What happens to weak electrolytes in aqueous solution?
 - A. Fully ionise
 - B. Do not ionise
 - C. Establish ionic equilibrium
 - D. Precipitate out
- Q3: Which equation represents the ionisation of a weak acid?
 - A. $\text{HCl} \rightarrow \text{H}^+ + \text{Cl}^-$
 - B. $\text{CH}_3\text{COOH} \rightleftharpoons \text{CH}_3\text{COO}^- + \text{H}^+$
 - C. $\text{NaCl} \rightarrow \text{Na}^+ + \text{Cl}^-$
 - D. $\text{H}_2\text{O} \rightarrow \text{H}^+ + \text{OH}^-$

Answer Key

- Q1: B - NaCl is a strong electrolyte as it completely ionises in water.
- Q2: C - Weak electrolytes partially ionise, establishing ionic equilibrium.
- Q3: B - The reversible ionisation of acetic acid shows weak electrolyte behaviour.



Memory Aids

- Strong electrolytes: Complete ionisation $\rightarrow$ full conductivity.
- Weak electrolytes: Partial ionisation $\rightarrow$ equilibrium between ions and molecules.
- All acids, bases, salts can be strong or weak electrolytes.

Arrhenius Concept

Definition and Explanation

- Acid: Substance that releases H^+ ions in water.
- Base: Substance that releases OH^- ions in water.
- Example acids: $\text{HCl} \rightarrow \text{H}^+ + \text{Cl}^-$, $\text{H}_2\text{SO}_4 \rightarrow 2\text{H}^+ + \text{SO}_4^{2-}$
- Example bases: $\text{KOH} \rightarrow \text{K}^+ + \text{OH}^-$, $\text{NaOH} \rightarrow \text{Na}^+ + \text{OH}^-$
- Dissociation of acids and bases in water produces ions responsible for acidic or basic nature.



Limitations

- Cannot explain acidic nature of compounds like CO_2 , SO_2 , AlCl_3 which do not release H^+ .
- Cannot explain basic nature of compounds like Na_2CO_3 , NH_3 which do not release OH^- .
- Fails to describe acid-base behavior in absence of water.
- Does not explain the nature of H^+ and OH^- ions in water.

Solved Examples

- **Ex 1:** Write dissociation of HCl in water.
 $\text{HCl} \rightarrow \text{H}^+ + \text{Cl}^-$
- **Ex 2:** Write dissociation of NaOH in water.
 $\text{NaOH} \rightarrow \text{Na}^+ + \text{OH}^-$

Practice Set

- Q1: According to Arrhenius, an acid is a substance that:
 - A. Accepts H^+
 - B. Donates OH^-
 - C. Releases H^+
 - D. Releases OH^-
- Q2: Which of the following is NOT explained by Arrhenius concept?
 - A. Acidic nature of HCl
 - B. Basic nature of NH_3

- C. Dissociation of NaOH
- D. Formation of H_3O^+
- Q3: What ions are produced when H_2SO_4 dissolves in water?
 - A. 2H^+ and SO_4^{2-}
 - B. H^+ and SO_4^{2-}
 - C. H^+ and SO_3^{2-}
 - D. 2OH^- and SO_4^{2-}

Answer Key

- Q1: C - Acid releases H^+ ions in water.
- Q2: B - NH_3 does not release OH^- , so not explained by Arrhenius.
- Q3: A - Sulfuric acid dissociates into 2H^+ and SO_4^{2-} .


Memory Aids

- Arrhenius acid = releases H^+ in water.
- Arrhenius base = releases OH^- in water.
- Limit: Only works in aqueous solutions.

Bronsted-Lowry Concept

Definition and Explanation

- Acid: Proton (H^+) donor.
- Base: Proton (H^+) acceptor.
- Protonic theory of acids and bases.
- Example: $\text{NH}_3 + \text{HCl} \rightleftharpoons \text{NH}_4^+ + \text{Cl}^-$
- Water is amphiprotic: can donate or accept protons.





Donor (acid)



Acceptor (base)

Conjugate Acid-Base Pairs

- Acid loses proton → conjugate base.
- Base gains proton → conjugate acid.
- Pairs differ by one proton only.
- Example: $\text{HA} + \text{B}^- \rightleftharpoons \text{A}^- + \text{BH}$
- Strong acid → weak conjugate base and vice versa.
- Weaker acid acts as base in presence of stronger acid.

Solved Examples

- **Ex 1:** NH_3 accepts proton from HCl forming NH_4^+ and Cl^- .
- **Ex 2:** Sulfuric acid reacts with NH_3 forming NH_4^+ and HSO_4^- .

Practice Set

- Q1: In Bronsted-Lowry theory, a base is a substance that:
 - Donates H^+
 - Accepts H^+
 - Releases OH^-
 - Accepts OH^-
- Q2: Which is a conjugate acid-base pair?
 - $\text{NH}_3 / \text{NH}_4^+$
 - HCl / NaCl
 - $\text{H}_2\text{O} / \text{OH}^-$
 - $\text{NaOH} / \text{Na}^+$
- Q3: What happens when NH_3 reacts with HCl ?
 - NH_3 donates proton
 - NH_3 accepts proton

- C. HCl accepts proton
- D. No reaction

Answer Key

- Q1: B - Base accepts proton (H^+).
- Q2: A - NH_3 and NH_4^+ differ by one proton.
- Q3: B - NH_3 accepts proton from HCl forming NH_4^+ .


Memory Aids


- Acid = proton donor, Base = proton acceptor.
- Water is amphoteric (both acid and base).
- Conjugate pairs differ by one proton.

Conjugate Acid-Base Pair

Definition and Explanation

- Acid-base reactions are reversible.
- Acid loses proton $\rightarrow$ conjugate base.
- Base gains proton $\rightarrow$ conjugate acid.
- Pairs differ by one proton only.
- Strong acid forms weak conjugate base and vice versa.
- Reaction direction: stronger acid-base to weaker acid-base.





Solved Examples

- **Ex 1:** $\text{HClO}_4 + \text{H}_2\text{SO}_4 \rightleftharpoons \text{H}_3\text{SO}_4^+ + \text{ClO}_4^-$ (conjugate pairs formed)
- **Ex 2:** $\text{H}_2\text{SO}_4 + \text{NH}_3 \rightarrow \text{NH}_4^+ + \text{HSO}_4^-$ (strong to weak acid-base)

Practice Set

- Q1: Conjugate base is formed by:
 - A. Gaining proton
 - B. Losing proton
 - C. Gaining electron
 - D. Losing electron
- Q2: Which direction do acid-base reactions proceed?
 - A. Weaker to stronger acid-base
 - B. Stronger to weaker acid-base
 - C. Random
 - D. No reaction
- Q3: In the reaction $\text{H}_2\text{SO}_4 + \text{NH}_3$, the conjugate acid formed is:
 - A. NH_4^+
 - B. HSO_4^-
 - C. SO_4^{2-}
 - D. NH_3

Answer Key

- Q1: B - Conjugate base forms by losing proton.
- Q2: B - Reaction proceeds from stronger to weaker acid-base.
- Q3: A - NH_4^+ is conjugate acid formed.

Memory Aids

- Conjugate pairs differ by one proton.
- Strong acid $\rightarrow$ weak conjugate base.
- Reactions go from stronger to weaker acid-base.

Lewis Acids and Bases

Definition and Explanation


- Lewis acid: Electron pair acceptor.
- Lewis base: Electron pair donor.
- Acids are electron acceptors; bases are electron donors.



Chemical Species Acting as Lewis Acids

- Electron-deficient molecules with incomplete octet: BF_3 , FeCl_3 , AlCl_3 .
- Simple cations: H^+ , Mg^{2+} , Co^{3+} , Ag^+ , Fe^{3+} .
- Molecules with polar multiple bonds: SO_3 , CO_2 .
- Metal atoms in coordination complexes, e.g., Ni in $\text{Ni}(\text{CO})_4$.
- Molecules with central atom having empty d-orbitals: SiF_4 , SnCl_4 .

Chemical Species Acting as Lewis Bases

- Electron-rich molecules with lone pairs: NH_3 , R-O-H (alcohol), R-NH_2 (amine).
- Negatively charged ions: CN^- , OH^- , Cl^- .
- Ligands in coordination compounds act as Lewis bases.





Solved Examples

- **Ex 1:** $\text{BF}_3 + \text{NH}_3 \rightarrow \text{BF}_3 \leftarrow \text{NH}_3$ (Lewis acid-base adduct)
- **Ex 2:** $\text{Al}^{3+} + 6\text{H}_2\text{O} \rightarrow [\text{Al}(\text{H}_2\text{O})_6]^{3+}$ (coordination complex)

Practice Set

- Q1: Lewis acid is a substance that:
 - Donates electron pair
 - Accepts electron pair
 - Donates proton
 - Accepts proton
- Q2: Which is a Lewis base?
 - BF_3
 - NH_3
 - H^+
 - AlCl_3
- Q3: What type of bond forms between Lewis acid and base?
 - Ionic
 - Covalent
 - Coordinate covalent
 - Metallic

Answer Key

- Q1: B - Lewis acid accepts electron pair.
- Q2: B - NH_3 donates lone pair, acts as Lewis base.
- Q3: C - Lewis acid-base bond is coordinate covalent.


Memory Aids

- Lewis acid = electron pair acceptor.
- Lewis base = electron pair donor.
- All Lewis bases are also Bronsted bases.

Ionisation of Acids and Bases



Ionisation constant of water and pH scale

- Water ionisation constant: $K_w = [H^+][OH^-] = 1.0 \times 10^{-14}$ at 298 K.
- $pH = -\log[H^+]$; $pOH = -\log[OH^-]$; $pH + pOH = 14$.
- $pH < 7$: acidic; $pH = 7$: neutral; $pH > 7$: basic solution.
- pH scale ranges from 1 (strong acid) to 14 (strong base).




Ionisation of acids and bases

- Acid ionisation: $HA \rightleftharpoons H^+ + A^-$, with $K_a = \frac{[H^+][A^-]}{[HA]}$.
- Degree of ionisation (α) for acid: $\alpha = \sqrt{\frac{K_a}{C}}$, where C is acid concentration.
- Base ionisation: $BOH \rightleftharpoons B^+ + OH^-$, with $K_b = \frac{[B^+][OH^-]}{[BOH]}$.
- Degree of ionisation (α) for base: $\alpha = \sqrt{\frac{K_b}{C}}$.
- Stronger acid/base has higher K_a or K_b and greater α .



Ostwald's dilution law

- For binary electrolyte AB dissociating as $\text{AB} \rightleftharpoons \text{A}^+ + \text{B}^-$:
- Initial concentration: $[\text{AB}] = C$, $[\text{A}^+] = 0$, $[\text{B}^-] = 0$.
- At equilibrium: $[\text{AB}] = C(1 - \alpha)$, $[\text{A}^+] = C\alpha$, $[\text{B}^-] = C\alpha$.
- Dissociation constant: $K_a = \frac{C\alpha^2}{1-\alpha}$.
- For weak electrolytes, $\alpha \ll 1$, so $K_a \approx C\alpha^2$ and $\alpha = \sqrt{\frac{K_a}{C}}$.
- Strong electrolytes: $\alpha = 1$; weak electrolytes: $\alpha < 1$.





Relation between K_a and K_b

- $\text{p}K_a = -\log K_a$; $\text{p}K_b = -\log K_b$.
- $\text{p}K_a + \text{p}K_b = \text{p}K_w = 14$ at 298 K.
- Polybasic acids and polyacidic bases ionise stepwise with constants $K_1, K_2, K_3, \dots$
- Higher order ionisation constants decrease: $K_1 > K_2 > K_3 > \dots$
- Overall ionisation constant: $K = K_1 \times K_2 \times K_3 \times \dots$

Factors affecting acid strength

- Depends on H-A bond strength and polarity.
- Acid strength increases with larger atom size ($\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$).
- Acid strength increases with higher electronegativity of central atom ($\text{CH}_4 < \text{NH}_3 < \text{H}_2\text{O} < \text{HF}$).



Common ion effect

- Adding a common ion suppresses dissociation of a weak electrolyte.

Hydrolysis of salts and pH of their solutions

- Salts ionise in water to hydrated ions; ions react with water (hydrolysis).
- Salts from strong acid + strong base: neutral solution (pH = 7).
- Salts from strong acid + weak base: acidic solution (pH < 7).
- Salts from weak acid + strong base: basic solution (pH > 7).
- Hydrolysis constant (K_h) and degree of hydrolysis (h) depend on K_w , K_a , K_b , and salt concentration.

Salt	Hydrolysis	Resulting solution	Hydrolysis constant (K_h)	Degree of hydrolysis (h)	pH
Weak acid and Strong base	Anionic	Alkaline, pH > 7	$K_h = \frac{K_w}{K_a}$	$h = \sqrt{\frac{K_h}{C}}$	$\text{pH} = \frac{1}{2} [\text{p}K_w + \text{p}K_a + \log C]$
Strong acid and Weak base	Cationic	Acidic, pH < 7	$K_h = \frac{K_w}{K_b}$	$h = \sqrt{\frac{K_h}{C}}$	$\text{pH} = \frac{1}{2} [\text{p}K_w - \text{p}K_b - \log C]$
Weak acid and Weak base	Anionic and Cationic both	Neutral, pH = 7 (If $K_a = K_b$)	$K_h = \frac{K_w}{K_a K_b}$	$h = \sqrt{K_h}$	$\text{pH} = \frac{1}{2} [\text{p}K_w + \text{p}K_a - \text{p}K_b]$

Solved Examples

Ex 1

- Calculate degree of ionisation (α) for a weak acid with $K_a = 1.0 \times 10^{-5}$ and concentration 0.1 M.
- Solution: $\alpha = \sqrt{\frac{K_a}{C}} = \sqrt{\frac{1.0 \times 10^{-5}}{0.1}} = 0.01$.

Ex 2

- Find pH of a solution with $[\text{H}^+] = 1.0 \times 10^{-3}$ M.
- Solution: $\text{pH} = -\log(1.0 \times 10^{-3}) = 3$.

Practice Set

- Q1: What is the pH of pure water at 298 K?

- A. 0
- B. 7
- C. 14
- D. 1
- Q2: For a weak acid, which expression gives the degree of ionisation (α)?
 - A. $\alpha = K_a \times C$
 - B. $\alpha = \sqrt{\frac{K_a}{C}}$
 - C. $\alpha = K_a / C$
 - D. $\alpha = C / K_a$
- Q3: If $pK_a = 5$, what is pK_b for its conjugate base?
 - A. 9
 - B. 14
 - C. 5
 - D. 19

Answer Key

- Q1: B - Pure water is neutral with pH 7 at 298 K.
- Q2: B - Degree of ionisation for weak acid is $\sqrt{\frac{K_a}{C}}$.
- Q3: A - $pK_a + pK_b = 14$, so $pK_b = 14 - 5 = 9$.

Memory Aids

- $pH + pOH = 14$ always at 298 K.
- Stronger acid: higher K_a , lower pK_a .
- Ostwald's law: $\alpha = \sqrt{\frac{K_a}{C}}$ for weak electrolytes.
- Acid strength increases down halogen group ($HF < HCl < HBr < HI$).
- Common ion effect reduces ionisation of weak electrolytes.

Buffer Solutions

Definition and Classification

- Solutions that resist pH change on dilution or addition of small amounts of acid/base.
- Two types of buffers:
 - **Simple buffers:** Solutions of salts of weak acid and weak base (e.g., CH_3COONH_4).
 - **Mixed buffers:** Mixtures of two solutions, further divided into:
 - **Acidic buffers:** Weak acid + salt of weak acid with strong base (e.g., $CH_3COOH + CH_3COONa$), $pH < 7$.
 - **Basic buffers:** Weak base + salt of weak base with strong acid (e.g., $NH_4OH + NH_4Cl$), $pH > 7$.

Solved Examples

- **Ex 1:** Identify the type of buffer formed by mixing CH_3COOH and CH_3COONa .
 - Mixture of weak acid and its salt with strong base.
 - Hence, acidic buffer with $\text{pH} < 7$.

Practice Set

- Q1: Which of the following is a simple buffer?
 - A. $\text{CH}_3\text{COOH} + \text{CH}_3\text{COONa}$
 - B. $\text{NH}_4\text{OH} + \text{NH}_4\text{Cl}$
 - C. $\text{CH}_3\text{COONH}_4$
 - D. $\text{HCl} + \text{NaCl}$
- Q2: What is the pH range of acidic buffers?
 - A. $\text{pK}_a \pm 1$
 - B. $(\text{pK}_w - \text{pK}_b) \pm 1$
 - C. 7 ± 1
 - D. 14 ± 1
- Q3: Which mixture forms a basic buffer?
 - A. $\text{CH}_3\text{COOH} + \text{CH}_3\text{COONa}$
 - B. $\text{NH}_4\text{OH} + \text{NH}_4\text{Cl}$
 - C. $\text{HCl} + \text{NaCl}$
 - D. $\text{CH}_3\text{COONH}_4$

Answer Key

- Q1: C - Simple buffer is salt of weak acid and weak base (ammonium acetate).
- Q2: A - Acidic buffer range is $\text{pK}_a \pm 1$.
- Q3: B - Basic buffer formed by weak base and its salt with strong acid ($\text{NH}_4\text{OH} + \text{NH}_4\text{Cl}$).

Memory Aids

- Simple buffer = salt of weak acid + weak base (e.g., $\text{CH}_3\text{COONH}_4$).
- Acidic buffer = weak acid + salt of weak acid with strong base ($\text{pH} < 7$).
- Basic buffer = weak base + salt of weak base with strong acid ($\text{pH} > 7$).

pH of Buffer Solutions

Acidic Buffer pH

- $\text{pH} = \text{pK}_a + \log ([\text{salt}]/[\text{acid}])$
- Also written as Henderson-Hasselbalch equation: $\text{pH} = \text{pK}_a + \log ([\text{conjugate base}]/[\text{acid}])$

Basic Buffer pH

- $\text{pOH} = \text{pK}_b + \log ([\text{salt}]/[\text{base}])$
- Also: $\text{pOH} = \text{pK}_b + \log ([\text{conjugate acid}]/[\text{base}])$

Solved Examples

- **Ex 1:** Calculate pH of acidic buffer with $\text{pK}_a = 4.76$, $[\text{salt}] = 0.1 \text{ M}$, $[\text{acid}] = 0.1 \text{ M}$.
 - $\text{pH} = 4.76 + \log(0.1/0.1) = 4.76 + 0 = 4.76$
- **Ex 2:** Calculate pOH of basic buffer with $\text{pK}_b = 4.75$, $[\text{salt}] = 0.2 \text{ M}$, $[\text{base}] = 0.1 \text{ M}$.
 - $\text{pOH} = 4.75 + \log(0.2/0.1) = 4.75 + 0.301 = 5.05$

Practice Set

- Q1: For acidic buffer, pH is given by?
 - A. $\text{pK}_a + \log ([\text{acid}]/[\text{salt}])$
 - B. $\text{pK}_a + \log ([\text{salt}]/[\text{acid}])$
 - C. $\text{pK}_b + \log ([\text{salt}]/[\text{base}])$
 - D. $\text{pK}_b + \log ([\text{base}]/[\text{salt}])$
- Q2: If $\text{pK}_b = 4.75$, $[\text{salt}] = 0.1 \text{ M}$, $[\text{base}] = 0.1 \text{ M}$, pOH of basic buffer is?
 - A. 4.75
 - B. 5.05
 - C. 4.45
 - D. 3.75
- Q3: Calculate pH of acidic buffer with $\text{pK}_a = 3.75$, $[\text{salt}] = 0.05 \text{ M}$, $[\text{acid}] = 0.1 \text{ M}$.
 - A. 3.45
 - B. 3.75
 - C. 4.05
 - D. 3.25

Answer Key

- Q1: B - Henderson-Hasselbalch equation for acidic buffer.
- Q2: A - $\text{pOH} = 4.75 + \log(0.1/0.1) = 4.75$.
- Q3: A - $\text{pH} = 3.75 + \log(0.05/0.1) = 3.75 - 0.3 = 3.45$.

Memory Aids

- Acidic buffer pH uses pK_a and ratio of salt to acid.
- Basic buffer pOH uses pK_b and ratio of salt to base.
- Remember: $\text{pH} + \text{pOH} = 14$.

Buffer Capacity and Range

Buffer Capacity

- Buffer capacity = moles of acid/base added per litre of buffer $\div$ change in pH.
- Indicates buffer's ability to resist pH change.

Buffer Range

- Range of pH over which buffer is effective.
- For acidic buffers: $\text{pK}_a \pm 1$.
- For basic buffers: $(\text{pK}_w - \text{pK}_b) \pm 1$.

Solved Examples

- **Ex 1:** Calculate buffer capacity if 0.01 moles acid added to 1 L buffer causing pH change from 5.0 to 4.8.
 - Change in pH = 0.2
 - Buffer capacity = $0.01 / 0.2 = 0.05 \text{ mol/L per pH unit}$.

Practice Set

- Q1: Buffer capacity measures:
 - A. pH of buffer
 - B. Amount of acid/base buffer can neutralize per pH change
 - C. pK_a value
 - D. Concentration of buffer components

- Q2: Buffer range of acidic buffer is:
 - A. $\text{pK}_a \pm 1$
 - B. $(\text{pK}_w - \text{pK}_b) \pm 1$
 - C. 7 ± 1
 - D. 14 ± 1
- Q3: If buffer capacity is 0.1 mol/L per pH unit, moles of acid added causing 0.5 pH change is:
 - A. 0.05
 - B. 0.2
 - C. 0.1
 - D. 0.5

Answer Key

- Q1: B - Buffer capacity shows resistance to pH change by neutralizing added acid/base.
- Q2: A - Acidic buffer effective range is $\text{pK}_a \pm 1$.
- Q3: B - Moles = capacity $\times$ pH change = $0.1 \times 0.5 = 0.05$ mol.

Memory Aids

- Buffer capacity = moles acid/base added $\div$ pH change.
- Acidic buffer range near pK_a , basic near $(\text{pK}_w - \text{pK}_b)$.

Solubility vs Solubility Product

Definitions and Differences

- **Solubility:** Amount of solute dissolved in 1 L of solvent (usually in grams).
- **Solubility Product (K_{sp}):** Equilibrium constant for a saturated solution showing ion concentrations (in mol/L) at a given temperature.
- Solubility depends on temperature, pressure, pH, and other chemicals.
- K_{sp} depends only on temperature.

Solved Examples

- **Ex 1:** Calculate solubility if $K_{sp} = 1.6 \times 10^{-10}$ for AgCl (1:1 salt).
 - Solubility, $s = \sqrt{K_{sp}} = \sqrt{(1.6 \times 10^{-10})} = 1.26 \times 10^{-5}$ mol/L

Practice Set

- Q1: What does solubility measure?
 - A) Ion concentration at equilibrium
 - B) Amount of solute dissolved per liter
 - C) Temperature effect on salt
 - D) Pressure effect on salt
- Q2: Which factor affects K_{sp} ?
 - A) pH
 - B) Temperature
 - C) Pressure
 - D) Common ion presence
- Q3: If solubility of salt is 2×10^{-3} mol/L, what is K_{sp} for a 1:1 salt?
 - A) 4×10^{-6}
 - B) 2×10^{-3}
 - C) 2×10^{-6}
 - D) 4×10^{-3}

Answer Key

- Q1: B - Solubility is the amount of solute dissolved per liter of solvent.
- Q2: B - K_{sp} depends only on temperature.
- Q3: A - $K_{sp} = s^2 = (2 \times 10^{-3})^2 = 4 \times 10^{-6}$.

Memory Aids

- Solubility = grams per liter; K_{sp} = ion product constant.
- K_{sp} changes only with temperature.
- Solubility affected by common ions, pH, pressure.

Solubility Product Constant

Equilibrium and Expression

- For salt M_xX_y : $M_xX_y(s) \rightleftharpoons xM^{y+}(aq) + yX^{x-}(aq)$
- $K_{sp} = [M^{y+}]^x [X^{x-}]^y = (xs)^x (ys)^y = x^x y^y s^{x+y}$
- Solubility, $s = \left(\frac{K_{sp}}{x^x y^y} \right)^{\frac{1}{x+y}}$
- At equilibrium, $K_{sp} = Q_{sp}$. If not, Q_{sp} indicates reaction direction.

Effect of Common Ion

- Adding a common ion decreases solubility (common ion effect).
- Example: For AgCl, if NaCl (common Cl^-) is added, solubility decreases.
- $K_{\text{sp}} = [\text{Ag}^+][\text{Cl}^-] = S'(S' + x)$, where x is added ion concentration.
- Increase in solubility on adding inert salt indicates complex ion formation.

Solved Examples

- **Ex 1:** Calculate solubility of AgCl if $K_{\text{sp}} = 1.8 \times 10^{-10}$ and 0.01 M NaCl is added.
 - $K_{\text{sp}} = S'(S' + 0.01) \approx S' \times 0.01$ (if $S' \ll 0.01$)
 - $S' = K_{\text{sp}} / 0.01 = 1.8 \times 10^{-8} \text{ mol/L}$ (much less than pure water solubility)

Practice Set

- Q1: What happens to solubility when a common ion is added?
 - A) Increases
 - B) Decreases
 - C) No change
 - D) Depends on temperature
- Q2: For salt M_xX_y , K_{sp} expression is?
 - A) $[\text{M}^{y+}][\text{X}^{x-}]$
 - B) $[\text{M}^{y+}]^x[\text{X}^{x-}]^y$
 - C) $(xs)(ys)$
 - D) $(x+y)s$
- Q3: If $Q_{\text{sp}} > K_{\text{sp}}$, what occurs?
 - A) No precipitation
 - B) Precipitation
 - C) Equilibrium
 - D) Complex formation

Answer Key

- Q1: B - Common ion decreases solubility by shifting equilibrium.
- Q2: B - $K_{\text{sp}} = [\text{M}^{y+}]^x[\text{X}^{x-}]^y$.
- Q3: B - Ionic product greater than K_{sp} causes precipitation.

Memory Aids

- Common ion effect lowers solubility.
- K_{sp} expression includes ion concentrations raised to their stoichiometric powers.
- $Q_{sp} > K_{sp}$ means precipitation.

Applications of Solubility Product

Key Uses

- Calculate solubility of sparingly soluble salts from K_{sp} .
- Predict precipitation by comparing ionic product (Q) with K_{sp} .
- Explain precipitation of common salt by adding HCl gas (increases Cl^- concentration).
- Use in qualitative analysis to separate ions based on solubility and common ion effect.

Solved Examples

- **Ex 1:** Predict if precipitation occurs when $[Ag^+] = 1 \times 10^{-4} \text{ M}$ and $[Cl^-] = 1 \times 10^{-3} \text{ M}$, K_{sp} of AgCl = 1.8×10^{-10} .
 - Ionic product = $(1 \times 10^{-4})(1 \times 10^{-3}) = 1 \times 10^{-7} > K_{sp}$
 - Precipitation occurs.

Practice Set

- Q1: What indicates precipitation?
 - A) Ionic product $< K_{sp}$
 - B) Ionic product $= K_{sp}$
 - C) Ionic product $> K_{sp}$
 - D) No ions present
- Q2: What happens when HCl gas is passed through impure NaCl solution?
 - A) NaCl dissolves more
 - B) Pure NaCl precipitates
 - C) No change
 - D) NaCl decomposes
- Q3: Which application uses solubility product in qualitative analysis?
 - A) Identifying gases
 - B) Separating basic radicals
 - C) Measuring pH
 - D) Calculating molarity

Answer Key

- Q1: C - Precipitation occurs if ionic product $> K_{sp}$.
- Q2: B - Increased Cl^- causes pure NaCl to precipitate.
- Q3: B - Solubility product helps separate basic radicals.

Memory Aids

- Precipitation if $Q > K_{sp}$.
- HCl increases Cl^- , causing NaCl precipitation.
- Qualitative analysis uses solubility and common ion effect.

Solubility Products of Electrolytes

Classification and Formulas

- Salts classified by electrolyte formula: binary (1:1), ternary (1:2 or 2:1), quaternary (1:3), 3:2 types.
- Examples:
 - Binary (1:1): AgCl
 - Ternary (1:2): $PbCl_2$
 - Quaternary (1:3): $Fe(OH)_3$
- Solubility (s) related to K_{sp} differently for each type:
 - 1:1 salts: $s = \sqrt{K_{sp}}$
 - 1:2 salts: $s = \sqrt[3]{(K_{sp}/4)}$
 - 2:1 salts: $s = \sqrt[3]{(K_{sp}/4)}$
 - 1:3 salts: $s = \sqrt[4]{\frac{K_{sp}}{27}}$
 - 3:2 salts: $s = \sqrt[5]{\frac{K_{sp}}{108}}$

Type of salt	Example	Relation between s and K_{sp}
Binary electrolyte 1 : 1 (AB)	AgCl, $BaSO_4$, etc.	$s = \sqrt{K_{sp}}$ or $K_{sp} = s^2$
Ternary electrolyte 1 : 2 (AB_2)	$PbCl_2$, $Ca(OH)_2$, etc.	$s = \sqrt[3]{K_{sp}/4}$ or $K_{sp} = 4s^3$
Ternary electrolyte 2 : 1 (A_2B)	Ag_2CrO_4 , Ag_2CO_3 , etc.	$s = \sqrt[3]{K_{sp}/4}$ or $K_{sp} = 4s^3$
Quaternary electrolyte 1 : 3 (AB_3)	$Fe(OH)_3$, $Al(OH)_3$, etc.	$s = \sqrt[4]{K_{sp}/27}$ or $K_{sp} = 27s^4$
3 : 2 type of electrolyte (A_3B_2)	$Ca_3(PO_4)_2$, $Ba_3(PO_4)_2$ etc.	$s = \sqrt[5]{K_{sp}/108}$ or $K_{sp} = 108s^5$

Solved Examples

- **Ex 1:** Calculate solubility of PbCl_2 if $K_{\text{sp}} = 1.7 \times 10^{-5}$.
 - For 1:2 salt, $s = \sqrt[3]{(K_{\text{sp}}/4)} = \sqrt[3]{(1.7 \times 10^{-5}/4)} = 0.016 \text{ mol/L}$

Practice Set

- Q1: Which salt is binary 1:1?
 - A) PbCl_2
 - B) AgCl
 - C) $\text{Fe}(\text{OH})_3$
 - D) $\text{Al}(\text{OH})_3$
- Q2: Solubility formula for 1:2 salt is?
 - A) $s = \sqrt{K_{\text{sp}}}$
 - B) $s = \sqrt[3]{(K_{\text{sp}}/4)}$
 - C) $s = K_{\text{sp}}^2$
 - D) $s = K_{\text{sp}}/4$
- Q3: For $\text{Fe}(\text{OH})_3$ (1:3 salt), solubility is?
 - A) $\sqrt[4]{\frac{K_{\text{sp}}}{27}}$
 - B) $\sqrt[3]{K_{\text{sp}}}$
 - C) $\sqrt{K_{\text{sp}}}$
 - D) $\sqrt[5]{\frac{K_{\text{sp}}}{108}}$

Answer Key

- Q1: B - AgCl is binary 1:1 salt.
- Q2: B - For 1:2 salts, $s = \sqrt[3]{(K_{\text{sp}}/4)}$.
- Q3: A - For 1:3 salts, $s = \sqrt[4]{\frac{K_{\text{sp}}}{27}}$.

Memory Aids

- Binary (1:1): $s = \sqrt{K_{\text{sp}}}$
- Ternary (1:2 or 2:1): $s = \sqrt[3]{(K_{\text{sp}}/4)}$
- Quaternary (1:3): $s = \text{4th root of } (K_{\text{sp}}/27)$
- 3:2 salts: 5th root of $(K_{\text{sp}}/108)$