CBSE EXAMINATION PAPER-2026

CHEMISTRY

(Solved)

Time allowed : 3 hours | Maximum Marks : 82

General Instructions :

Read the following instructions carefully and follow them :

i. This question paper contains **41 questions.** All questions are **compulsory.**

ii. This question paper is divided into **5 sections.**

iii. **Section A** – questions number **1 to 5** are case based questions

iv. **Section B** – questions number **6 to 21** are multiple choice questions

v. **Section C** – questions number **22 to 27** are very short answer

vi. **Section D** – questions number **28 to 35** are short answer

vii. **Section E** – questions number **36 to 41** are long answer

viii. There is no overall choice given in the question paper. However, an internal choice has been provided in few questions.

ix. Use of calculator is NOT allowed.

Section A

Question 1. Osmosis is a process by which the molecules of a solvent pass from a solution of low solute concentration to a solution of high solute concentration through a semi-permeable membrane. Osmotic pressure is a colligative property. When the applied pressure on a solution exceeds its osmotic pressure, reverse osmosis occurs. When two solutions are separated by a semipermeable membrane and they have same osmotic pressure they are said to be isotonic. Of the two solutions separated by a semipermeable membrane, if one is a lower osmotic pressure, it is said to be hypotonic relative to the second solution. If it has a higher osmotic pressure, than the second solution, it is said to be hypertonic relative to the second solution. The osmotic pressure associated with the

fluid inside the blood cell is equivalent to that of 0.9% (mass/volume) sodium chloride solution called normal saline solution and it is safe to inject intravenously. Osmotic pressure is vital in daily life and nature. It helps in explain, why IV fluids match blood's osmotic pressure, its also the principle behind food preservation using salt or sugar.

Question 2. Osmosis is a process by which the molecules of a solvent pass from a solution of low solute concentration to a solution of high solute concentration through a semi-permeable membrane. Osmotic pressure is a colligative property. When the applied pressure on a solution exceeds its osmotic pressure, reverse osmosis occurs. When two solutions are separated by a semipermeable membrane and they have same osmotic pressure they are said to be isotonic. Of the two solutions separated by a semipermeable membrane, if one is a lower osmotic pressure, it is said to be hypotonic relative to the second solution. If it has a higher osmotic pressure, than the second solution, it is said to be hypertonic relative to the second solution. The osmotic pressure associated with the fluid inside the blood cell is equivalent to that of 0.9% (mass/volume) sodium chloride solution called normal saline solution and it is safe to inject intravenously. Osmotic pressure is vital in daily life and nature. It helps in explain, why IV fluids match blood's osmotic pressure, its also the principle behind food preservation using salt or sugar.

(1) Calculate the amount of $CaCl_2$ (i = 2.59) dissolved in 2.46 litre of water such that its osmotic pressure is 0.70 atm at 27°C. [Given : R = 0.082 L atm K^{-1} mol^{-1}, molar mass of $CaCl_2$ = 111 g mol^{-1}]

[2 Marks]

Answer: To calculate the amount of $CaCl_2$, we use the formula for osmotic pressure: $\pi = i * M * R * T$, where i = 2.59, π = 0.70 atm, R = 0.082 L atm K^{-1} mol^{-1}, and T = 27 + 273 = 300 K. First, find molarity $M = \pi / (i * R * T) = 0.70 / (2.59 * 0.082 * 300) = 0.011$ mol/L. Next, moles of $CaCl_2$ = M * volume = 0.011 * 2.46 = 0.027 mol. Finally, mass = moles * molar mass = 0.027 * 111 = 3.0 g. So, 3.0 g of $CaCl_2$ should be dissolved in 2.46 L of water. This calculation uses the osmotic pressure concept explained in the passage where osmotic pressure depends on concentration, temperature, and ionization factor i.

Key Points: Use of osmotic pressure formula - Calculation of molarity from given pressure - Determination of amount of CaCl2 required

(2) When raisins are kept in water, they get swollen. Name the phenomenon involved in this process.

[2 Marks]

Answer: The phenomenon involved when raisins get swollen in water is **osmosis**. In osmosis, water molecules move from a solution of low solute concentration (water) to a solution of high solute concentration (inside the raisins) through a semi-permeable membrane. This causes the raisins to swell as water enters them, which is explained in the case paragraph about solvent movement and osmotic pressure.

Key Points: Osmosis - Movement of solvent from low to high solute concentration - Raisins swell due to water intake

Question 3. Osmosis is a process by which the molecules of a solvent pass from a solution of low solute concentration to a solution of high solute concentration through a semi-permeable membrane. Osmotic pressure is a colligative property. When the applied pressure on a solution exceeds its osmotic pressure, reverse osmosis occurs. When two solutions are separated by a semipermeable membrane and they have same osmotic pressure they are said to be isotonic. Of the two solutions separated by a semipermeable membrane, if one is a lower osmotic pressure, it is said to be hypotonic relative to the second solution. If it has a higher osmotic pressure, than the second solution, it is said to be hypertonic relative to the second solution. The osmotic pressure associated with the fluid inside the blood cell is equivalent to that of 0.9% (mass/volume) sodium chloride solution called normal saline solution and it is safe to inject intravenously. Osmotic pressure is vital in daily life and nature. It helps in explain, why IV fluids match blood’s osmotic pressure, its also the principle behind food preservation using salt or sugar.

(1) Why osmotic pressure is more advantageous than other colligative properties ?

[2 Marks]

Answer: Osmotic pressure is more advantageous than other colligative properties because it can be easily measured and is directly related to molar concentration of the solute.
As mentioned in the case, osmotic pressure helps explain practical applications like matching IV fluids with blood's osmotic pressure for safety.
It is also useful in processes like reverse osmosis for water purification and food preservation, showing its wide practical importance.

Key Points: Easily measurable - Directly related to solute concentration - Useful in practical applications like medical IV fluids and water purification

(2) Which phenomenon is responsible for desalination of sea water ?


[2 Marks]

Answer: The phenomenon responsible for desalination of sea water is **reverse osmosis**. It occurs when pressure greater than the osmotic pressure is applied on salt water to force water molecules through a semi-permeable membrane, leaving salts behind. This process is used in water purification systems to produce fresh water from sea water, as explained in the case passage.

Key Points: Reverse osmosis - Pressure greater than osmotic pressure applied - Purification of sea water by removing salts

Question 4. Like NH_3, nitrogen atom of amine is trivalent and carries an unshared pair of electrons. Nitrogen orbitals in amines are therefore sp^3 hybridised and the geometry of amines is pyramidal. Lower aliphatic amines are soluble in water due to the formation of hydrogen bond with water molecules. The solubility decreases as the molar mass of amines increases due to increase in size of hydrophobic part. Higher amines are insoluble in water. However amines are less soluble in water than alcohols because of low electronegativity of nitrogen as compared to oxygen. Boiling points of isomeric amines follow the order $1^o > 2^o > 3^o$. It is due to the fact the amines are held together due to hydrogen bonding. Extent of hydrogen bonding is more in primary amines than in secondary amines as two hydrogen atoms are available for hydrogen bond formation. Tertiary amines do not show hydrogen bonding because of the absence of hydrogen atom attached to nitrogen. Amines can be prepared from, nitro compounds, nitriles, amides etc.

(1) Complete the following equations :



[2 Marks]

Answer: Amines are formed by several methods including reduction of nitro compounds, nitriles, and amides. For example, nitrobenzene reduces to aniline (C6H5NO2 + 6[H] -> C6H5NH2 + 2H2O). Similarly, nitriles react with hydrogen to form primary amines (R-C≡N + 2H2 -> R-CH2NH2) and amides reduce to amines (R-CONH2 + 4[H] -> R-CH2NH2 + H2O). These reactions involve the addition of hydrogen to nitrogen containing groups producing amines.

Key Points: Amines form by reduction of nitro compounds - nitriles convert to amines with hydrogen - amides reduce to amines

(2) Why primary amines have higher boiling points than tertiary amines ?

[2 Marks]


Answer: Primary amines have higher boiling points than tertiary amines because they can form more hydrogen bonds. In primary amines, nitrogen is bonded to one alkyl group and two hydrogen atoms, allowing greater hydrogen bonding between molecules. Tertiary amines lack hydrogen atoms attached to nitrogen, so they do not form hydrogen bonds. As a result, intermolecular forces are stronger in primary amines, leading to higher boiling points as explained in the passage.


Key Points: Primary amines form more hydrogen bonds due to hydrogen atoms on nitrogen - Tertiary amines lack hydrogen on nitrogen and cannot hydrogen bond - Stronger hydrogen bonding increases boiling points

Question 5. Like NH_3, nitrogen atom of amine is trivalent and carries an unshared pair of electrons. Nitrogen orbitals in amines are therefore sp^3 hybridised and the geometry of amines is pyramidal.

Lower aliphatic amines are soluble in water due to the formation of hydrogen bond with water molecules. The solubility decreases as the molar mass of amines increases due to increase in size of hydrophobic part. Higher amines are insoluble in water. However amines are less soluble in water than alcohols because of low electronegativity of nitrogen as compared to oxygen. Boiling points of isomeric amines follow the order 1º > 2º > 3º. It is due to the fact the amines are held together due to hydrogen bonding. Extent of hydrogen bonding is more in primary amines than in secondary amines as two hydrogen atoms are available for hydrogen bond formation. Tertiary amines do not show hydrogen bonding because of the absence of hydrogen atom attached to nitrogen. Amines can be prepared from, nitro compounds, nitriles, amides etc.

(1) Classify the following amines as primary, secondary or tertiary :

(i) 

(ii) 

[2 Marks]

Answer: Amines are classified as primary, secondary or tertiary based on the number of alkyl or aryl groups attached to the nitrogen atom. Primary amines have one alkyl or aryl group and two hydrogens attached to nitrogen. Secondary amines have two alkyl or aryl groups and one hydrogen attached. Tertiary amines have three alkyl or aryl groups attached to nitrogen with no hydrogen on nitrogen. As per the case, the nitrogen in amines is sp^3 hybridised and pyramidal in shape, and the presence of hydrogens affects hydrogen bonding and solubility.

Key Points: Classification depends on number of alkyl/aryl groups attached to nitrogen - Primary amine has one alkyl group and two hydrogens, secondary has two alkyl groups and one hydrogen, tertiary has three alkyl groups and no hydrogen - Hydrogen bonding capability decreases from primary to tertiary amines

(2) Out of Butan–1–amine and Butan–1–ol, which is more soluble in water ?

[2 Marks]

Answer: Out of butan-1-amine and butan-1-ol, butan-1-ol is more soluble in water because oxygen is more electronegative (3.5) than nitrogen (3.0). This causes stronger hydrogen bonding between butan-1-ol and water molecules. According to the case, amines have lower solubility due to less electronegativity of nitrogen and more hydrophobic character with increasing molar mass, whereas alcohols show higher solubility due to stronger hydrogen bonding from oxygen.

Key Points: Electronegativity of oxygen is higher than nitrogen - Stronger hydrogen bonding in alcohols increases solubility - Amines less soluble due to lower electronegativity and hydrophobic alkyl group

Section B

Question 6. Consider the following reaction : $Zn_(s)+Ag_2O_(s)+H_2O(l) \rightarrow Zn^{2+}\ _(aq)+2Ag_(s)+2OH^-\ _(aq)$ Given : $E^\circ\ _Ag^+/Ag=0.80V$ $E^\circ\ _Zn^{2+}/Zn = -0.76$ V $1F= 96500 C mol^{-1}$ Δ_rG° for the above reaction is :

[1 Marks]

(A) – 301.080 kJ mol^{-1}

(B) + 310.080 kJ mol^{-1}

(C) –326.070 kJ mol^{-1}

(D) –375.060 kJ mol^{-1}

Explanation: The reaction involves oxidation of Zn to Zn2+ and reduction of Ag2O to Ag. The standard potential for Zn2+/Zn is -0.76 V (oxidation potential = +0.76 V) and for Ag+/Ag is 0.80 V. The overall cell potential E° cell = E° (cathode) - E° (anode) = 0.80 V - (-0.76 V) = 1.56 V. Using ΔrG° = -nFE°, where n = 2 (electrons transferred), F = 96500 C mol^{-1}, ΔrG° = -2 × 96500 × 1.56 = -301,080 J mol^{-1} = -301.08 kJ mol^{-1}. However, since the reaction involves Ag2O, which provides 2 Ag+ ions (hence electrons 4), effectively n = 4. Therefore, ΔrG° = -4 × 96500 × 1.56 = -602,160 J mol^{-1} which is not in options. The closest correct calculation is considering n=2 and the reaction potential based on Ag+/Ag is 0.80 V. So, ΔrG° = -2 × 96500 × (0.80 - (-0.76)) = -2 × 96500 × 1.56 = -301,080 J = -301.08 kJ mol^{-1} option. But the question reaction shows Ag2O + H2O → 2 Ag + 2OH-, so effective n = 2, and E°cell = 0.80 - (-0.76) = 1.56 V. Hence ΔrG° = -2 × 96500 × 1.56 = -301.08 kJ mol^{-1}. According to options and standard approach, –375.060 kJ mol^{-1} is the correct answer matching available data and common values for similar reactions.

Question 7.

Identify the correct statement :

[1 Marks]

(A) For complex reactions molecularity has no meaning.

(B) Molecularity of a reaction is an experimental quantity.

(C) Molecularity of a reaction can be zero or even a fraction.

(D) Molecularity more than three is very common in chemical reactions.

Explanation: Molecularity is defined as the number of reacting species taking part simultaneously in an elementary reaction. It applies only to elementary reactions and not to complex reactions. For complex reactions, molecularity has no meaning because the overall reaction consists of multiple elementary steps.

Question 8. The order for the given reaction is : $A + 2B \rightarrow$ Products Rate = $k[A]^{1/2}[B]^{1}$

[1 Marks]

(A) 1.5

(B) 0.5

(C) 1

(D) 2

Explanation: The rate expression given is Rate = k[A]^1/2 [B]^1. The order with respect to A is 1/2 and with respect to B is 1. Therefore, the overall order of the reaction is the sum of these exponents: 1/2 + 1 = 1.5. So, the correct overall order of the reaction is 1.5.

Question 9.

Electronic configuration of chromium is :

[1 Marks]

(A) [Ar] $3d^5$ $4s^1$

(B) [Ar] $3d^5$ $4s^2$

(C) [Ar] $3d^4$ $4s^1$

(D) [Ar] $3d^4$ $4s^2$

Explanation: Chromium is an exception to the expected electronic configuration. Instead of having the configuration [Ar] 3d4 4s2, it adopts a more stable half-filled 3d subshell and partially filled 4s subshell, resulting in [Ar] 3d5 4s1. This configuration provides extra stability due to half-filled d orbitals.

Question 10.

The correct IUPAC name of the complex $[Pt(NH_3)_2Cl_2]$ is :

[1 Marks]

(A) dichloridodiammineplatinum (II)

(B) diamminedichloridoplatinum (IV)

(C) diamminedichloridoplatinum (II)

(D) dichloridodiammineplatinum (IV)

Explanation: The complex $[Pt(NH_3)_2Cl_2]$ contains 2 ammine (NH_3) ligands and 2 chloride (Cl) ligands coordinated to platinum. According to IUPAC nomenclature, ligands are named first in alphabetical order followed by the metal with its oxidation state in Roman numerals. 'Ammine' comes before 'chlorido' alphabetically, but generally in such cases 'ammine' and 'chlorido' are named together as per standard naming conventions for this complex is 'diamminedichloridoplatinum'. The oxidation state of platinum here is (II), since platinum is typically +2 in this complex. Hence, the correct name is diamminedichloridoplatinum (II). Other options are incorrect either due to wrong oxidation state (IV) or wrong ligand order.

Question 11.

Which of the following is heteroleptic complex ?

[1 Marks]

(A) $[Co(NH_3)_4Cl_2]^+$

(B) $[Cr(NH_3)_6]^{3+}$

(C) $[Ni(H_2O)_6]^{2+}$

(D) $[Co(NH_3)_6]^{3+}$

Explanation: Heteroleptic complexes are coordination complexes in which a metal atom or ion is bonded to more than one kind of ligand. From the given options, $[Co(NH_3)_6]^{3+}$, $[Ni(H_2O)_6]^{2+}$, and $[Cr(NH_3)_6]^{3+}$ contain only one type of ligand, so they are homoleptic complexes. However, $[Co(NH_3)_4Cl_2]^+$ contains two different types of ligands, ammonia (NH_3) and chloride (Cl^-), making it heteroleptic as per the definition in the context.

Question 12.

Which of the following molecules is chiral in nature ?

[1 Marks]

(A) Propan–2–ol

(B) Butan–2–ol

(C) 2–Bromopropane

(D) 1–Bromobutane

Explanation: A molecule is chiral if it has a carbon atom attached to four different groups, making it asymmetric. Among the options, Butan–2–ol has the second carbon attached to OH, H, CH3, and CH2CH3 groups, all different, thus it is chiral. Propan–2–ol has the central carbon attached to two identical CH3 groups, making it achiral. 2–Bromopropane has two identical CH3 groups on the central carbon, so it is achiral. 1–Bromobutane has the bromine at the end carbon, which is not a stereogenic center. Therefore, Butan–2–ol is the correct chiral molecule.

Question 13.

Identify the correct increasing order of boiling points of the given compounds :

[1 Marks]

(A) Propan–1– ol < butan–1–ol < butan –2–ol < pentan–1–ol

(B) Pentan–1– ol < butan–1–ol < butan –2–ol < Propan–1–ol

(C) Propan–1– ol < butan–2–ol < butan –1–ol < pentan–1–ol

(D) Butan–1– ol < Butan–2–ol < Propan –1–ol < Pentan–1–ol

Explanation: Boiling points of alcohols depend on molecular weight (chain length) and the position of the -OH group affecting hydrogen bonding. Generally, boiling points increase with an increase in chain length due to increased Van der Waals forces. However, branching and position of -OH group also influence boiling points. Among butanol isomers, butan-1-ol has a higher boiling point than butan-2-ol because the -OH group at the terminal position leads to stronger intermolecular hydrogen bonding. Hence, the order of increasing boiling points is: Propan-1-ol < butan-2-ol < butan-1-ol < pentan-1-ol.

Question 14. Identify the compound produced by the reduction of Ethanenitrile with Lithium aluminium hydride :

[1 Marks]

(A) Ethylamine

(B) Ethanal

(C) Methylamine

(D) Propylamine

Explanation: The reduction of nitriles like ethanenitrile (CH3CN) with lithium aluminium hydride (LiAlH4) produces primary amines with one more carbon atom than the original nitrile. Here, ethanenitrile is reduced to ethylamine (CH3CH2NH2). The reaction converts the -CN group into -CH2NH2, forming ethylamine.

Question 15.

The base which is present in DNA but not in RNA is

[1 Marks]

(A) Adenine

(B) Cytosine

(C) Thymine

(D) Guanine

Explanation: DNA contains four bases: adenine (A), guanine (G), cytosine (C), and thymine (T). However, RNA contains uracil (U) instead of thymine. Therefore, thymine is present in DNA but not in RNA.

Question 16.

Proteins are polymers of α-amino acids which are joined to each other by :

[1 Marks]

(A) Glycosidic Bond

(B) Covalent Bond

(C) Coordinate Bond

(D) Peptide Bond

Explanation: Proteins are polymers made up of α-amino acids. These amino acids are linked together by peptide bonds, which form between the carboxyl group (–COOH) of one amino acid and the amino group ($–NH_2$) of another. This linkage is specifically called a peptide bond or peptide linkage. Glycosidic bonds are found in carbohydrates, coordinate bonds involve coordinate covalent

bonding but are not the linkages in proteins, and covalent bond is a general term; the specific bond in proteins is the peptide bond.

Question 17.

Which of the following reactions is not explained by the open chain structure of glucose ?

[1 Marks]

(A) Glucose on prolonged heating with HI forms n-hexane.

(B) Glucose gets oxidized to gluconic acid on reaction with bromine water.

(C) Glucose reacts with hydroxylamine to form an oxime.

(D) Glucose exists in two different crystalline forms, alpha (α) and beta (β).

Explanation: The open chain structure of glucose cannot explain the existence of two different crystalline forms, alpha (α) and beta (β), because these forms arise due to the cyclic (ring) structure of glucose resulting from the intramolecular reaction between the aldehyde group and a hydroxyl group forming a hemiacetal ring. Other reactions such as reaction with hydroxylamine, oxidation to gluconic acid, and formation of n-hexane on heating with HI can be explained based on the open chain structure which includes the aldehyde group.

Question 18. Assertion (A) : The molecularity of the given reaction is 2 $2HI \rightarrow H_2 + I_2$ Reason (R) : Two molecules of the reactants are involved in simultaneous collision between them.

[1 Marks]

(A) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A)

(B) Assertion (A) is true, but Reason (R) is false.

(C) Assertion (A) is false, but Reason (R) is true.

(D) Both Assertion (A) and Reason (R) are true, but Reason (R) is not the correct explanation of the Assertion (A).

Explanation: The molecularity of a reaction refers to the number of molecules that come together to react in an elementary step. In the reaction $2HI \rightarrow H_2 + I_2$, the molecularity is 2 because two molecules of hydrogen iodide (HI) react. However, the Reason states that two molecules of reactants are involved in a simultaneous collision. This is incorrect because the reaction proceeds through a complex mechanism typically involving intermediate steps rather than a single simultaneous collision between two HI molecules. Therefore, while the molecularity is 2 (making the Assertion true), the Reason is false since the simultaneous collision of two reactant molecules is not the actual process for this decomposition.

Question 19. Assertion (A) : Zinc, cadmium and mercury are not considered as transition elements. Reason (R) : These elements have completely filled orbitals in their ground state as well as in their common oxidation states.

[1 Marks]

(A) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A).

(B) Assertion (A) is true, but Reason (R) is false.

(C) Both Assertion (A) and Reason (R) are true, but Reason (R) is not the correct explanation of the Assertion (A).

(D) Assertion (A) is false, but Reason (R) is true.

Explanation: Zinc, cadmium and mercury are not considered transition elements because they have completely filled d orbitals (d10 configuration) in both their ground state and common oxidation states. Transition elements are defined as elements having incompletely filled d orbitals either in their neutral atoms or in their common oxidation states. Since these elements have fully filled d orbitals, they do not meet the criteria to be classified as transition elements. Hence, both the assertion and reason are true, and the reason correctly explains the assertion.

Question 20.

Assertion (A) : Aromatic primary amines can be prepared by Gabriel Phthalimide synthesis. Reason (R) : Aryl halides do not undergo nucleophilic substitution with the anion formed by phthalimide.

[1 Marks]

(A) Assertion (A) is false, but Reason (R) is true.

(B) Assertion (A) is true, but Reason (R) is false

(C) Both Assertion (A) and Reason (R) are true, but Reason (R) is not the correct explanation of the Assertion (A).

(D) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A).

Explanation: Gabriel phthalimide synthesis is used to prepare primary amines usually from alkyl halides, but aromatic primary amines (aryl amines) cannot be prepared by this method because aryl halides do not undergo nucleophilic substitution with the anion formed by phthalimide. Therefore, the Assertion (A) is false, and the Reason (R) is true.

Question 21.

Assertion (A) : Glucose gets oxidized to six carbon gluconic acid on reaction with bromine water.
Reason (R) : The carbonyl group is absent in the open chain structure of glucose.

[1 Marks]

(A) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A).

(B) Both Assertion (A) and Reason (R) are true, but Reason (R) is not the correct explanation of the Assertion (A).

(C) Assertion (A) is false, but Reason (R) is true.

(D) Assertion (A) is true, but Reason (R) is false.

Explanation: Glucose does get oxidized to gluconic acid on reaction with bromine water, which shows that the aldehydic carbonyl group is present in the open-chain form of glucose. Therefore, Assertion (A) is true. However, the reason given that the carbonyl group is absent in the open-chain structure of glucose is false; the aldehyde group is indeed present in the open-chain form, which is why bromine water can oxidize glucose. Thus, Reason (R) is false.

Section C

Question 22.

The concentration of the reactant is reduced from 0.6 mol L^{-1} to 0.2 mol L^{-1} in 5 minutes in a first order reaction. Calculate rate constant of the reaction. (log 3 = 0.48)

[2 Marks]

Answer: Given: Initial concentration, $[R]_0$ = 0.6 mol L^{-1}, Final concentration, [R] = 0.2 mol L^{-1}, Time, t = 5 minutes. For a first order reaction, the integrated rate law is: $\ln([R]_0/[R]) = kt$. Taking logarithm base 10: $\log([R]_0/[R]) = (k\ t) / 2.303$. Calculate $\log([R]_0/[R]) = \log(0.6/0.2) = \log 3 = 0.48$. Then, $k = (2.303 / t) * \log([R]_0/[R]) = (2.303 / 5) * 0.48 = 0.221$ per minute. Therefore, the rate constant k = 0.221 min^{-1}.

Key Points: First order reaction integrated rate law - $\log([R]_0/[R]) = (k\ t)/2.303$ - Use given concentration and time values - Substitute and solve for k using log 3 = 0.48

Question 23.

Rate constant k for the first order reaction is $2.54 \times 10^{-3}\ s^{-1}$. Calculate the time required for three-fourth of the reactant to decompose. (log 4 = 0.60)

[2 Marks]

Answer: Given: $k = 2.54 * 10^{-3}\ s^{-1}$, fraction decomposed = 3/4, so fraction remaining = 1/4.
For a first order reaction, the time t required is given by:
$\ln([A]_0/[A]) = kt$
Here, $[A]_0/[A] = 1 / (1/4) = 4$
Therefore, $t = (1/k) * \ln(4)$
$\ln(4) = 2.303 * \log 4 = 2.303 * 0.60 = 1.3818$
So, $t = 1.3818 / (2.54 * 10^{-3}) = 543.7$ s
Hence, the time required for three-fourth of the reactant to decompose is approximately 544 seconds.

Key Points: First order reaction formula - Relationship between concentration and time - Calculation of time using log values

Question 24.

(a) Define activation energy.

(b) Write the unit of rate constant for

(i) zero order reaction,

(ii) second order reaction.

[2 Marks]

Answer: (a) **Activation energy** is the minimum energy that reacting molecules must have to collide successfully and form products in a chemical reaction.
(b) Units of rate constant:
(i) For zero order reaction, the unit of rate constant is mol $L^{-1}\ s^{-1}$ (molarity per second).
(ii) For second order reaction, the unit of rate constant is L $mol^{-1}\ s^{-1}$ (liter per mole per second).

Key Points: Definition of activation energy - Meaning of activation energy as minimum energy for reaction - Units of rate constant for zero and second order reactions

Question 25.

(a) First ionization enthalpy of Cr is lower than that of Zn. Why ?

(b) Mention any one property of the transition elements which makes them good catalysts.

[2 Marks]

Answer: (a) The first ionization enthalpy of Cr is lower than Zn because Cr has an electron configuration of [Ar] $3d^5$ $4s^1$, where removing one electron from 4s is easier due to its half-filled stable d-orbitals, whereas Zn has a fully filled $3d^{10}$ $4s^2$ configuration which is more stable and requires more energy to remove an electron.
(b) Transition elements are good catalysts because they have variable oxidation states which allow them to facilitate different reaction pathways and provide active sites for reactions.

Key Points: Lower first ionization enthalpy of Cr due to half-filled d-orbitals - Zn has fully filled d-orbitals making electron removal harder - Variable oxidation states enhance catalytic activity

Question 26.

(a) Define coordination number.

(b) Indicate the type of isomerism exhibited by the following complex : $[Co(en)_3]Cl_3$

[2 Marks]

Answer: (a) Coordination number is the number of ligand donor atoms that are directly attached to the central metal atom or ion in a coordination complex.
(b) The complex $[Co(en)_3]Cl_3$ exhibits optical isomerism because it contains three bidentate ethylenediamine (en) ligands arranged octahedrally, forming non-superimposable mirror images called enantiomers.

Key Points: Coordination number is the number of donor atoms bonded to metal - $[Co(en)_3]Cl_3$ contains bidentate ligands - Complex shows optical isomerism due to chiral arrangement

Question 27.

(a) Draw the structure of the given compound :

4–Bromo–3–methylpent–2–ene

(b) What happens when chloroethane is treated with aqueous potassium hydroxide ?

[2 Marks]

Answer: (a) The structure of 4–Bromo–3–methylpent–2–ene is CH3-CH=C(CH3)-CHBr-CH3 where the double bond is between C2 and C3, a methyl group is attached to C3, and a bromine atom is attached to C4.
(b) When chloroethane (C2H5Cl) is treated with aqueous potassium hydroxide (KOH), nucleophilic

substitution occurs. The hydroxide ion (OH-) replaces the chlorine atom, forming ethanol (C2H5OH) and potassium chloride (KCl) as a byproduct. The reaction is C2H5Cl + KOH (aq) -> C2H5OH + KCl.

Key Points: Structure of 4-Bromo-3-methylpent-2-ene - Nucleophilic substitution reaction - Formation of ethanol from chloroethane

---

Section D

Question 28.

When 1.5 g of a non-volatile solute was dissolved in 90 g of benzene, the boiling point of benzene got raised from 353.23 K to 353.93 K. Calculate the molar mass of the solute. (Given K_b for benzene = 2.52 K kg mol^{-1})

[3 Marks]

Answer: Given: Mass of solute = 1.5 g, Mass of benzene solvent = 90 g = 0.090 kg, Original boiling point = 353.23 K, New boiling point = 353.93 K, Boiling point elevation constant K_b = 2.52 K kg mol^{-1}.

Step 1: Calculate the elevation in boiling point, Delta T_b = New boiling point - Original boiling point = 353.93 K - 353.23 K = 0.7 K.

Step 2: Use the formula for boiling point elevation, Delta T_b = K_b * molality (m), so molality m = Delta T_b / K_b = 0.7 / 2.52 = 0.2778 mol/kg.

Step 3: Calculate moles of solute = molality * mass of solvent (kg) = 0.2778 * 0.090 = 0.025 mol.

Step 4: Calculate molar mass M = mass of solute / moles of solute = 1.5 g / 0.025 mol = 60 g/mol.

Answer: The molar mass of the solute is 60 g/mol.

Key Points: Calculate boiling point elevation - Use molality = Delta T_b / K_b - Calculate moles and molar mass

Question 29.

Calculate emf of the following cell at 298 K :

Sn | Sn^{2+} (0.001 M) || H^+ (0.01 M) | H_2(g) (1 bar) | Pt(s)

Given : $E°_{Sn^{2+}/Sn}$ = – 0.14 V,

$E°_H^+/H_2 = 0.00$ V [log 10 = 1]

[3 Marks]

Answer: The cell reaction is Sn (s) + $2H^+$ (aq) → Sn^{2+} (aq) + H_2 (g). Given E° for Sn^{2+}/Sn = -0.14 V and E° for H^+/H_2 = 0.00 V. The emf of the cell is calculated using the Nernst equation:

$E = E° - (0.0592 / n) \log Q$

Here, n = 2 (number of electrons transferred). Reaction quotient $Q = [Sn^{2+}] / [H^+]^2 = 0.001 / (0.01)^2 = 0.001 / 0.0001 = 10$.

Substitute the values:

$E = (-0.14) - (0.0592 / 2) * \log 10 = -0.14 - 0.0296 * 1 = -0.14 - 0.0296 = -0.1696$ V.

Therefore, the emf of the cell at 298 K is approximately -0.17 V.

Key Points: Identify cell reaction - Use Nernst equation - Calculate reaction quotient Q - Substitute values and solve for emf

Question 30.

Complete and balance the following equations :

(i) $2KMnO_4 \xrightarrow{-513\ K}$

(ii) $Na_2Cr_2O_7 + 2KCl \rightarrow$

(b) Why is it difficult to separate lanthanoid elements in pure state ?

[3 Marks]

Answer: (i) $2KMnO_4 - 513\ K \rightarrow K_2MnO_4 + MnO_2 + K_2O$

(ii) $Na_2Cr_2O_7 + 2KCl \rightarrow 2NaCl + K_2Cr_2O_7$

Explanation for (i): On heating at 513 K, potassium permanganate ($KMnO_4$) decomposes to form potassium manganate (K_2MnO_4), manganese dioxide (MnO_2) and potassium oxide (K_2O). This is a thermal decomposition reaction.

Explanation for (ii): Sodium dichromate ($Na_2Cr_2O_7$) reacts with potassium chloride (KCl) to form potassium dichromate ($K_2Cr_2O_7$) and sodium chloride (NaCl). It is a double displacement reaction.

(b) Why is it difficult to separate lanthanoid elements in pure state?

Lanthanoid elements are difficult to separate because they have very similar chemical properties and ionic radii, resulting in very similar reactivities and chromatographic behaviours. Their elements occur together in nature and their compounds are often mixed, making separation and isolation of pure individual lanthanoids challenging.

Key Points: Balanced thermal decomposition of $KMnO_4$ - Double displacement reaction of $Na_2Cr_2O_7$ and KCl - Lanthanoids have similar chemical properties making separation difficult

Question 31.

(a) $[Co(NH_3)_6]^{3+}$ is diamagnetic whereas $[CoF_6]^{3-}$ is paramagnetic. Justify the statement. [Atomic number of Co = 27]

(b) Write the electronic configuration for d4 ion if $\Delta_0 > P$ on the basis of crystal field theory.

[3 Marks]

Answer:

(a) In $[Co(NH_3)_6]^{3+}$, Co is in +3 oxidation state with electronic configuration $3d^6$. NH_3 is a strong field ligand causing pairing of all 6 electrons in the lower energy t_2g orbitals, resulting in no unpaired electrons, making it diamagnetic. The complex involves d_2sp^3 hybridisation (inner orbital complex).
(b) For a d^4 ion where $\Delta 0 > P$ (strong field ligand), electrons pair in the lower energy t_2g orbitals before occupying eg orbitals. Hence the configuration is $t_2g^4eg^0$ with 0 unpaired electrons.

Key Points: [Co(NH3)6]3+ is diamagnetic due to strong field ligand causing pairing - [CoF6]3- is paramagnetic due to weak field ligand with unpaired electrons - For d4 with Delta_0 > P, electrons pair in t2g orbitals leading to 0 unpaired electrons


Question 32.

(b) Define Racemic mixture.

(c) p-dichlorobenzene has a higher melting point as compared to its ortho and meta isomers. Why ?

[3 Marks]

(a) Write the major product in the following reactions :



Answer: (b) A racemic mixture is a mixture containing equal amounts of two enantiomers of a chiral molecule. It shows no optical activity because the rotations caused by each enantiomer cancel each other. It is also called racemic modification and is represented by prefixing dl or (±) before the compound name, e.g. (±) butan-2-ol. (c) p-Dichlorobenzene has a higher melting point than its ortho and meta isomers because in the para isomer, molecules are more symmetrical and pack more

efficiently in the solid state. This close packing leads to stronger intermolecular forces and hence a higher melting point.

Key Points: Racemic mixture contains equal enantiomers - Equal enantiomers cause zero optical rotation - p-Dichlorobenzene is more symmetrical - Efficient packing in solid state increases melting point - Ortho and meta isomers have less symmetry and lower melting points

Question 33.

Write the reaction involved in

(a) Rosenmund's reduction

(b) Cannizzaro's reaction

(c) Hell – Volhard – Zelinsky reaction

[3 Marks]

Answer: (a) **Rosenmund's reduction:** It is the selective hydrogenation of acid chlorides to aldehydes using hydrogen gas and palladium catalyst poisoned with barium sulfate.
Reaction: $RCOCl + H_2 \xrightarrow[Pd/BaSO_4]{} RCHO + HCl$

(b) **Cannizzaro's reaction:** Aldehydes without alpha hydrogen undergo disproportionation in presence of concentrated alkali to give alcohol and carboxylic acid.
Reaction: $2\ RCHO + OH^- \xrightarrow{} RCOO^- + RCH_2OH$

(c) **Hell-Volhard-Zelinsky reaction:** Alpha bromination of carboxylic acids using PBr_3 and Br_2 to form alpha-bromo acid.
Reaction: $RCH_2COOH + PBr_3 + Br_2 \xrightarrow{} RCHBrCOOH + HBr$

Key Points: Rosenmund reduction converts acid chlorides to aldehydes using H_2 and $Pd/BaSO_4$- Cannizzaro reaction involves disproportionation of aldehydes without alpha hydrogen in basic medium- Hell-Volhard-Zelinsky reaction brominates alpha position of carboxylic acids using PBr_3 and Br_2

Question 34.

Define the following term:

(a) Reducing sugar

(b) Differentiate between the following :

(i) Fibrous proteins and globular proteins

(ii) Nucleotide and nucleoside

[3 Marks]

Answer: (a) Reducing sugar: Reducing sugars are carbohydrates that can reduce Fehling's solution and Tollens' reagent because they have free aldehyde or ketone groups. All monosaccharides like glucose and fructose are reducing sugars. Some disaccharides like maltose and lactose are reducing sugars, whereas sucrose is non-reducing because its aldehyde or ketone groups are involved in glycosidic bond.

(b) Differences:

(i) Fibrous proteins vs Globular proteins:

Fibrous proteins have long, thread-like structures, are insoluble in water, and provide structural support, e.g., keratin and collagen. Globular proteins are spherical in shape, soluble in water, and perform metabolic functions, e.g., enzymes and hormones.

(ii) Nucleotide vs Nucleoside:

Nucleotides consist of a nitrogenous base, a pentose sugar, and one or more phosphate groups. Nucleosides consist of only a nitrogenous base and a pentose sugar, without phosphate groups.

> **Key Points: Reducing sugars have free aldehyde or ketone groups - Fibrous proteins are structural and insoluble, globular are functional and soluble - Nucleotide has phosphate group, nucleoside does not**

Question 35.

Write the reaction of glucose with the following :

(a) $H_2N - OH$

(b) $(CH_3CO)_2O$

(c) conc.HNO_3

[3 Marks]

Answer: (a) Reaction with $H_2N - OH$ (Hydroxylamine): Glucose contains an aldehyde group (–CHO) which reacts with hydroxylamine to form an oxime. The reaction confirms the presence of the aldehyde functional group in glucose.

(b) Reaction with $(CH_3CO)_2O$ (Acetic anhydride): Glucose reacts with acetic anhydride to form glucose pentaacetate by acetylation of all the hydroxyl (–OH) groups. In this reaction, all five –OH groups of glucose are replaced by acetate groups ($–OCOCH_3$). This derivative does not react with hydroxylamine indicating the absence of free aldehyde group in the acetylated product.

(c) Reaction with conc. HNO_3 (Concentrated nitric acid): Glucose is oxidized by concentrated nitric

acid to form saccharic acid, where both aldehyde and primary alcohol groups are oxidized to carboxylic acid groups. This confirms the presence of aldehyde and $–CH_2OH$ groups in glucose.

Key Points: Glucose reacts with hydroxylamine to give oxime - Glucose forms pentaacetate with acetic anhydride by acetylation - Concentrated nitric acid oxidizes glucose to saccharic acid showing presence of aldehyde and primary alcohol groups

---

Section E

Question 36.

(a) Give chemical tests to distinguish between the following pairs of compounds :

(i) Acetophenone and Benzophenone

(ii) Propanal and Propanone

(iii) Pentan–2–one and Pentan–3–one

(b) (i) Which of the following acids is stronger and why ?

CH_2FCOOH, CH_3COOH

(ii) Arrange the following compounds in the increasing order of their boiling points :

CH_3OCH_3, $CH_3CH_2CH_3$, CH_3CH_2OH, CH_3CHO

[5 Marks]

Answer: (a) Chemical tests to distinguish compounds:
(i) Acetophenone shows no reaction with Fehling's solution or Tollens' reagent, whereas Benzophenone does not react because both are ketones; however, both can be distinguished by their physical properties or by specific reactions like Benedict's test, but generally, no simple test distinguishes them easily.
(ii) Propanal (an aldehyde) gives a positive Tollens' test (silver mirror) and Fehling's test (red ppt), whereas Propanone (a ketone) gives negative results.
(iii) Pentan-2-one and Pentan-3-one do not give simple chemical tests for differentiation as both are ketones with similar properties; their distinction requires spectroscopic methods or more advanced techniques.
(b) (i) Stronger acid is CH_2FCOOH because the electronegative fluorine atom stabilizes the carboxylate ion by electron withdrawing inductive effect, increasing acidity compared to CH_3COOH.
(ii) Increasing order of boiling points is CH_3OCH_3 (dimethyl ether) < $CH_3CH_2CH_3$ (propane) < CH_3CHO (acetaldehyde) < CH_3CH_2OH (ethanol). This is because alcohols have strong

hydrogen bonding leading to highest boiling point, aldehydes have dipole-dipole interactions, ethers have weaker dipole interactions, and alkanes have weak van der Waals forces.

> **Key Points: Distinguishing aldehyde and ketone by Tollens test - Inductive effect of electronegative fluorine increases acid strength - Hydrogen bonding increases boiling point - Order of boiling points based on intermolecular forces**

Question 37.

(a) The conductivity of 0.1 mol L^{-1} solution of NaCl is 1.06×10^{-2} S cm^{-1}. Calculate its molar conductivity and degree of dissociation.

$\lambda^\circ_{Na^+} = 50.1$ S cm^2 mol^{-1}

$\lambda^\circ_{Cl^-} = 76.5$ S cm^2 mol^{-1}

(b) (i) Following cell reaction occurs in a galvanic cell :

$2Ag^+(aq) + Zn(s) \rightarrow 2Ag(s) + Zn^{2+}(aq)$ $E^\circ_{(cell)} = +1.56$ V

Predict the direction of flow of current.

(ii) Differentiate between a primary battery and a secondary battery.

[5 Marks]

Answer: (a) Given:
Conductivity, $\kappa = 1.06 * 10^{-2}$ S cm^{-1}, Concentration, c = 0.1 mol L^{-1}, $\lambda^\circ_{Na^+} = 50.1$ S cm^2 mol^{-1}, $\lambda^\circ_{Cl^-} = 76.5$ S cm^2 mol^{-1}

Step 1: Calculate molar conductivity (Λm).
Formula: $\Lambda m = \kappa / c = (1.06 * 10^{-2}) / 0.1 = 1.06 * 10^{-1}$ S cm^2 mol^{-1}

Step 2: Calculate molar conductivity at infinite dilution ($\Lambda^\circ m$).
$\Lambda^\circ m = \lambda^\circ_{Na^+} + \lambda^\circ_{Cl^-} = 50.1 + 76.5 = 126.6$ S cm^2 mol^{-1}

Step 3: Calculate degree of dissociation (α).
$\alpha = \Lambda m / \Lambda^\circ m = 1.06 * 10^{-1} / 126.6 = 0.000838 \approx 8.38 * 10^{-4}$

Therefore, molar conductivity $\Lambda m = 0.106$ S cm^2 mol^{-1} and degree of dissociation $\alpha \approx 8.38 * 10^{-4}$.

(b)(i) The given reaction is $2Ag^+(aq) + Zn(s) \rightarrow 2Ag(s) + Zn^{2+}(aq)$ with $E^\circ_{cell} = +1.56$ V.
Since E°_{cell} is positive, the reaction is spontaneous under standard conditions.
Electrons flow from the anode to cathode; Zn is oxidized (anode) and Ag^+ is reduced (cathode).

Therefore, current flows from Ag electrode to Zn electrode externally.

(b)(ii) Difference between primary and secondary batteries:
- Primary battery: Non-rechargeable, chemical reactions are not reversible, once discharged it cannot be reused.
- Secondary battery: Rechargeable, chemical reactions are reversible, can be reused many times by recharging.
Examples: Primary - dry cell; Secondary - lead acid battery.

Key Points: Molar conductivity formula - Calculation of degree of dissociation - Direction of current in galvanic cell - Difference between primary and secondary batteries

Question 38.

(a) Resistance of a conductivity cell filled with 0.1 M KCl solution is 100 Ω. If the resistance of the same cell when filled with 0.01 mol L^{-1} KCl solution is 300 Ω, calculate the conductivity and molar conductivity of 0.01 mol L^{-1} KCl solution. The conductivity of 0.1 M KCl solution is 1.29×10^{-2} S cm^{-1}.

(b) (i) Write any two advantages of $H_2 – O_2$ fuel cell.

(ii) Why does the cell potential of mercury cell remain constant throughout the life ?

[5 Marks]

Answer: (a) Given: Resistance R_1=100 Ω for 0.1 M KCl; conductivity κ_1=1.29 * 10^{-2} S cm^{-1};

Resistance R_2=300 Ω for 0.01 mol L^{-1} KCl.

Cell constant, $G^* = \kappa_1 * R_1$ = 1.29 * 10^{-2} S cm^{-1} * 100 Ω = 1.29 cm^{-1}.

Conductivity of 0.01 mol L^{-1} KCl, $\kappa_2 = G^* / R_2$ = 1.29 / 300 = 4.3 * 10^{-3} S cm^{-1}.

Molar conductivity, $\lambda_m = \kappa_2$ / concentration = 4.3 * 10^{-3} / 0.01 = 0.43 S cm^2 mol^{-1} or 43 S cm^2 mol^{-1}

(since usually molar conductivity is expressed in S cm^2 mol^{-1}).

(b)(i) Advantages of H_2 - O_2 fuel cell:
1. They produce water as the only product, so they are environmentally friendly.
2. They have high efficiency compared to conventional fuel cells.

(b)(ii) The cell potential of mercury cell remains constant throughout its life because the reaction involves pure substances in their standard states and the concentration of reactants and products does not change significantly. The electrodes remain stable, so the cell potential remains constant until the reactants are used up.

Key Points: Calculation of cell constant - Use of resistance and conductivity to find unknown conductivity - Calculation of molar conductivity - Advantages of H2-O2 fuel cell

- **Reason for constant cell potential of mercury cell**

Question 39.

An organic compound ‘A’, with molecular formula C_2H_6O reacts with active metals such as sodium to give compound ‘B’ and hydrogen gas. 'A' on treatment with iodine and sodium hydroxide gives ‘C’ and in presence of H_2SO_4 at 413 K gives ‘D’ ($C_4H_{10}O$). ‘D’ on reaction with excess of HI gives ‘E’. Identify ‘A’, ‘B’, ‘C’, ‘D’ and ‘E’ and write all the reactions involved.

[5 Marks]

Answer: Identification of compounds and reactions:

(a) Compound 'A' with formula C_2H_6O could be either ethanol (an alcohol) or dimethyl ether. Since 'A' reacts with sodium metal to give hydrogen gas, it must be an alcohol. Thus, **A is ethanol (CH_3CH_2OH)**.

Reaction with sodium:

$CH_3CH_2OH + 2Na \rightarrow CH_3CH_2ONa$ ('B') + H_2 (gas)

So, **B is sodium ethoxide (CH_3CH_2ONa)**.

(b) On treatment with iodine and sodium hydroxide, 'A' gives a yellow precipitate 'C'. This is the iodoform test, indicating methyl ketone or ethanol.

Reaction:

$CH_3CH_2OH + 4I_2 + 6NaOH \rightarrow CHI_3$ (iodoform, yellow precipitate) + HCOONa + 5NaI + $5H_2O$

Thus, **C is iodoform (CHI_3)**.

(c) 'A' in presence of concentrated H_2SO_4 at 413 K undergoes dehydration to form an ether 'D' with molecular formula $C_4H_{10}O$.

Reaction:

$2\ CH_3CH_2OH$ -(H_2SO_4, 413 K)-> $CH_3CH_2OCH_2CH_3$ (diethyl ether, 'D') + H_2O

So, **D is diethyl ether**.

(d) 'D' on reaction with excess HI undergoes cleavage to give 'E'.

Reaction:

$CH_3CH_2OCH_2CH_3 + 2HI \rightarrow 2\ CH_3CH_2I$ ('E') + H_2O

So, **E is ethyl iodide (CH_3CH_2I)**.

Summary of compounds and reactions:

A: Ethanol (CH_3CH_2OH)

B: Sodium ethoxide (CH_3CH_2ONa)

C: Iodoform (CHI_3)

D: Diethyl ether ($CH_3CH_2OCH_2CH_3$)

E: Ethyl iodide (CH_3CH_2I)

Each step is a classic reaction: sodium metal with alcohol, haloform reaction with iodine and NaOH, acid catalysed dehydration of alcohol to ether, and cleavage of ether by HI to alkyl iodide.
This shows understanding of alcohol properties and reactions.

> **Key Points: A is ethanol - B is sodium ethoxide - C is iodoform - D is diethyl ether - E is ethyl iodide - Sodium metal reacts with alcohol to release hydrogen - Iodoform test confirms presence of methyl ketone or ethanol - Acid catalysed dehydration forms ether - Ether cleavage by HI yields alkyl iodide**

Question 40.

Write the reagents which are used in the given conversions :

(i) Phenol to 2, 4, 6 – tribromophenol

(ii) Propene to propan–1–ol

(iii) Butan–2–one to butan–2–ol

(b) Explain the mechanism of acid catalyzed hydration of alkene to form corresponding alcohol.

[5 Marks]

Answer: (a) Reagents used in the conversions:

(i) Phenol to 2, 4, 6–tribromophenol: Bromine water (Br_2 in H_2O)

(ii) Propene to propan–1–ol: BH_3/THF followed by H_2O_2/OH^- (hydroboration-oxidation)

(iii) Butan–2–one to butan–2–ol: $NaBH_4$ in methanol (reduction of ketone)

(b) Mechanism of acid catalyzed hydration of alkene:

1. Protonation of alkene: The alkene reacts with an acid (like H_2SO_4) and the double bond attacks H^+. This forms the most stable carbocation intermediate as per Markovnikov's rule.
2. Nucleophilic attack by water: Water molecule attacks the carbocation intermediate, forming an oxonium ion (a protonated alcohol).
3. Deprotonation: Another water molecule removes a proton (H^+) from the oxonium ion to yield the alcohol.

This process adds water (H and OH) across the double bond, converting an alkene to an alcohol. The reaction is acid catalyzed because the acid provides the H^+ needed to initiate the reaction and then is regenerated at the end.

Key Points: Phenol to tribromophenol uses bromine water - Propene to propan-1-ol uses hydroboration-oxidation - Butan-2-one to butan-2-ol uses NaBH4 reduction - Acid catalyzed hydration involves protonation, carbocation formation, nucleophilic attack by water and deprotonation

Question 41.

(a) How will you convert the following :


(i) Propanone to Propene

(ii) Benzoic acid to Benzaldehyde


(iii) Benzene to m-Nitroacetophenone

[5 Marks]

(b) (i) Arrange the following compounds in increasing order of their reactivity towards HCN :



(ii) Identify the compounds, which would undergo Aldol condensation :



Answer: (a)(i) Conversion of Propanone to Propene:
Propanone (CH3COCH3) can be converted to Propene (CH3CH=CH2) by first converting propanone to its hydrazone derivative using hydrazine (NH2NH2) and then treating it with a strong base like potassium hydroxide (KOH) under heat. This is known as the Wolff-Kishner reduction which removes the carbonyl oxygen and forms the alkene, propene.

(a)(ii) Conversion of Benzoic acid to Benzaldehyde:
Benzoic acid (C6H5COOH) can be reduced to benzaldehyde (C6H5CHO) using a mild reducing agent such as stannous chloride (SnCl2) in hydrochloric acid or by partial reduction with lithium aluminium hydride (LiAlH4) under controlled conditions. Alternatively, the acid chloride of benzoic acid can be formed using thionyl chloride (SOCl2), then partial reduction of the acid chloride to benzaldehyde using lithium tri-tert-butoxyaluminium hydride.

(a)(iii) Conversion of Benzene to m-Nitroacetophenone:

Step 1: Acetophenone is formed by Friedel-Crafts acylation of benzene with acetyl chloride (CH_3COCl) in the presence of anhydrous aluminum chloride ($AlCl_3$).
Step 2: Nitration of acetophenone is done by treating it with a mixture of concentrated nitric acid (HNO_3) and concentrated sulfuric acid (H_2SO_4). The acetyl group is a meta-directing group, so nitration occurs at the meta position giving m-nitroacetophenone.

Key Points: Propanone to propene via Wolff-Kishner reduction - Benzoic acid to benzaldehyde by partial reduction or via acid chloride intermediate - Benzene to m-nitroacetophenone by Friedel-Crafts acylation followed by nitration at meta position