

OUR BEST
ANNUAL
PROGRAM



4 STEP PROGRAM

Covers 5,000 SQ FT

KEEP OUT OF REACH OF CHILDREN
CAUTION
See back for additional precautionary statements.

ACTIVE INGREDIENTS:	
2,4-D††	1.21%
Mecoprop-p††††	0.61%
OTHER INGREDIENTS:	
Total	98.18%
	100.00%

†0.0121 lbs 2,4-D acid equivalent per pound of product
††0.0061 lbs MCPP-p acid equivalent per pound of product

28-0-3

NET WT 14.29 lb (6.48 kg)



23616-3



STEP²

Weed Control plus Lawn Food²


DO NOT USE ON ST. AUGUSTINEGRASS, FLORATAM,
DICHONDRA, LIPPIA, CARPETGRASS OR BENTGRASS LAWNS.



Apply in Late Spring

STEP²

Weed Control plus Lawn Food²



Now it's Easy for You to Have a Better Lawn



Start anytime
You can begin the recommended Annual Program anytime of the year. Just check the month and select the appropriate STEP® product listed.

Simple and easy
No more guesswork. You simply apply the right product – one STEP® at a time.

Guaranteed results
Each STEP® product is designed to the specific season to help give you a greener lawn.

SPREADER SETTINGS

Spreading Tips: Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scott's® spreader. Covers 5,000 sq ft This package applies 2.86 lb of product per 1,000 sq ft

Spreader Settings	Directions	Spreader Settings	Directions
Scott's® Broadcast Rotary Spreaders 3 1/2	Each pass should be within 5 feet of each other to ensure overlap. 	Drop Spreaders 5	Make sure spreader wheel completely overlaps previous wheel marks. 

For additional spreader settings, please call 1-800-543-TURF (8873).

GET EXPERT ADVICE FOR YOUR LAWN NOW!

Want information on what to use and when? Sign up today to get FREE expert advice specifically for your lawn.

Consumer Helpline:
1-800-543-TURF
(8873)

Visit our website:
www.scotts.com



Well-fed lawns are more efficient at using available water.

Feeding with Scott's® lawn food helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn against heat, drought and other stresses.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This product applies 1.5 pounds of 2,4-D per acre per application.

APPLICATION PRECAUTIONS

- Apply when weeds are actively growing when daytime temperatures are consistently above 60 °F.
- Apply on a calm day.
- Apply to moist grass. To help particles stick to weed leaves, the grass should be thoroughly watered or wet from dew before applying this product.
- Grass should not touch the bottom of a drop spreader. Tall grass should be mowed a day or two before application.

APPLICATION RESTRICTIONS

- Do not allow product to contact people or pets directly or through drift. Keep people and pets off lawn during application. Do not allow people or pets to enter treated areas until dusts have settled.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, seeps, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product runoff, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.
- On hilly and terraced lawns, wait until after the second mowing of the year but NOT before mid-March. In the fall, DO NOT treat centipede after the first of October and DO NOT treat bahia after the first of November.
- Do not use this product on flowers, shrubs, fruit or vegetable plants or on creeping bentgrass, St. Augustine, Floratam and dichondra lawns, Or. Bermuda grass, bahia, centipede grass, zoysia lawns, under certain conditions, some temporary discoloration may result.
- Apply only to grass. Do not allow granules or dust to drift onto flowers, shrubs, fruit or vegetables.
- The maximum number of broadcast applications per treatment site is two per year.
- Allow a minimum of 30 days between applications.

EXTREME TEMPERATURES

- Do not apply when temperatures are consistently above 90 °F.

WATERING OR RAIN

- Do not water or mow lawn for at least 24 hours after application. Granules must remain on weed leaves for 24 hours to be effective.
- Do not apply if rain is expected within the next 24 hours.

IMPORTANT

WHAT TO EXPECT

- Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for control.
- This product will not prevent weed seeds from producing new plants.
- Control the following weeds: aster, bindweed, bitterroot, black medic, buckhorn, bull thistle, burdock, butterweed, Canada thistle, carpetweed, catnip (coat dandelion), chickweed, stitchwort (Stellaria sp.), chionox, cinquefoil, clover (hop), clover (red), clover (white), culant, curly dock, dandelion, dichondra, dollarweed/pennycress, English daisy, evening primrose, flax, foxtail, ground ivy, hawkweed, heath, henbit, henwood, knotweed (seedling), limonum, lupine, lupinus (seedling), lupine (leafhopper), mallow, monardella, morning glory, mustard, pepperweed/pigeonweed, pigweed, plantain (broadleaf), purslane, radish (Richardia spp.), ragwort, red sorrel (sheep sorrel), scarlet pimpernel, shepherd-purse, smartweed, sowthistle, sparrowweed/lawn burweed, violet, wild carrot, yellow, yellow nodules.

SEEDING/SODDING/SPRINGING

- Wait until the new grass has been mowed at least four times before applying STEP® 2 Weed Control Plus Lawn Food.
- If you are planning to seed, sod or spring, you should wait at least one month after you have applied STEP® 2 Weed Control Plus Lawn Food.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a dry place. Reseal opened bag by folding top down and securing. Store in original container in a secure area away from children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, shoes and socks. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID
If in eyes: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment. For emergency information call 1-800-543-8873.

ENVIRONMENTAL HAZARDS:
This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.
This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.
EPA Reg. No. 538-282 EPA Est. No. 538-OH-1 *CAS # 94-75-7 **CAS # 16484-77-8

STEP® 2 Weed Control Plus Lawn Food 28-0-3 F 643
Guaranteed Analysis

Total Nitrogen (N)	28%
2.7% Ammoniacal Nitrogen	
13.9% Urea Nitrogen	
10.5% Other Water Soluble Nitrogen*	
0.9% Water Insoluble Nitrogen*	
Soluble Potash (K ₂ O)	3%
Sulfur (S)	4%
4.0% Combined Sulfur (S)	

Derived from: Ammonium Sulfate, Methyleneurea, Urea and Potassium Sulfate.
*8.5% slowly available Nitrogen from Methyleneurea.
Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatoreg-fbi-nc.com>.
Notice: To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling if this product not in accordance with directions.

SMG Soil and Guarantees by
BYNE The Scott's Company
14111 Scottslawn Rd.
Marysville, Ohio 43041
©2017 The Scott's Company LLC
World rights reserved.

Scott's No-Quibble Guarantee®
If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

SCOTT'S® STEP® 2 WEED CONTROL PLUS LAWN FOOD: 9M 23616-3

FPO

0 32247 23616 4
PROOF OF PURCHASE
VID-BB
BGS467

COVERS 5,000 SQUARE FEET