

10

FOR GENERAL

警告体征

原发性免疫系统缺陷病

原发性免疫系统缺陷病导致儿童和成人频繁地受感染和难以治疗。每500人里有1人被确诊。如果您或您身边的人有以下两个或更多的警告体征，咨询您的医生关于患有原发性免疫系统缺陷病的可能性。

- 1 一年中的耳部感染次数超过4次。
- 2 一年中出现两次或更多次严重的鼻窦感染。
- 3 使用抗生素两个月或更长时间，收效甚微。
- 4 一年中出现两次或更多次的肺部感染。
- 5 婴儿体重不增加或生长不正常。
- 6 经常出现深层皮肤或器官脓肿。
- 7 口腔或皮肤出现持续性鹅口疮。
- 8 需要静脉抗生素，以清除感染。
- 9 两次或两次以上的深部感染，包括败血症。
- 10 有原发性免疫缺陷病家族史。

Presented as a public service by:



Jeffrey Modell
Foundation

Curing PI.
Worldwide.



Funding was made possible in part by a grant from the U.S. Centers for Disease Control and Prevention (CDC).



National Heart,
Lung, and Blood
Institute (NHLBI)



National Institute of
Allergy and Infectious
Diseases (NIAID)



Baxter
BioScience

Biotest
From Nature for Life

CSL Behring
Biotherapies for Life™

GRIFOLS

KEDRION
BIOPHARMA

octapharma

These warning signs were developed by the Jeffrey Modell Foundation Medical Advisory Board. Consultation with Primary Immunodeficiency experts is strongly suggested. © 2013 Jeffrey Modell Foundation

For information or referrals, contact the Jeffrey Modell Foundation: info4pi.org | 866-INFO-4-PI