

# 10 Warning Signs of Primary Immunodeficiency

FOR ADULTS

**Primary Immunodeficiency (PI) causes children and adults to have infections that come back frequently or are unusually hard to cure. 1:500 persons are affected by one of the known Primary Immunodeficiencies. If you or someone you know is affected by two or more of the following Warning Signs, speak to a physician about the possible presence of an underlying Primary Immunodeficiency.**

- 1** Two or more new ear infections within 1 year.
- 2** Two or more new sinus infections within 1 year, in the absence of allergy.
- 3** One pneumonia per year for more than 1 year.
- 4** Chronic diarrhea with weight loss.
- 5** Recurrent viral infections (colds, herpes, warts, condyloma).
- 6** Recurrent need for intravenous antibiotics to clear infections.
- 7** Recurrent, deep abscesses of the skin or internal organs.
- 8** Persistent thrush or fungal infection on skin or elsewhere.
- 9** Infection with normally harmless tuberculosis-like bacteria.
- 10** A family history of PI.

Presented as a public service by:



Jeffrey Modell  
Foundation

Curing PI.  
Worldwide.



Funding was made possible in part by a grant from the U.S. Centers for Disease Control and Prevention (CDC).



National Heart,  
Lung, and Blood  
Institute (NHLBI)



Plasma Protein Therapeutics Association



National Institute of  
Allergy and Infectious  
Diseases (NIAID)



Eunice Kennedy Shriver  
NICHD  
National Institute of Child Health  
& Human Development



Rare Disease



Biotherapies for Life™



These warning signs were developed by the Jeffrey Modell Foundation Medical Advisory Board.  
Consultation with Primary Immunodeficiency experts is strongly suggested. © 2024 Jeffrey Modell Foundation

For information or referrals, contact the Jeffrey Modell Foundation: [info4pi.org](http://info4pi.org)