

Accuris™ Taq DNA Polymerase

Description

Highly soluble with an increased affinity for template DNA, Accuris Taq DNA Polymerase is suitable for a variety of assay conditions and templates. Improved yield, activity, sensitivity and speed are achieved by modifying the polymerase with the addition of hydrophilic residues. A proprietary 5X buffer system with PCR enhancers and optimal levels of MgCl₂ has been created to partner with Accuris Taq DNA Polymerase and is supplied in this package

- Performs across a wide range of DNA templates including genomic DNA and GC-rich and AT-rich sequences
- Proprietary 5X reaction buffer includes enhancers for maximizing enzyme activity and reaction speed
- Improved solubility and template affinity

Storage

Upon receipt, immediately store at -20°C. Avoid excessive freeze/thaw cycles. When stored as directed, this product will retain its activity for 12 months from date of receipt. The product may also be stored at 4°C for up to one month.

Limitations of Use

For research purposes only. Not intended for therapeutic or diagnostic use.

Quality Control

Accuris enzymes and reagents are tested under general assay conditions for activity, reproducibility, efficiency, heat activation, sensitivity, and absence of nuclease contamination and nuclease activity. This product is manufactured under a comprehensive quality management system, following ISO 9001:2008 standards.

General Guidelines

1. Reaction Buffer

The 5X reaction buffer supplied with the Accuris Taq DNA polymerase has been formulated for maximum efficiency, sensitivity and successful PCR with a variety of difficult templates. Proprietary PCR enhancers and 15mM MgCl₂ are included in the buffer. Use of additional PCR enhancers may have a negative effect on the reaction.

2. Template

For PCR of complex genomic DNA, 5ng - 500ng of template DNA may be added per reaction. Do not add more than 100ng of DNA for cDNA or plasmid DNA

3. Primers

Primers should have a predicted melting temperature of approximately 60°C, using default Primer 3 settings (<http://frodo.wi.mit.edu/primer3>). The final primer concentration should be 0.2µM to 0.6µM.

4. Annealing Temperature

An initial annealing temperature of 55°C is recommended. If nonspecific products appear, increase the temperature in 2°C increments. Alternately, a temperature gradient may be performed.

5. Extension

The polymerase performs optimally at 72°C. Extension time is dependent upon amplicon complexity and length. Generally, 30 seconds per kb is recommended for eukaryotic genomic DNA and cDNA. A five second extension is sufficient for shorter amplicons.

Technical Support

For trouble-shooting and tech support, contact us at info@accuris-usa.com or call 908 769-5555.

Accuris is not responsible for consequential or incidental damages, whether direct or indirect, resulting from use of this product. Accuris guarantees the performance of this product as described when used in accordance with these instructions. slj 09/21

Reaction setup

Prepare the reaction on ice as follows:

Component	25 µl reaction	50 µl reaction	Final concentration
Accuris 5x Taq Reaction Buffer	5 µl	10 µl	1X
100mM dNTPs (25mM each)	0.25 µl	0.5 µl	1mM
Forward Primer (10µM)	1.0 µl	2.0 µl	400 nM
Reverse Primer (10µM)	1.0 µl	2.0 µl	400 nM
Template DNA	<100ng cDNA, <500ng genomic		variable
Accuris Taq DNA Polymerase (5u/µl)	0.1µl - 0.5 µl	0.20 µl - 1 µl	variable
PCR-grade water	to final reaction volume		

For other volumes, adjust the amount of each component accordingly.

Gently mix the solution. If needed, spin briefly in a microcentrifuge to bring reaction mixture to the bottom of the tube. Transfer samples to a thermal cycler with the block preheated to 95°C and begin cycling.

Routine PCR Cycling

Step	Temperature	Time
Initial denaturation	95°C	1 minute
	95°C	15 seconds
55-40 cycles	55°C to 67°C*	15 seconds
	72°C	30 seconds per Kb

*Annealing temperature determined by user

Package contents and reordering

Taq Polymerase is supplied at a concentration of 5 units/µl. Available in 500, 1000 and 6000 unit packages with 5X Taq buffer.

Accuris Taq DNA Polymerase, sample pack

Catalog number PR1000-S
Includes 10 µl of enzyme and buffer.

Accuris Taq DNA Polymerase, 500 units

Catalog number PR1000-500
Includes 100 µl of enzyme and 4ml of 5X buffer in 1ml aliquots.


Accuris Taq DNA Polymerase, 1000 units

Catalog number PR1000-1000
Includes 200 µl of enzyme in 100 µl aliquots, and 8ml of 5X buffer in 1ml aliquots.

Accuris Taq DNA Polymerase, 6000 units

Catalog number PR1000-6000
Includes 1200 µl of enzyme in 100 µl aliquots, 48ml of 5X buffer in 1ml aliquots.

Accuris offers a full line of PCR enzymes and master mixes. Visit www.accuris-usa.com.



Accuris™
**Taq Polymerase
PR1000**

5 units/µl, supplied with 5X Buffer

Package contains:
500 units Taq Polymerase (1x100 µl)
4ml 5X Accuris Taq Buffer (4x1ml)
Store at -20°C upon receipt

PH: 908.769.5555 EM: Info@accuris-usa.com