

Check the product label for actual catalog number, lot and expiry date.

dNTP Sets & Mixes

CAT.#	SIZE	COMPONENTS	COMPONENT COMPOSITION
dNTP Mixes			
NUM0101	1 ml	2 x 0.5 ml - 25 mM dNTP Mix	Aqueous solution of equal concentration of 25 mM each of 4 dNTPs, pH 7.0
NUM0201	1 ml	2 x 0.5 ml - 10 mM dNTP Mix	Aqueous solution of equal concentration of 10 mM each of 4 dNTPs, pH 7.0
dNTP Set			
NUS0101	4 x 0.25 ml	0.25 ml - 100 mM dATP	Aqueous solution of 100 mM dATP, pH 7.0
		0.25 ml - 100 mM dCTP	Aqueous solution of 100 mM dCTP, pH 7.0
		0.25 ml - 100 mM dGTP	Aqueous solution of 100 mM dGTP, pH 7.0
		0.25 ml - 100 mM dTTP	Aqueous solution of 100 mM dTTP, pH 7.0

Storage In the dark at -20°C.

APPLICATIONS

- All molecular biology applications including dNTPs, like:
- cDNA synthesis
 - Standard PCR, Long and high-fidelity PCR
 - qPCR
 - Sequencing

BENEFITS

- Highest quality, >99% HPLC pure dNTPs for high & reproducible yields
- Pure from DNA contamination and from PCR inhibitors
- Highly stable remain pure after weeks at room temperature, after 30 freezing thawing cycles and during the 40 PCR cycles
- Available in ready-to use mixes and sets for maximized flexibility

PRODUCT DETAILS

highQu dNTP sets and mixes meet all highest industry standards and allow for unrivaled performance of your PCR and other DNA synthesis reactions.

Produced under the stringent quality monitoring conditions, they guaranty reproducible results. More than 99% HPLC purity eliminates inhibitions of PCR and allows for increased yields with higher dNTP concentrations.

Exceptional stability eliminates dNTP usability concerns related to short term ambient temperature shipments, room temperature storage or PCR exceeding 40 cycles.

PREPARATION OF dNTP MIXES FROM A SET

- Highly concentrated solutions require thorough mixing before the use.
- The optimal dNTP mix shall have equal concentrations of all 4 dNTPs.
- To prepare from a set of 4 dNTPs mixes of common concentrations, follow the guidelines below:

Use same volume of each from four 100 mM dNTP solutions:	PCR Water	Resulting 1 ml Mix concentration:
20 µl	920 µl	2 mM dNTP
25 µl	900 µl	2.5 mM dNTP
100 µl	600 µl	10 mM dNTP
250 µl	-	25 mM dNTP

PROTOCOL RECOMMENDATIONS FOR STANDARD PCR

- Typical concentration of each dNTP in the reaction is 0.2 – 0.25 mM. Higher concentration increase yields, however Mg²⁺ ions bind to dNTPs, therefore, both components shall be present in coordinated concentrations. Too high dNTPs and magnesium concentrations reduce PCR fidelity.
- Mix well each dNTP and magnesium solution, to avoid concentration fluctuations.
- Use final 3 mM MgCl₂ with 0.25 mM each dNTP concentration for routine PCR.

Starting dNTP Mix conc.	Vol. of dNTP mix for 50 µl rxn	Final Mg ²⁺ conc. in rxn	Vol. of 50 mM MgCl ₂ for 50 µl rxn to achieve desired conc.
10 mM	1.25 µl	2 mM	2 µl
25 mM	0.5 µl	3 mM	3 µl

IN VITRO RESEARCH USE ONLY

ORDERING

T: +49 7250 33 13 401
F: +49 7250 33 11 413
order@highQu.com
www.highQu.com

SALES

T: +49 7250 33 13 401
F: +49 7250 33 11 413
sales@highQu.com

TECHNICAL SUPPORT

T: +49 7250 33 13 401
F: +49 7250 33 11 413
tech-support@highQu.com