

Post-Operative Instructions: Dental Implants

It is imperative that you follow these instructions carefully to maximize your healing and improve the long-term outcome of your dental implant(s).

Do not disturb the wound. Avoid spitting and touching the wound for a few days after surgery. There may be a metal stump slightly protruding through the gum tissue.

Some bleeding or redness is normal for 24 hours. Excessive bleeding (where your mouth fills up rapidly with blood) can be controlled by biting on a gauze pad placed directly on the bleeding wound for 30 minutes. If bleeding continues, please call the office for further instructions.

Swelling is a normal occurrence after surgery. To minimize swelling, apply an ice pack on the cheek in the area of surgery. Apply the ice continuously, as much as possible, for the first 36 hours. You may place a towel between the ice pack and your skin for comfort.

Drink plenty of fluids. Avoid hot liquids or foods. Soft foods and liquids can be eaten on the day of surgery. You may return to a normal diet 1–2 days after surgery unless otherwise directed.

Warm saltwater rinses can be made by mixing ½ teaspoon of salt in a cup of warm water and should be used at least 4–5 times a day, especially after meals. Alternatively, you may use Peridex™ mouth rinse as directed.

Brushing your teeth and the healing abutments is no problem. Be gentle initially with brushing the surgical areas.

Keep physical activities to a minimum for several days following surgery. Avoid bending over, heavy lifting, or strain. Keep in mind that you are probably not taking normal nourishment. This may weaken you and further limit your ability to exercise.