



The Royal College of Emergency Medicine

Patron: HRH The Princess Royal

Glossary of terms used in College examinations

The Royal College of Emergency Medicine uses several terms in examinations that may cause confusion. The following definitions are intended as a guide to the understanding of these terms. It is important to read the questions carefully and to understand the term in the context of that question. Examiners and candidates are advised to be rigorous in the use of these terms.

Abnormality

This is any feature in an examination or investigation which is outside the standard deviation of the population being studied. A **Clinical** abnormality however would be a pathologically relevant abnormality and would not include the presence of tubes, prostheses etc.

Assessment

History taking, physical examination and use of investigations.

Characteristics

Something that describes a condition, or piece of equipment that is consistently present in that condition or is pretty fundamental to how the piece of equipment works e.g. mercury is characteristic of the content of thermometers

Class of drug

This is the generic name for the type of drug with a particular pharmacological affect e.g. anticoagulant, antihypertensive etc.

Clinical findings

This may include symptoms, signs and vital signs. It is information gleaned from the clinical evaluation, but not the results of investigations even bedside ones (e.g. BM or Urine Dipstick)

Commonest/Common

>75% incidence, or prevalence

Condition

This would suggest a well know pathological entity or diagnosis that should be mentioned as contributing to the presenting complaint.

Criteria

This refers to the fact that there is a formal international/national guideline or scoring system that allows you to define the seriousness of a condition e.g. CURB-65 score for pneumonia etc. Each criterion may be a clinical sign, measurement, or bedside observation that helps discriminate in some way for the management of the patient.

Definitive management/treatment

This may include things you would do in the department but usually requires you to list the operation or procedure that will cure or contain the condition.

This may also refer to the gold standard treatment which has been proven to give best results, even if not available in the institution where you work.

Disposition

Where the patient is sent following care in the ED including follow-up if discharged.

ED management

This requires you to list actions that are life or limb saving or that might improve the course of the condition if done within the ED. It is not definitive management. This may however include analgesia, referral to specialty team etc

Essential

This indicates life saving treatments/management steps that are the priority, and would not normally include things like analgesia, communication etc.

Factor

A contributing element or cause for the condition.

Features

This is used in a variety of ways

In the medical history – it indicated symptoms

In the examination – examination findings

In results – abnormalities that are clinically relevant or might simply be the presence of an ETT or central line i.e. abnormality

If describing equipment or procedures, it is how the equipment looks, or key elements of the procedure.

Clinical features can be symptoms or signs

Immediate

This indicates what you will do now, rather than include within the general list of investigations or treatments that a patient needs.

Implication

Something that is suggested or hinted at.

Important

Used to indicate something that needs treatment or has a very high chance of recurring e.g. important complications are those that you warn patients about, or that you specifically wish to exclude if a patient deteriorates

Indicators

This is used in the context of a clinical evaluation. It should include history, examination and investigations that might indicate that a particular diagnosis is likely.

Investigations

Specific tests undertaken to make a diagnosis or monitor the patient's condition. They may include bedside tests such as urine dipstick or BM unless otherwise specified.

Management

Aspects of care including treatment, supportive care and disposition/disposal. This does not normally include investigations unless an investigation leads to an immediate change in the treatment, i.e. blood gas to check the correct Oxygen level is being given.

Measures

Actions that can be taken which may include physical procedures, prescriptions, referrals etc.

Most likely

This requires the commonest or best know item. For example, if asked for the most likely organisms causing a UTI – you should list E Coli.

Pathophysiological sequence of events

This requires you to list in time order, the events that happen on a cellular, or hormonal level, leading to the current condition. For example, if a lactate is high in the presence of sepsis, you could suggest –

- Hypotension
- Poor organ perfusion
- Tissue hypoxia
- Anaerobic metabolism
- Glycolysis and lactate build up

Pathognomic

Refers to a symptom or sign that if present, would always lead to a particular diagnosis

Pre-alert or Standby

These mean the same thing i.e. a radio call from an ambulance crew to inform ED of an acutely unwell/injured patient arriving imminently.

Principles

These are the ideal or essential themes of a treatment or plan. e.g. Principles of drug treatment do not usually require doses or routes but might include “broad spectrum antibiotics” or “antihistamines”

Rarely

<10% of the time

Recommended

This is the best treatment according to a National guideline or accepted practice

Symptoms

This is what the patient complains of.

Signs

This is what you identify by examination and may include abnormal observations/measurements of vital parameters.

Steps in a management plan

Actions that may include giving treatment, support or referring, if it included an investigation, the investigation must lead to a change in the management plan.

Strategy

This is your plan of action, and would normally include a list of investigations, prescriptions, physical treatments, in a particular order.

Treatment

Measures undertaken to cure or stabilise the patient's condition. This includes oxygen, fluids, drugs, and may also mean surgery. It does not include investigations.

Usual/normal

>90% of the time

Abbreviations that may be used in the examinations

ACP	Advance Care Practitioner
ACS	Acute Coronary Syndrome
AECU	Ambulatory Emergency Care Unit
AF	Atrial Fibrillation
AKI	Acute Kidney Injury
ALS	Advanced Life Support
ALTE	Apparent Life-Threatening Event
ARCP	Annual Review of Competency Progression
ARDS	Acute Respiratory Distress Syndrome
ATLS	Advanced Trauma Life Support
AXR	Abdominal x-ray
BHCG	Beta Human Chorionic Gonadotropin
BiPAP	Bilateral Positive Airway Pressure
BLS	Basic Life Support
BM	blood sugar reading
BMI	Body Mass Index
BNF	British National Formulary
BP	Blood Pressure
BRUE	Brief resolved unexplained event
CAMHS	Child and adolescent mental health service
C-spine	Cervical Spine
CES	Cauda Kidney Disease
CN	Cranial Nerve
COPD	Chronic Obstructive Pulmonary Disease
COVID-19	Coronavirus 19
CPAP	Continuous Positive Airway Pressure
CRP	C-Reactive Protein
CSF	Cerebrospinal Fluid
CT 1/2/3	Core Trainee years 1-3 of training
CT scan	Computerised Tomography Scan
CV	Curriculum Vitae
CXR	Chest x-ray
DATIX	Internet based incident reporting system
DGH	District General Hospital
DIC	Disseminated Intravascular Coagulation

DNACPR	Do Not Attempt Cardiopulmonary Resuscitation
DOAC	Direct oral anticoagulant
DVLA	Driver and Vehicle Licensing Agency
DVT	Deep Vein Thrombosis
ECG	Electrocardiogram
ECHO	Echocardiogram
ECMO	Extracorporeal Membrane Oxygenation
ED	Emergency Department
eGFR	Estimated glomerular filtration rate
ENP	Emergency Nurse Practitioner
ENT	Ear Nose and Throat
ESR	Erythrocyte Sedimentation Rate
FAST	Focussed Assessment with Sonography for Trauma
FBC	Full Blood Count
FY1/2	Foundation Year doctor in year 1 or 2 of their foundation training
GCS	Glasgow Coma Score
GMC	General Medical Council
GORD	Gastro-oesophageal reflux disease
GP	General Practitioner
GTN	Glyceryl trinitrate
Hb	Haemoglobin
HELLP	Hypertension, Elevated Liver Enzymes, Low Platelets
HIV	Human Immunodeficiency Virus
HR	Heart Rate
ICD	Implantable cardiac defibrillator
ICP	Intracranial Pressure
ICU	Intensive Care Unit
IDDM	Insulin Dependent Diabetes Mellitus
IHD	Ischaemic Heart Disease
INR	International Normalised Ratio
IV	Intravenous
IVDU	Intravenous Drug User
JVP	Jugular Venous Pressure
LVH	Left Ventricular Hypertrophy
LVF	Left Ventricular Failure
LP	Lumbar Puncture

LFT	Liver Function Tests
LMN	Lower Motor Neurone
LMP	Last menstrual period
LRTI	Lower Respiratory Tract Infection
mcg	Microgram
mg	Milligram
Mg/dl	Milligram/deciliter
MI	Myocardial Infarction
mL	Millilitre
MR	Magnetic Resonance
MRI	Magnetic Resonance Imaging
MRSA	Methicillin Resistant Staphylococcus Aureus
NAI	Non-accidental Injury
NICE	National Institute for Health and Clinical Excellence
NIDDM	Non-insulin dependent diabetes mellitus
NIV	Non-Invasive Ventilation
NOAC	Novel oral anticoagulant
NSAID	Non-Steroidal Anti-Inflammatory Drug
NSTEMI	Non-ST elevation myocardial infarction
OGD	Oesophago-gastro duodenoscopy
OPG	Orthopantomogram
OSCE	Objective Structured Clinical Examination
PALS	Patient Advice and Liaison Service
PCI	Percutaneous coronary intervention
PEA	Pulseless Electrical Activity
PEEP	Positive End Expiratory Pressure
PEFR	Peak Expiratory Flow Rate
PEP	Post exposure prophylaxis
PPE	Personal Protective Equipment
QI	Quality improvement
QIP	Quality Improvement Project
RR	Respiratory Rate
RCEM	Royal College of Emergency Medicine
ROSC	Return of Spontaneous Circulation
RTC	Road Traffic Collision
SIGN	Scottish Intercollegiate Guidelines Network

SpO₂	Oxygen Saturations
STEMI	ST elevation myocardial infarction
SUFE	Slipped Upper Femoral Epiphysis
SVT	Supraventricular tachycardia
TELP	Treatment escalation/limitation plan
TEMP	Temperature
TFT	Thyroid Function Tests
TIA	Transient Ischaemic Attack
TPD	Training Programme Director
U&E's	Urea & Electrolytes
UMN	Upper Motor Neurone
URTI	Upper Respiratory Tract Infection
USS	Ultrasound Scan
UTI	Urinary Tract Infection
VF	Ventricular Fibrillation
VT	Ventricular Tachycardia
WBPA	Workplace base assessment
WCC	White cell count

Normal Values

Haematology

Haemoglobin	11.5 - 16.6g/dl
White blood cells	4 - 11 x 10 ⁹ /L
Platelets	150 - 450 10 ⁹ /L
MCV	80 - 96 fl
MCHC	32 - 36 g/dl
Neutrophils	2 - 7.5 x 10 ⁹ /L
Lymphocytes	1.5 - 4 x 10 ⁹ /L
Monocytes	0.3 - 1 x 10 ⁹ /L
Eosinophils	0.1- 0.5 x 10 ⁹ /L
Basophils	<0.2 x 10 ⁹ /L
Reticulocytes	<2 %
Haematocrit	0.35 - 0.49
Red Cell distribution width	11 - 15%

Biochemistry

Sodium	135 - 145 mmol/L
Potassium	3 - 4.5 mmol/L
Urea	2.5 - 7.5 mmol/L
Glucose	3.5 - 5 mmol/L
Creatinine	35 - 135 µmol/L
Alanine aminotransferase	5 - 35 U/L
Gamma GT	<65 U/L
Alkaline phosphatase	30 - 135 U/L
AST	<40 U/L
Total Protein	60 – 80 g/l
Albumin	35 - 50 g/L
Globulin	2.3 - 3.5 g/dl
Amylase	<70 U/L
Total bilirubin	3 - 17 µmol/L
Calcium	2.1 - 2.5 mmol/L
Chloride	95 – 105 mmol/L
Phosphate	0.8 - 1.4 mmol/L

Blood gases

pH	7.35 - 7.45
pO ₂	11 - 14 KPa
PCO ₂	4.5 - 6 KPa
Base excess	-2 to +2 mmol/L
Bicarbonate	24 - 30
Lactate	<2 mmol/L